

Antioxidant Properties and Phenolic Composition of *Salvia virgata* Jacq.

Gökçe Şeker KARATOPRAK^{1,*}, Selen ILGÜN², Müberra KOŞAR¹

¹Erciyes University, Faculty of Pharmacy, Department of Pharmacognosy, 38039 Melikgazi, Kayseri, TURKEY, ²Erciyes University, Faculty of Pharmacy, Department of Pharmaceutical Botany, 38039 Melikgazi, Kayseri, TURKEY

Methanol and water extracts of *Salvia virgata* Jacq. aerial parts were examined for its phenolic contents and antioxidant activities. Both extracts were investigated for their total phenolic contents, total flavonoids and flavonols, qualitative-quantitative compositions (by HPLC-PDA analysis). Free radical scavenging activities (using DPPH• and ABTS^{•+}), iron(II) chelation activity and also the peroxidation level was determined by TBA method. The antioxidant activity of extracts were given as IC₅₀ values and compared with standards; butylated hydroxyanisole, butylated hydroxytoluene, rosmarinic acid, ascorbic acid, and gallic acid. Methanol extract was found to be active for DPPH• radical scavenging activity where as water extract was more active for ABTS^{•+} radical scavenging. The methanol extract enriched in phenolic compound and rosmarinic acid as the main component in both extracts has been identified.

Key words: *Salvia virgata*, Antioxidant activity, HPLC

Salvia virgata Jacq.'ın Antioksidan Özellikleri ve Fenolik İçeriği

Salvia virgata Jacq.'ın toprak üstü kısımlarının metanol ve su ekstresinin fenolik içeriği ve antioksidan özellikleri belirlenmiştir. Her iki ekstrenin de toplam fenolik, toplam flavonoit ve flavonol içeriği, kalitatif-kantitatif kompozisyonları (YPSK ile), serbest radikal süpürücü aktiviteleri (DPPH• ve ABTS^{•+}), demir(II) ile şelat oluşumu ve TBA yöntemi ile peroksidasyon seviyeleri incelenmiştir. IC₅₀ değeri olarak verilen antioksidan aktiviteler butil hidroksianisol, butil hidroksitoluen, rozmarinik asit, askorbik asit ve gallik asit standartları ile karşılaştırılmıştır. Metanol ekstresi DPPH• radikalini süpürücü etkide aktif bulunurken, su ekstresi ABTS^{•+} radikalini süpürücü etkide aktif bulunmuştur. Metanol ekstresinin fenolik bileşiminin daha zengin olduğu ve her iki ekstrede de rozmarinik asitin ana bileşen olduğu tespit edilmiştir.

Anahtar kelimeler: *Salvia virgata*, Antioksidan aktivite, YPSK

*Correspondence: E-mail: gskaratoprak@gmail.com; Tel: +903524380486/28176

INTRODUCTION

Numerous lines of evidence have indicated that free radicals play a critical role in a variety of pathological conditions such as brain dysfunction, cancer, heart diseases and inflammation by damaging cellular components of DNA, proteins and lipids. Therefore antioxidants present in human diet

may play an important role in disease prevention (1-3). Recently intense addition has been devoted to natural sources of phenolic antioxidants, not only for the prevention and treatment of diseases but also for protecting food quality by preventing oxidative deterioration of lipids (4, 5). Synthetic antioxidants such as butylated hydroxyanisole (BHA), butylated

hydroxytoluen (BHT) and tert-butyl hydroquinone have been used to protect food products against lipid peroxidation. Many researchers have focused on the search for natural compounds with antioxidant properties because of the harmful effects of synthetic antioxidants in the human body (6, 7).

The genus *Salvia* with about 900 species, is one of the most widespread members of the *Lamiaceae* family (2, 8). *Salvia* have been used since ancient times in folk medicine and have been subjected to extensive researches and it features prominently in the pharmacopoeias of many countries throughout the world (7-9). *Salvia officinalis* serves as standard sage is well known with its antioxidant activity and used as by this purpose. Hydroxycinnamic acid derivatives (caffeic and rosmarinic acids), flavonoids (luteolin, apigenin and glycosides) and diterpenoids (carnosol, carnosic acid and methyl carnosol) are responsible by the antioxidant effects of *Salvia* species (10).

Salvia genus is represented by totally 95 species in Turkey (11). In Anatolian folk medicine the infusions of *Salvia* species generally have been used as carminative, antiperspirant, diuretic and for wound healing (12). In Turkey, *S. virgata* Jacq. is known as 'yılancık' and its leaves used for externally wound healing. The decoction of *S. virgata* is also used for leukemia (13). The objective of this work was to evaluate the antioxidant activity of *S. virgata*. Therefore water extract and methanol extract were investigated in different *in vitro* antioxidants test systems. Furthermore the total phenols, total flavonoids, total flavonols were also analyzed. The phenolic compositions of the extracts were identified by HPLC-PDA analysis.

EXPERIMENTAL

Plant material and reagents

Air-dried aerial plant material (*S. virgata* Jacq., *Lamiaceae*) was collected from University of Uludağ /Bursa on 22/06/2009. A voucher specimen has been deposited at the Uludağ University Herbarium of Faculty of Arts and Sciences (BULU 29050). Chromatographic standards *p*-OH benzoic acid (65-85-0), caffeic acid (331-39-5), *o*-

coumaric acid (614-60-8), rosmarinic acid (20283-92-5), luteolin (491-70-3) were purchased from Sigma Chemical Company (St. Louis, MO, USA). Ultra-pure water was used throughout and was prepared using a Millipore Milli-RO 12 plus system (Millipore Corp., Billerica, MA, USA). All remaining reagents were of the highest purity available and obtained from the Sigma Chemical Company (St. Louis, MO, USA).

Preparation of the extracts

Air-dried *S. virgata* was cut into small pieces and sequentially extracted with sufficient amount of 70% methanol and water for 24 hrs at 40 °C in a water bath with shaking and was filtered. This procedure was repeated three times using the same batch of starting material and the resultant filtrates were combined and the solvent was removed in vacuo (40 °C). All extracts stored at -20 °C until the analysis time. Prior to the analysis, an aliquot of each extract was dissolved and filtered through 0.45µm membrane (Whatman, Maidstone, UK) and used in all subsequent experiments.

Determination of total phenolics, flavonoids, and flavonols

Total phenols were estimated as gallic acid equivalents (GAE), per gram of extract (14). To ca. 6.0 mL H₂O, 100 µL sample (conc. 2-6 mg/mL) were transferred into a 10.0 mL volumetric flask, to which 500 µL undiluted Folin-Ciocalteu reagent were subsequently added. After 1 min, 1.5 mL 20% (w/v) Na₂CO₃ were added and the volume was made up to 10.0 mL with H₂O. After 2 hrs incubation at 25 °C, the absorbance was measured at 760 nm and compared to a gallic acid calibration curve. The data are presented as the mean value of triplicate analyses.

Total flavonoids and total flavonols were estimated as rutin equivalents (RE), expressed as mg rutin eq./g extract (15). One milliliter of plant extract in methanol (2.5 mg/mL) was mixed with 1 mL aluminum trichloride in ethanol (20 g/L) and diluted with ethanol to 25 mL. The absorbance at 415 nm was read after 40 min at 20 °C. Blank samples were prepared from 1 mL plant extract (conc. 2.5 mg/mL) and 1 drop of acetic acid, and diluted to 25 mL. The rutin calibration curve was

prepared in ethanolic solutions using the same procedure. All determinations were carried out in quadruplicate and the mean values were calculated.

For total flavonols, the rutin calibration curve was prepared by mixing 2 mL of 0.5–0.015 mg/mL rutin in ethanol with 2 mL (20 g/L) aluminum trichloride and 6 mL (50 g/L) sodium acetate. The absorbance at 440 nm was read after 2.5 hrs at 20 °C. The same procedure was carried out with 2 mL of plant extract (2.5 mg/mL) instead of the rutin solution. All determinations were carried out in quadruplicate and the mean values were calculated.

Qualitative–quantitative chromatographic analysis

The liquid chromatographic apparatus (Shimadzu LC 10Avp, Kyoto, Japan) consisted of an in-line degasser, pump and controller coupled to a SPD-M10 Avp photodiode array detector equipped with an automatic injector interfaced to Class VP chromatography manager software (Shimadzu, Kyoto, Japan). Separations were performed on a 250 x 4.6 mm i.d., 5 µm particle size, reverse-phase Discovery-C18 analytical column (Supelco, Bellefonte, PA, USA) operating at room temperature (22 °C) at a flow rate of 1 mL/min. Detection was carried out between the wavelengths of 200 and 550 nm. Elution was carried out using a ternary non-linear gradient of the solvent mixture MeOH/H₂O/CH₃COOH (10:88:2,

v/v/v) (solvent A), MeOH/H₂O/CH₃COOH (90:8:2, v/v/v) (solvent B) and MeOH (solvent C).

Components were identified by comparison of their retention times to those of authentic standards under identical analysis conditions and UV spectra using our in house PDA-library. A 10 min equilibrium time was allowed between injections. All the standard and sample solutions were injected triplicate.

1,1-Diphenyl-2-picrylhydrazyl radical (DPPH•) scavenging activity

The ability of the extracts to scavenge DPPH• was determined by the method of Gyamfi et al. (1999) (16). A 50 µL aliquot of each extract, in 50 mM Tris-HCl buffer (pH 7.4), was mixed with 450 µL of Tris-HCl buffer and 1.0 mL of 0.1 mM DPPH• in MeOH. After 30 min incubation in darkness and at ambient temperature, the resultant absorbance was recorded at 517 nm. The percentage inhibition was calculated using eq. 1. Estimated IC₅₀ values are presented as the mean value of triplicate analyses.

$$\% \text{ inhibition} = \frac{[(\text{Abs}_{\text{control}} - \text{Abs}_{\text{sample}}) / \text{Abs}_{\text{control}}] \times 100}{\text{Abs}_{\text{control}}} \quad (\text{Eq. 1})$$

ABTS•+ radical scavenging activity

The determination of ABTS•+ radical scavenging was carried out as described by Re et al. (17). The ABTS•+ radical was generated by reacting an (7 mmol/L) ABTS•+ aqueous solution with K₂S₂O₈ (2.45 mmol/L, final concentration) in the dark for 12–16 h, at

Table 1. Gradient flow of HPLC

Time(min)	Solvent A (%)	Solvent B (%)	Solvent C (%)
0	85	15	0
15	70	30	0
18	60	40	0
30	60	40	0
35	100	0	0
37	85	0	15
48	70	0	30
50	85	15	0

Table 2. Extract yields, total phenols, flavonoids and flavonols

Extracts*	Yield ^a (%)	Total Phenols ^b (mg _{GAE} /g _{extract})	Total Flavonoids ^c (mg _{RE} /g _{extract})	Total Flavonols ^d (mg _{RE} /g _{extract})
SME	25.94	195.22 ± 0.25	62.20 ± 0.57	15.24 ± 1.12
SWE	24.99	120.14 ± 2.27	14.17 ± 0.83	9.62 ± 0.60

^a Extract yields expressed as milligrams of extract per gram (dry weight) of root material. ^bTotal phenolics expressed as mg gallic acid per gram (dry weight) of extract. ^c Total flavonoids expressed as mgrutin per gram (dry weight) of extract. ^d Total flavonols expressed as mgrutin per gram (dry weight) of extract. SME 70% methanol extract; SWE water extract.

ambient temperature, and adjusting the Abs_{734nm} to 0.700 (70.020) with ethanol. The samples were diluted, such that, a 15 mL sample when added to 1.485 mL ABTS^{•+} resulted in a 20–80% inhibition of the blank absorbance. After 1.485 mL ABTS^{•+} solution was added to 15 mL sample, the absorbance at 734 nm was recorded 1 min after initial mixing and subsequently (for 15 min in total). The results are expressed as the Trolox equivalent antioxidant capacity (TEAC, mmol/L Trolox) at 1, 5, 10 and 15 min times and as the relative antioxidant activity (AUC), as described by Re et al. (17).

Iron(II) chelation

The ability of the extract to chelate iron(II) was carried out as described by Carter (1971) (18). To 200 µL of extract dissolved in an appropriate solvent was added 100 µL (2.0 mmol/L) FeCl₂·4H₂O and 900 µL methanol. After 5 min incubation, the reaction was initiated by the addition of 400 µL (5.0 mmol/L) ferrozine. After a further 10 min incubation period, the absorbance at 562 nm was recorded. EDTA was used as a positive control. The Fe²⁺ chelating activity was calculated using eq. 1. The values are presented as the mean of three measurements.

Ascorbate-Iron(III)-catalyzed phospholipid peroxidation

The ability of the extracts to scavenge hydroxyl radicals was determined by the method of Aruoma et al. (19). Bovine brain extract (Folch type VII) was mixed with 10 mM phosphate-buffered saline (PBS, pH 7.4) and sonicated in an ice bath until an

opalescent suspension was obtained, containing 5 mg/mL phospholipid liposomes. The liposomes (0.2 mL) were combined with 0.5 mL of PBS buffer, 0.1 mL of 1 mM FeCl₃, and 0.1 mL of extract dissolved in PBS. Peroxidation was initiated by adding 0.1 mL of 1 mM ascorbic acid. The mixture was incubated at 37 °C for 60 min, after which, 50 µL of 2.0% butylated hydroxytoluene (BHT) was added to each tube, followed by 1 mL of 2.8% trichloroacetic acid (TCA) and 1 mL of 1.0% 2-thiobarbituric acid (TBA) in 0.05 M NaOH. The samples were vortexed and heated in a water bath at 100 °C for 20 min. The reaction was stopped by a 5 min ice H₂O bath. To each tube was added 2 mL of *n*-butanol, and the mixture was vigorously vortexed. After centrifugation, the extent of oxidation was estimated from the absorbance of the organic layer at 532 nm. The percentage inhibition was calculated using eq 1, where the controls contain all the reaction reagents except the extract or positive control substance, and the IC₅₀ values were estimated using a nonlinear regression algorithm (Sigma Plot 2001 version 7.0). The values are presented as the mean values of five measurements.

Statistical analysis

Data are presented as mean values ± 95% confidence interval. Analysis of variance was performed using ANOVA procedures. Significant differences between means were determined by Tukey's pairwise comparison test at a level of $p < 0.05$. IC₅₀ values were estimated using a nonlinear regression algorithm unless stated otherwise.

RESULTS

Extract yields, total phenolic, flavonoid and flavonol contents

S. virgata herb was extracted with 70% methanol and water. The results of extract yields, total phenols, total flavonoids and total flavonols are presented in Table 2. Aqueous methanol extract contained the highest amount of total phenol, flavonoids, flavonols. According to the Table 2, the yields of both extracts were also found to be almost same.

Qualitative–quantitative chromatographic analysis

The qualitative–quantitative analyses of the extracts, carried out using an HPLC apparatus coupled to a PDA detector, are presented in Table 3. Phenolic compounds were identified and quantified at 280, 320, and 360 nm as benzoates, hydroxycinnamates, and flavonoids, respectively. Gallic, *p*-OH-benzoic, caffeic, *o*-coumaric, rosmarinic acids, luteolin-7-*O*-glycoside, and luteolin were identified by comparison with the retention times and UV spectra of authentic standards, while quantitative data were calculated from their calibration curves. For *S. virgata*, the aqueous methanol extract was found to contain phenolics in high amount as measured by both UV spectrophotometry and HPLC. Rosmarinic acid was the main compound in both extracts. The rosmarinic acid concentration of methanolic extract was found as 66.94 ± 0.47 mg/g (w/w) and water extract 26.81 ± 0.05 mg/g (w/w). *o*-Coumaric acid and caffeic acid were found in the *S.*

virgata extracts as the other hydroxycinnamic acids. Also it was found that the extracts contain more flavonoid glycosides than aglycones. The luteolin concentrations of the extracts were 0.97 ± 0.03 mg/g for methanolic extract and 0.22 ± 0.00 mg/g for water extract. As well known from the literature, rosmarinic acid was found as the principal component (5, 20-25).

1,1-Diphenyl-2-picrylhydrazyl radical (DPPH•) scavenging activity

1,1-Diphenyl-2-picrylhydrazyl (DPPH•) is a kind of stable organic radical. The DPPH• oxidative assay is used worldwide in the quantification of radical-scavenging capacity. The DPPH• alcohol solution is deep purple and possesses a characteristic absorption maximum between 515-517 nm. When reduced to hydrazine derivative by an antioxidant *via* electron or hydrogen atom transfer reactions, this absorption maximum decreases (25-27). The reactive rate and the ability of the antioxidant depend on the rate and the peak value of disappearance of the DPPH•. All sage extracts managed to scavenge the DPPH• radicals at physiological pH and did so in a concentration-dependent fashion. IC₅₀ values defined as the concentration required to scavenge 50% of the available free radicals, estimated by non linear regression for all the extracts are presented in Figure 2. According to Figure 2, the aqueous methanol extract was more active than (IC₅₀: 0.16 mg/mL) the water extract (IC₅₀: 0.20 mg/mL). Also aqueous methanol extract was as active as ascorbic acid. The IC₅₀ value of

Table 3. HPLC Results of *S. virgata* extracts

Compounds	Equation and r^2 of standards	Extracts (mg/g)	
		SWE mean \pm SD (n=3)	SME mean \pm SD (n=3)
<i>p</i> -OH-benzoic acid	$y=13918x + 7,95 \quad r^2=0,998$	0.83 ± 0.013	0.69 ± 0.040
caffeic acid	$y=47251x+191,94 \quad r^2=0,996$	0.77 ± 0.001	0.97 ± 0.004
<i>o</i> -coumaric acid	$y=25443x -12,69 \quad r^2=0,998$	0.90 ± 0.002	3.25 ± 0.240
rosmarinic acid	$y=23903x -29,76 \quad r^2=0,998$	26.81 ± 0.047	66.94 ± 0.471
luteolin	$y=42001x -84,34 \quad r^2=0,997$	0.22 ± 0.003	0.97 ± 0.031

SWE: water extracts; SME: 70% methanol extracts




Figure 1. HPLC chromatogram of *Salvia virgata* extracts and standards.

A; Standards, B; Methanol extract, C; Water extract, 1; *p*-OH-benzoic acid, 2; caffeic acid, 3; *o*-coumaric acid, 4; rosmarinic acid, 5; luteolin

rosmarinic acid was found 0.20 mg/mL. The radical scavenging of rosmarinic acid was higher than ascorbic acid, BHT and BHA. It is estimated that the concentration of rosmarinic acid as main component is related to DPPH[•] radical scavenging activity. The strong superoxide scavenging activities of the

rosmarinic acid and its derivatives like their DPPH[•] radical scavenging activities may also be attributed to the presence of the caffeic acid moiety which is contained in sage fractions (28).

ABTS^{•+} radical scavenging

The DPPH radical is frequently used to determine the free radical scavenging activity of natural products (29). Occasionally, however, it is inappropriate to use this particular free radical due to sample solubility or spectral interference (30). An alternative synthetic radical is ABTS^{•+} a moderately stable nitrogen-centered radical species. Although the principle underpinning the use of ABTS^{•+} and DPPH[•] free radicals is essentially identical, ABTS^{•+} based models are more versatile as both non polar and polar samples can be assessed and spectral interference is minimized as the absorption maximum used is 760 nm, a wavelength not normally encountered with natural products (31). The extracts analyzed at two different concentrations and the percentage inhibition was plotted as function of concentration and the total antioxidant activity was calculated using Trolox calibration curve. As can be seen in Figure 3, all extracts and positive controls manage to inhibit the ABTS^{•+} radical in a concentration-dependent fashion. Also it is found that aqueous methanol extract and water extract were more active than ascorbic acid. Water extract inhibit the ABTS^{•+} radical both two concentrations (0.25 mg/mL and 0.5 mg/mL). Rosmarinic acid (TEAC 1.23-2.26 mg/mL) was found to be more active than BHT (TEAC 0.57-1.17 mg/mL).

Iron(II) chelation activity

Foods are often contaminated with transition metal ions which may be introduced by processing methods. Bivalent transition metal ions play an important role as catalysts of oxidative processes, leading to the formation of hydroxyl radicals and hydroperoxide decomposition reactions *via* Fenton chemistry processes can be delayed by iron chelation and deactivation (32). Therefore, the ability of the extracts to chelate iron (II) ions was evaluated and expressed as Na₂EDTA equivalents ($\text{mg}_{\text{Na}_2\text{EDTA}}/\text{g}_{\text{extract}}$). The results are presented in Figure 4. The aqueous methanol extract and water extract showed an activity on Iron (II) chelation. Flavonoids that are presented in these extracts are responsible for this activity. The structural characteristics of flavonoids are efficient on binding the Iron (II) (17). None of the extracts were not as active as EDTA.

Ascorbate-Iron(III)-catalyzed phospholipid peroxidation

Polyunsaturated fatty acid rich phospholipids are important biologically relevant substrates and are susceptible to hydroxyl radical-mediated degradation. Furthermore, it is considered that the oxidative deterioration of lipids is the principal cause of the development of rancidity in lipid-rich products (33, 34).




Figure 2. Effect of the extracts of *S. virgata* and positive controls on DPPH[•] radical scavenging.

AscAs, ascorbic acid; BHT, Butylated hydroxytoluene; BHA, Butylated hydroxyanisole; GA, gallic acid; RA, rosmarinic acid; SME 70% methanol extract; SWE water extract. Values are presented as means \pm standard error. Bars with the same letter (a-e) are not significantly ($p < 0.05$) different.

Therefore, it was considered important to assess the ability of the *S.virgata* extracts to protect phospholipids against hydroxyl radical mediated oxidative degradation. In the ascorbate-iron(III)-catalyzed hydroxyl radical-mediated degradation of bovine brain-derived phospholipid liposomes, a sample capable of inhibiting the formation of TBARS can be described as an antioxidant capable of protecting phospholipids at physiological pH from oxidative degradation via a mechanism of hydroxyl radical scavenging. As seen in Figure 5, none of the *S.virgata* extracts were not as active as positive controls. Aqueous

methanol extract was found to be more active than water extract.

DISCUSSION

As well known from the literature, rosmarinic acid was found as the principal component. Also rosmarinic acid, carnosic acid, carnosol and their derivatives in sage species are main components and these compounds are responsible for their antioxidant activities (5, 20, 21, 23, 24, 35, 36).




Figure 3. Effect of the extracts of *S.virgata* and positive controls on ABTS^{•+} radical scavenging.

AscAs, ascorbic acid; BHT, Butylated hydroxytoluene; BHA, Butylated hydroxyanisole; GA, gallic acid; RA, rosmarinic acid; SME 70% methanol extract; SWE water extract. Values are presented as means \pm standard error. Bars with the same letter (a-g) and number (1-6) are not significantly ($p < 0.05$) different.




Figure 4. Effect of the extracts of *S.virgata* and positive control on Iron (II) chelation.

EDTA (ethylenediaminetetraaceticacid); SME 70% methanol extract; SWE water extract. Data are presented as means ± standard error.




Figure 5. Effect of the extracts of *S.virgata* and positive controls on Hydroxyl radical-mediated phospholipid degradation.

BHT, butylated hydroxytoluene; BHA, butylated hydroxyanilin; GA, gallic acid; RA, rosmarinic acid; SME 70% methanol extract; SWE water extract. Data are presented as means ± standard error. Bars with the same letter (a-c) are not significantly ($p < 0.05$) different.

In current study the aqueous methanol extract is determined to have the higher phenolic composition and higher rosmarinic acid level. Therefore it's found to be the most effective extract. Evaluation of relationship between antioxidant activity of plant products and their phenolic content has been discussed in several studies. Some authors found a

correlation while others found no relationship (35) between total phenolic content and antioxidant activity. Göger et al. also demonstrated that the 50% methanol extract of *S.virgata* was found to have significantly higher activity than the other extracts on reducing the iron (III) to iron (II) (12). Phenolic acids and flavonoids are dissolved in

polar solvents so they show strong activity in the polar systems like, iron(III) to iron(II) reduction activity and DPPH[•] scavenging activity. Therefore the fractions which are rich by the phenolic acids and flavonoids found to be efficient in these assays.

It is estimated that the concentration of main component rosmarinic acid is related to the scavenging activity against DPPH[•] radical. The strong superoxide scavenging activities of the rosmarinic acid and its derivatives may also be attributed to the presence of the caffeic acid moiety like their DPPH[•] radical-scavenging activities (28). The results showed that phenolic acids (rosmarinic acid derivatives, sagecoumarin, sagerinic acid, salvianolic acid) has potent activity while the flavonoid glycosides (luteolin and apigenin glycosides) all display weak to moderate activities, though they are still better than trolox (28). Also Kosar et al. reported that carnosic acid and rosmarinic acid were found to be the principal radical scavengers in sage by the HPLC- post column DPPH[•] method (5).

The ABTS assay is based on the generation of a blue/green ABTS^{•+}, which is applicable to both hydrophilic and lipophilic antioxidant systems; whereas DPPH assay uses a radical dissolved in organic media and is, therefore, applicable to hydrophobic systems (36). The difference between antioxidant capacities of the extracts determined by the two assays is agreed with the principles of the assays.

Plant extracts are rich in phenolic compounds and therefore these phenolics complex with transition metal ions, rendering them unable to participate in metal- catalyzed initiation (37). From the represented data *S.virgata* extracts may be able to afford protection against oxidative damage by complexing free iron(II) ions.

Free radical mediated degradation of phospholipids is resulted with cell death and also considered to be responsible for the oxidative deterioration and off flavor development of foods. Non polar antioxidants are concentrated at the lipid-air interface and demonstrate high protection in emulsions against the polar antioxidants presented in aqueous phase (28, 30) used the same method for *Sideritis* extracts and compared this method with some other antioxidant tests.

According to the data presented the extracts were able to protect phospholipids by scavenging hydroxyl radicals before the reaction of susceptible components within the lipid matrix. Therefore the extracts especially the methanol extract have a potent protection ability of biological membrane from the actions of reactive hydroxyl radical.

CONCLUSION

Overall, the data suggest that the *S.virgata* used in this study possess promising antioxidant properties *in vitro*. This plants also can be used to developing ingredient for the new antioxidant agents for different materials such as cosmetics and dietary foods. Further work should be carried out to determine whether extracts possess *in vivo* activities; otherwise, their potential use in the functionalization of cosmetics and foods cannot be confidently assumed.

REFERENCES

1. Generalic I, Skroza D, Ljubenkovic I, Katalinic A, Burcul F, Katalinic V, Influence of the phenophase on the phenolic profile and antioxidant properties of Dalmatian sage, Food Chem 127, 427-433, 2012.
2. Kosar M, Goger F, Baser KHC, *In vitro* antioxidant properties and phenolic composition of *Salvia halophila* Hedge. from Turkey, Food Chem 129, 374-379, 2011.
3. Latte KP, Kolodziej H, Antioxidant properties of phenolic compounds from *Pelargonium reniforme*, J Agric Food Chem 52, 4899-4902, 2004.
4. Matkowski A, Zielin'ska S, Oszmian'ski J, Lamer-Zarawska E, Antioxidant activity of extracts from leaves and roots of *Salvia miltiorrhiza* Bunge., *S. przewalskii* Maxim., and *S. verticillata* L., Bioresource Technol 99, 7892-7896, 2008.
5. Kosar M, Dorman HJD, Baser KHC, Hiltunen R, An Improved HPLC post-column methodology for the identification of free radical scavenging phytochemicals in complex mixtures, Chromatographia 60, 11-12, 2004.
6. Kosar M, Dorman HJD, Baser KHC, Hiltunen R, Screening of free radical scavenging compounds in water extracts of *mentha* samples using a postcolumn derivatization method, J Agric Food Chem 52, 5004-5010, 2004.

7. Kosar M, Goger F, Baser KHC, *In vitro* Antioxidant properties and phenolic composition of *Salvia virgata* Jacq. from Turkey, J Agric Food Chem 56, 2369–2374, 2008.
8. Tepe B. Antioxidant potentials and rosmarinic acid levels of the methanolic extracts of *Salvia virgata* (Jacq), *Salvia staminea* (Montbret & Aucher ex Benth) and *Salvia verbenaca* (L.) from Turkey, Bioresource Technol 99, 1584-1588, 2008.
9. Loizzoa M, Tundisa R, Confortia F, Menichinia F, Bonesia M, Nadjafib F, Giuseppe Fregac N, Menichinia F, *Salvia leriifolia* Benth. (Lamiaceae) extract demonstrates *in vitro* antioxidant properties and cholinesterase inhibitory activity, Nutr Res 30, 823-830, 2010.
10. Deans SG, Simpson EJM. Antioxidants from *Salvia officinalis*, In: The Genus *Salvia*, Ed(s): Kintzios SE, pp 185-192, Harwood Academic Publishers, Amsterdam, 2000.
11. Senol SF, Orhan İ, Celep F, Kahraman A, Doğan M, Yılmaz G, Sener B, Survey of 55 Turkish *Salvia* taxa for their acetylcholinesterase inhibitory and antioxidant activities, Food Chem 120, 34-43, 2010.
12. Göger F, Determination of antioxidant activity and the compositions of *Salvia virgata* Jacq. and *Salvia halophila* Hedge. Master of thesis, Anadolu University Graduate Institute of Health Science Department of Pharmacognosy, Eskişehir, Turkey, 2006.
13. Baytop T, Türkiye’de Kullanılan Tıbbi Bitkiler. In: Türkiye’de Bitkiler ile Tedavi Geçmişte ve Bugün, pp. 143, İkinci Baskı, Nobel Tıp Kitabevleri, İstanbul, Turkey, 1999.
14. Singleton VL, Orthofer R, Lamuela-Raventos RM, Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. In: Methods in Enzymology, Ed(s): Packer L, pp 152–315, Academic Press, San Diego, 1999.
15. Miliauskas G, Venskutonis PR, Van Beek TA, Screening of radical scavenging activity of some medicinal and aromatic plant extracts, Food Chem 85, 231–237, 2004.
16. Gyamfi MA, Yonamine M, Aniya Y, Free-radical scavenging action of medicinal herbs from Ghana *Thonningia sanguinea* on experimentally induced liver injuries, Gen Pharm 32, 661–667, 1999.
17. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice- Evans C, Antioxidant activity applying an improved ABTS radical cation decolorisation assay, Free Radical Bio Med 26, 1231–1237, 1999.
18. Carter P, Spectrophotometric determination of serum iron at the submicrogram level with a new reagent (ferrozine), Anal Biochem 40, 450–458, 1971.
19. Aruoma OI, Spencer J, Warren D, Jenner P, Butler J, Halliwell B, Characterization of food antioxidants, illustrated using commercial garlic and ginger preparations, Food Chem 60, 149–156, 1997.
20. Cuvelier ME, Berset C, Richard H, Antioxidant constituents in sage (*Salvia officinalis*), J Agric Food Chem 42, 665-669, 1994.
21. Adzet T, Canigüeral S, Iglesias J, A chromatographic survey of polyphenols from *Salvia* species, Biochem Syst Ecol 16, 29-32, 1988.
22. Lu Y, Foo LY, Polyphenolics of *Salvia*, Phytochem 59, 117–140, 2002.
23. Miura K, Kikuzaki H, Nakatani N, Antioxidant activity of chemical components from sage (*Salvia officinalis* L.) and thyme (*Thymus vulgaris* L.) measured by the oil stability index method, J Agric Food Chem 50, 1845-1851, 2002.
24. Matsingou TC, Petrakis N, Kapsokefalou M, Salifoglou A, Antioxidant activity of organic extracts from aqueous infusions of sage, J Agric Food Chem 51, 6696-6701, 2003.
25. Lu Y, Foo LY, Antioxidant activities of polyphenols from sage (*Salvia officinalis*), Food Chem 75, 197–202, 2001.
26. Bandoniene D, Murkovic M, Pfannhauser W, Venskutonis PR, Gruzdiene D, Detection and activity evaluation of radical scavenging compounds by using DPPH free radical and on-line HPLC-DPPH methods, Eur Food Res Technol 214, 143-147, 2002.
27. Deng J, Cheng W, Yang GA, Novel antioxidant activity index (AAU) for natural products using the DPPH assay, Food Chem 125, 1430–1435, 2011.
28. Pizzale L, Bortolomeazzi R, Vichi S, Ubergger E, Conte LS, Antioxidant activity of sage (*Salvia officinalis* and *S. fruticosa*) and oregano (*Origanum onites* and *O. onites*) extracts related to their phenolic compound content, J Agric Food Chem 82, 1645-1651, 2002.
29. Siddhuraju P, Becker K, Studies on antioxidant activities of mucuna seed (*Mucuna pruriens* var *utilis*) extract and various nonprotein amino/imino acids through *in vitro* models. J Sci Food Agric 83, 1517–1524, 2003.
30. Koleva II, Van Beek TA, Linsen JPH, De Groo A, Evstatieva LN, Screening of plant extracts for antioxidant activity: A comparative study on three testing methods, Phytochem Anal 13, 8–17, 2002.

31. Pukalskas A, Van Beek TA, de Waard P, Development of a triple hyphenated HPLC-radical scavenging detection-DAD-SPENMR system for the rapid identification of antioxidants in complex plant extracts, *J Chromatogr A* 1074, 81–88, 2005.
32. Dorman HJD, Hiltunen R, Fe(II) reductive and free radical scavenging properties of summer savory (*Satureja hortensis* L.) extract and subfractions, *Food Chem* 88,193–199, 2004.
33. Halliwell B, Antioxidants: the basics – what they are and how to evaluate them, *Adv Pharmacol* 38, 3–20,1997.
34. Melidou M, Riganakos K, Galaris D, Protection against nuclear DNA damage offered by flavonoids in cells exposed to hydrogen peroxide: The role of iron chelation, *Free Radical Bio Med* 39, 159 –1600, 2005.
35. Velioglu YS, Mazza G, Oomah BD, Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products, *J Agr Food Chem* 46, 4113–4117, 1998.
36. Floegel A, Kim DO, Chung SJ, Koo SI, Chun OK, Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods, *J Food Compos Anal* 24, 1043–1048, 2011.
37. Dorman HJD, Bachmayer O, Kosar M, Hiltunen R, Antioxidant properties of aqueous extracts from selected Lamiaceae species grown in Turkey, *J Agr Food Chem* 52, 762-770, 2004.

Received: 08.10.2015

Accepted: 25.02.2016