

Diagnostic Value of Diffusion Weighted Imaging in Sacroiliitis

Sakroileitlerde Difüzyon Ağırlıklı Görüntülemenin Tanı Değeri

Yasemin Durum Polat¹, Can Çeviko², Abdullah Utku Şeno², Emir Hüseyin Navaei¹

¹Adnan Menderes University Faculty of Medicine, Department of Radiology, Aydın, Turkey

²Akdeniz University Faculty of Medicine, Department of Radiology, Antalya, Turkey



Keywords

Sacroiliac joint, diffusion weighted imaging magnetic resonance imaging, inflammatory back pain

Anahtar Kelimeler

Sakroiliak eklem, difüzyon ağırlıklı görüntülemenin manyetik rezonans görüntüleme, enflamatuvar bel ağrısı

Received/Geliş Tarihi : 20.12.2016

Accepted/Kabul Tarihi : 01.03.2017

doi:10.4274/meandros.33042

Address for Correspondence/Yazışma Adresi:

Yasemin Durum Polat MD,
Adnan Menderes University Faculty of
Medicine, Department of Radiology, Aydın,
Turkey
Phone : +90 505 229 15 35
E-mail : yasemindurum@gmail.com
ORCID ID: orcid.org/0000-0002-4452-6632

©Meandros Medical and Dental Journal, Published by Galenos Publishing House.
This is article distributed under the terms of the Creative Commons Attribution NonCommercial 4.0 International Licence (CC BY-NC 4.0).

Abstract

Objective: Early diagnosis of acute sacroiliitis and related disorders are quite important for preventing the irreversible results of these diseases (such as ankylosis) and increasing the efficiency of treatment. Magnetic resonance imaging (MRI) is the most important modality in early diagnosis. Routine MRI examinations should include short tau inversion recovery and contrast enhanced fat saturated T1-weighted sequences. In our study, we evaluated the role of diffusion weighted imaging (DWI) MRI sequence in the diagnosis of acute sacroiliitis. Inter-observer and intra-observer reliability was also evaluated.

Materials and Methods: Seventy patients suffering from back pain and underwent sacroiliac joint MRI between October 2008 and December 2009 were retrospectively evaluated. Patients who were under age of 18 and had MRI examinations without contrast were excluded. Routine MRI sequences and DWI sequences (b100, b600, b1000) of 84 sacroiliac joints of 42 patients were evaluated retrospectively. Evaluations were made independently without any information about patient's name and clinical findings. First, DWI MRI examinations were evaluated by two radiologists separately (one of the radiologist made another evaluation after one week). After one month, sacroiliac joint MRI's of the patients were evaluated for osteitis and bone marrow enhancement (positive MRI) by two radiologists and decisions were made by consensus. A contrast-enhanced MRI study was accepted as the gold standard for the diagnosis of active sacroiliitis in the statistical evaluation. The sensitivity and specificity of DWI for active sacroiliitis were calculated. Also intra-observer and inter-observer concordances were made by Cohen kappa test in SPSS (Statistical Package for the Social Sciences) version 16 software.

Results: For the diagnosis of acute sacroiliitis, sensitivity and specificity values of DWI MRI were calculated as 82.1% and 87.5% respectively in 84 sacroiliac joints. By using Cohen's kappa test, moderate intra-observer correlation for first radiologist [kappa=0.684 (p<0.05)], weak intra-observer correlation for second radiologist [kappa=0.369 (p<0.05)] and moderate inter-observer correlation [kappa=0.512 (p<0.01)] was found.

Conclusion: In conclusion, DWI MRI examination is a non-invasive (contrast material administration is not needed), sensitive, fast diagnostic method in the diagnosis of acute sacroiliitis. Besides, DWI MRI may have a potential as a screening method in suspected cases, if supported with large studies.

Öz

Amaç: Sakroiliitin erken tanısı, eklemde geri dönüşümü mümkün olmayan (ankiloz gibi) sonuçların önlenmesinde veya geciktirilmesinde oldukça önemlidir. Son dönemlerde tedavide uygulanan etkin yeni ilaçlar erken tanının önemini bir kez daha gündeme getirmiştir. Halen manyetik rezonans görüntüleme (MRG) erken tanıda kullanılan en önemli görüntüleme yöntemidir. Rutin incelemeler yağ baskılı T2 ağırlıklı (short tau inversion recovery) ve postkontrast yağ baskılı T1 ağırlıklı sekansları içermelidir. Son yıllarda özellikle serebral enfarkt değerlendirmesinde kullanılan difüzyon ağırlıklı görüntüleme (DAG) sekansının kas iskelet sisteminde kullanımı görece yenidir. Çalışmamızda aktif sakroiliitin tanısında DAG MRG'nin tanısal etkinliği araştırılmıştır. Ayrıca DAG'lerin değerlendirilmesinde, gözlemci içi ve gözlemciler arası değişkenlik araştırılmıştır.

Gereç ve Yöntemler: Ekim 2008 ile Aralık 2009 tarihleri arasında kliniğimize bel ağrısı yakınmasıyla başvuran 70 hasta retrospektif olarak araştırılmıştır. On sekiz yaş altında olan, tetkiklerinde kontrast madde kullanılmayan veya dosyalarına ulaşamayan olgular çalışma dışı bırakılarak 42 hastanın MRG kesitleri ile b100, b600, b1000 değerlerindeki difüzyon ağırlıklı görüntüleri, klinik bilgileri bilinmeksizin incelenmiştir. DAG'nin aktif sakroiliitin tespitindeki tanısal etkinliğinin araştırılması amacı ile 42 olgunun, 84 sakroiliak eklemde ait DAG iki ayrı radyolog tarafından (bir radyoloğun ikinci kez değerlendirmesiyle birlikte) birbirinden bağımsız olarak difüzyon değişikliği açısından değerlendirilmiştir. Bu değerlendirmeden 1 ay sonra iki radyolog birlikte rutin ve kontrastlı sekansları osteitis ve kemik iliğinde kontrast tutulumu (pozitif MRG) açısından tekrar değerlendirmiştir. Bulguların istatistiksel olarak değerlendirilmesinde 42 olguya ait 84 sakroiliak eklem için, aktif sakroiliitin tanısında, kontrastlı rutin sakroiliak eklem MRG incelemesi sonuçları altın standart olarak kabul edilerek DAG'lerin aktif sakroiliitin tanısındaki duyarlılık ve özgüllükleri hesaplandı. Ayrıca gözlemci içi ve gözlemciler arası uyum Cohen kappa testi ile SPSS (Statistical Package for the Social Sciences) sürüm 16 programında yapıldı.

Bulgular: DAG'nin aktif sakroiliiti saptamadaki saptamadaki duyarlılığı %82,1 ve özgüllüğü %87,5 olarak bulunmuştur. Çalışmamızda DAG'lerin değerlendirilmesinde, 1. radyolog için gözlemci içi uyum orta derecede, 2. radyolog için gözlem içi uyum düşük derecede, gözlemciler arası uyum orta derecede bulunmuştur [Cohen kappa testinde 1. radyolog için gözlem içi kappa değeri 0,684 ($p<0,05$), 2. radyolog için gözlemci içi kappa değeri 0,369 ($p<0,05$), gözlemciler arası kappa değeri 0,512 ($p<0,01$)]. İkinci radyolog için gözlemci içi uyumun düşüklüğü değerlendirme yapan radyoloğun klinik tecrübe azlığına bağlanmıştır.

Sonuç: Çalışmamızın sonuçlarına göre, enflamatuvar bel ağrısı yakınması olan olgularda akut sakroiliitin saptanmasında, DAG tarama yöntemi olarak kullanılabilir hızlı, kontrast maddeye ihtiyaç göstermeyen, invaziv olmayan, görece ucuz bir yöntem olarak gelecekte kullanım alanı bulabilecektir. Ancak bu konuda kesin bir önermede bulunmak için, daha geniş seriler ile yapılacak prospektif çalışmalara ihtiyaç vardır.

Introduction

Sacroiliac joint (SIJ) involvement can be seen in many inflammatory, infectious, metabolic and degenerative diseases (1,2). In this wide disease spectrum, most common involvement is seen in seronegative spondyloarthropathies. In this group especially in ankylosing spondylitis (AS) as a prototype disease, SIJ inflammation is the most common and characteristic form of involvement (3). Early diagnosis is crucial to prevent or delay irreversible outcomes such as ankylosis. For many decades radiography has been used for diagnosis and also follow up the disease. However imaging limitations due to complex anatomy of the joint and more importantly late onset of radiographic manifestations have been disadvantages of this modality. Over the time for imaging of inflammation this joint scintigraphy, ultrasonography, computed tomography and magnetic resonance imaging (MRI) have been started to be used (4-9). Now with MRI, early imaging findings that could not be detected before with other modalities can be seen. Short tau inversion recovery (STIR), fat-suppressed

T2-weighted and contrast-enhanced T1-weighted MRI sequences have been shown that are highly sensitive in detection of early inflammatory changes (6,7,9). Characteristic SIJ lesion types in MRI are evaluated in two groups; active inflammatory lesions (bone marrow edema, capsulitis, synovitis, enthesitis) and chronic inflammatory lesions (sclerosis, erosion, fat deposition, ankylosis) (10). In some recent studies, it has been reported that diffusion weighted imaging (DWI) MRI as a more advanced technique, has a similar diagnostic efficiency comparing to conventional MRI in early detection of SIJ inflammatory findings (7).

In this study, we aimed to investigate the efficacy of DWI MRI for detection of early sacroiliitis (SI).

Materials and Methods

Seventy patients suffering from back pain and underwent SIJ MRI between October 2008 and December 2009 were retrospectively evaluated. Patients under age of 18, with history of surgery and foreign metallic bodies, and also the patients who unable to be applied contrast agents or unable to

reach their patient files were excluded from study. Routine MRI sequences and DWI sequences (b100, b600, b1000) of 84 SIJs of 42 patients were evaluated retrospectively. Evaluations were made independently without any information about patient's name and clinical findings. First, DWI MRI examinations were evaluated by two radiologists separately (one of the radiologist made another evaluation after one week). After one month, SIJ MRI's of the patients were evaluated for osteitis and bone marrow enhancement (positive MRI) by two radiologists and decisions were made by consensus. The areas with diffusion changes in b100, b600 and b1000 sequences were compared statistically with areas we evaluated as active SI in STIR and contrast enhanced T1-weighted sequenced. According to obtained findings, the sufficiency and reliability of singly DWI in detection of active SI were studied and also concordance among the different diffusion values was observed. This imaging evaluation was made independent of clinical findings. DWI sequences were evaluated by two radiologists separately whereas routine and contrast enhanced sequences evaluated together collaboratively. In this project, intra-observer and inter-observer variability were also statistically studied.

Magnetic Resonance Imaging Protocols

All patients underwent conventional MR imaging and DWI at 1.5 T MR scanner (Avanto Siemens, Erlangen, Germany). In routine SIJ MR imaging, based on a sagittal survey image, SIJ oblique coronal and oblique transverse slices were obtained parallel to the long axis of sacrum and parallel to the short axis of sacrum (vertical to coronal slices) respectively. In this study MRI protocols were; pre-contrast series; oblique coronal plane T1A turbo spin eko (TSE) [the echo time/ the repetition time (TE/TR): 10/639] and STIR (TE/TR: 255/4000); oblique transverse plane T1A TSE (TE/TR: 11/482), T2A fat-suppressed TSE (TE/TR: 92/4560) slices, post-contrast (0.1 mmol/kg gadolinium bolus i.v. infusion) oblique coronal fat-suppressed T1A TSE (TE/TR: 10/845) and oblique transverse fat-suppressed T1A TSE (TE/TR: 11/482) as routine MRI sequences. Also, oblique coronal b100, b600, b1000 diffusion weighted sequences (TE/TR: 94/3400) were obtained. Other parameters for routine sequences were; field of view (FOV): 240 mm (STIR sequence FOV: 300 mm), matrix: 320x320, slice thickness: 4

mm. For DWI sequences these parameters were; FOV: 360 mm, matrix: 168x192, slice thickness: 4 mm, number of excitations: 3.

Statistical Analysis

For diagnosis of SI for 84 SIJs of 42 patients, contrast-enhanced routine SIJ MRI protocols were used as gold standard and compares with DWI findings. Based on acquired data, sensitivity and specificity of DWI were calculated. Intra-observer and inter-observer concordances were also made by Cohen kappa test in SPSS (Statistical Package for the Social Sciences) version 16.0 software.

Result

In our research in 28 of 84 SIJs, bone marrow edema (Figure 1, 2) and enhancement in the post-




Figure 1. Coronal short tau inversion recovery images show hyperintense lesions consistent with bone marrow edema in bilateral sacroiliac joints




Figure 2. Coronal short tau inversion recovery images show hyperintense lesions consistent with bone marrow edema in bilateral sacroiliac joints

contrast series were detected and were diagnosed as acute SI (Figure 3). Twenty three out of 28 SIJs who have been diagnosed as acute SI have been evaluated as positive by DWI (Figure 4, 5). According to this, for both methods (84 SIJs) the sensitivity and specificity have been calculated as 82.1% and 87.5% respectively. In our study, 20 patients without acute SI also showed no DWI findings. Among the cases with DWI negative finding we found SI in 3 patients on the right, in 6 patients on the left. In the retrospective evaluation of findings; in one patient who was diagnosed as acute SI by routine imaging, DWI evaluated as normal. We considered that DWI findings could not be noted due to the artifacts in this case. Similarly, in retrospective evaluation of cases with positive DWI findings but




Figure 3. Contrast enhanced fat-sat T1-weighted image shows enhancement in bilateral sacroiliac joints




Figure 4. Lesions are hyperintense on coronal diffusion-weighted images at b values of 1.000

negative routine exam findings, we think that the reason of the false positive evaluation was the artifacts. We found positive MRI findings in 6 cases among the 16 cases referred for inflammatory back pain, in 9 cases among the 14 cases with AS and 5 cases among the 12 cases complain of mechanic back pain. We noticed positive MRI finding in almost half of the cases with mechanic back pain. It should be remembered that DWI changes in bone marrow is also reported in trauma and infection in the literature (2). In our study intra and inter-observer coherence have been tested with the Cohen kappa test. According to the Cohen kappa test, in the evaluation of the results of the first radiologist and the gold standard test a significant coherence was detected with the kappa value as 0.684 ($p < 0.05$). According to this test sensitivity has been determined as 82.1% and specificity as 87.5%. Receiver operating characteristic area this analysis has been estimated as 0.848 (Graphic 1). Kappa test, weak intra-observer correlation for second radiologist [$\text{kappa} = 0.369$ ($p < 0.05$)] was found. Between the first and second radiologist, the inter-observer variability was again questioned with the Cohen kappa test and the kappa value has been detected as 0.512 ($p < 0.01$) (minimal coherence). Minimal coherence between observers was linked to lack of clinical experience and insufficient experience in interpretation of SIJ DWI of second observer. Increased consistency of the results of the second evaluation with the gold standard was linked to the improved experience and knowledge of SIJ anatomy of the observer. Intermediate consistence between first observer and the gold standard was





Figure 5. Lesions are hyperintense on axial diffusion-weighted images at b values of 1.000



Graphic 1. Receiver operating characteristic area for this analysis (0.848) showing graphic
ROC: Receiver operating characteristic

thought to be a possible consequence of false positive results due to some DWI artifacts. b100, b600 and b1000 DWI were evaluated consecutively. In SI cases diffusion signal changes were seen in all b values, however the highest accuracy was observed in b100 value images.

Discussion

In literature the usage of DWI in the SIJ is quite limited. In a study on 30 patients, Gaspersic et al. (11) has evaluated different treatment types upon enthesitis and osteitis of AS by using DWI and dynamic MRI and has detected the effectiveness of both treatment. It has been pointed out that DWI can be used to evaluate the treatment response in rheumatology patients. Bozgeyik et al. (7), in their study on 13 patients with inflammatory back pain, and the other patients with low (mechanic) back pain (total of 42 patients), they have compared the DWI findings of subchondral bone changes in early SI and bone marrow changes with post-contrast T1-TSE sequence. They reported that acute lesions in early SI can be detected with both DWI and contrast-enhanced T1-weighted sequences with a similar accuracy. They obtained different b values (DWI's; b100, 600 and 1000) and evaluated separately. They found that normal areas and areas of acute SI showed significant difference in apparent diffusion coefficient (ADC) values. Ai et al. (12) has researched the value of DWI in early detection of AS. In their study they have calculated the mean ADC values in DWI and whole-body DWI images in 16 AS patients, 18 patients with mechanical back pain and 18 healthy volunteers, and have found a significantly high ADC values in early AS patients comparing to other two groups. Dallaudière

et al. (13) has compared the ADC values of 95 patients (two separate groups with degenerative changes and spondyloarthritis) and have found significantly higher values in the second group. Sahin et al. (14) has reported a significant difference between ADC values of normal subchondral bone marrow and the areas with subchondral bone marrow edema in active SI patients. They found that DWI with ADC values may be for accurately diagnosing inflammatory SI. Sanal et al. (15) has studied the role of DWI in evaluation of activity of SI in AS patients for a long period. In their study with 21 AS patients and 7 healthy volunteers they have measured the contrast-to-noise ratio (CNR) values in subchondral bone marrow DWI and contrast enhanced series. As a result, they have shown that there is no significant difference in CNR values of DWI and contrast enhanced series in the active lesion areas detected on the STIR sequence. Also, they have found a statistically significant difference in comparison with control group.

Study Limitations

In our study, the most important limitations were limited number of patients, being a retrospective study and inexperience of the second observer in interpretation of SIJ DWI findings. Lack of patient follow up was the other limitation of this study. Moreover in our study correlation of imaging findings with laboratory data was not investigated. No availability of antibodies to streptolysin and C-reactive protein levels which have been widely used for detection of acute inflammation in SI based on current Assessment of Spondylo Arthritis international Societ classification criteria and therefore lack of research of correlation with laboratory findings was a significant limitation of our study.

Conclusion

We believe that by larger patient numbers and prospectively designed studies and with involvement of the investigation of correlation between laboratory and imaging findings and including patient follow up, more generalizable results will be obtained.

Ethics

Ethics Committee Approval: For study, approval of Ethics Committee of Scientific Research Ethics Committee of Akdeniz University Faculty of Medicine was obtained (approval no: 18).

Informed Consent: Consent form was not filled, because this study was done retrospectively.

Peer-review: Internally peer-reviewed.

Authorship Contributions

Concept: Y.D.P., C.Ç., Design: Y.D.P., C.Ç., A.U.Ş., Data Collection or Processing: Y.D.P., C.Ç., Analysis or Interpretation: Y.D.P., C.Ç., Literature Search: Y.D.P., C.Ç., Writing: Y.D.P., C.Ç., E.H.N.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Sutton D. Textbook of Radiology and imaging, 7th.edition, Elsevier Science Ltd, Edinburgh, UK 2002; 2: 1201-40.
2. Dahnert W. Radiology Review Manual, 6th.edition; 2007.
3. Dougados M, van der Linden S, Juhlin R, Huitfeldt B, Amor B, Calin A, et al. The European Spondylarthropathy Study Group preliminary criteria for the classification of spondylarthropathy. *Arthritis Rheum* 1991; 34: 1218-27.
4. Lacout A, Rousselin B, Pelage J. CT and MRI of Spine and Sacroiliac Involvement in Spondyloarthropathy. *AJR* 2008; 191: 1016-23.
5. Arslan H, Sakarya ME, Adak B, Unal O, Sayarlioglu M. et al. Duplex and color Doppler sonographic findings in active sacroiliitis. *Am J Roentgenol* 1999; 173: 677-80.
6. Guglielmi G, Scalzo G, Cascavilla A, Carrotti M, Salaffi F, Grassi W. et al. Imaging of the sacroiliac joint. involvement in seronegative spondylarthropathies. *Clinical Rheumatology* 2009; 28:1007-19.
7. Bozgeyik Z, Özgöçmen S, Karakoç E. Role of Diffusion-Weighted MRI in the Detection of Early Active Sacroiliitis. *AJR* 2008; 191: 980-6.
8. Bredella MA, Steinback LS, Morgan S, Ward M, Davis JC. et al. MRI of the sacroiliac joints in Patients with Moderate to Severe Ankylosing Spondylitis. *AJR* 2006; 187: 1420-6.
9. Algin O, Gokalp G, Baran B, Ocakoglu G, Yazici Z. Evaluation of sacroiliitis: contrast-enhanced MRI with subtraction technique. *Skeletal Radiology* 2009; 38: 983-8.
10. Sieper J, Rudwaleit M, Baraliakos X, Brandt J, Braun J, Burgos-Vargas R, et al. The Assessment of SpondyloArthritis international Society (ASAS) handbook: a guide to assess spondyloarthritis. *Ann Rheum Dis* 2009; 68: 1-44.
11. Gaspersic N, Sersa I, Jevtic V, Tomsic M, Praprotnik S. Monitoring ankylosing spondylitis therapy by dynamic contrast-enhanced and diffusion-weighted magnetic resonance imaging. *Skeletal Radiology* 2008; 2: 123-31.
12. Ai F, Ai T, Li X, Hu D, Zhang W, Morelli JN. Value of diffusion-weighted magnetic resonance imaging in early diagnosis of ankylosing spondylitis. *Rheumatol Int* 2012; 32: 4005-13.
13. Dallaudière B, Dautry R, Preux P, Perozziello A, Lincot J, Schouman-Claeys E, et al. Comparison of apparent diffusion coefficient in spondylarthritides axial active inflammatory lesions and type 1 modic changes. *European Journal of Radiology* 2014; 366-70.
14. Sahin N, Hacibeyoglu H, Ince Ö, Solak A, Uyar B, Erol Ö, et al. Is there a role for DWI in the diagnosis of sacroiliitis based on ASAS criteria? *Int J Clin Exp Med* 2015; 8: 7544-52.
15. Sanal HT, Yilmaz S, Kalyoncu U, Cinar M, Simsek İ, Erdem H, et al. Value of DWI in visual assessment of activity of sacroiliitis in longstanding ankylosing spondylitis patients. *Skeletal Radiol* 2013; 42: 289-93.