



The Level of Knowledge of Pelvic Floor Dysfunction After Delivery in Women who Attended to a Tertiary Center

Üçüncü Basamak bir Merkeze Başvuran Kadınlarda Doğumdan Sonra Pelvik Taban Disfonksiyonu Bilgi Düzeyi

Murat Ekin¹, Cihan Kaya¹, Emine Öztürk², Hüseyin Cengiz¹, Gülden Uzer³, Levent Yaşar¹

Abstract / Öz

Introduction: Pelvic floor disorders affect women in all age groups and cause poor quality of life with economic burden. The aim of the present study was to investigate the attitude of women who were admitted to our clinic about the relationship between mode of delivery and pelvic floor disorders and pelvic muscle exercises.

Methods: A total of 1316 women who had attended our outpatient gynecology clinic for various complaints were interviewed by an expert gynecologist. Demographic data including age, marital status, education, health insurance, menopausal status, parity, mode of deliveries, birth weights, and also lower urinary tract symptoms were included in the interview. All the participants were asked to complete a verbal modified Pelvic Risk Knowledge Score (PRKS) questionnaire regarding their knowledge about pelvic floor disorder risks.

Results: The mean modified PRKS was 4.95 ± 2.5 . Of the patients, 26.8% had a diagnosis of stress urinary incontinence, 14.3% had urgency, 19.8% had frequency, 11.1% had >stage 2 pelvic organ prolapse (POP), 7.4% did not even hear about pelvic muscle exercises, and 17.6% did not even inform about pelvic muscle exercises by a health care provider. PRKS was significantly higher in multiparous women than in primiparous women ($p < 0.0001$). Vaginal birth also significantly increased the PRKS with respect to cesarean delivery ($p = 0.006$). Women with cesarean deliveries had significantly increased PRKS with respect to nulliparous women ($p < 0.0001$).

Conclusion: Episiotomy, menopause, lower urinary tract symptoms, and POP are the factors that significantly increase PRKS.

Keywords: Pelvic floor disorders, pelvic muscle exercises, pelvic risk, vaginal birth

Amaç: Pelvik taban hastalıkları tüm yaş gruplarındaki kadınları etkiler ve ekonomik kayıpla birlikte kötü hayat kalitesine neden olur. Bu çalışmada kliniğimize başvuran kadınların doğum şekli ile pelvik kas egzersizleri ve pelvik taban hastalıkları arasındaki ilişki hakkındaki yaklaşımlarını incelemeyi amaçladık.

Yöntemler: Çeşitli şikayetlerle jinekoloji polikliniğimize başvurmış 1316 kadın uzman bir jinekolog tarafından sorgulandı. Yaş, evlilik durumu, eğitim, sağlık sigortası, menapoz durumu, parite, doğum şekilleri, doğum kiloları ve alt üriner sistem semptomları gibi demografik bilgiler sorgulamaya dahil edildi. Sonrasında tüm katılımcıların pelvik taban hastalıkları riski hakkında bilgilerini ölçmek amacıyla modifiye sözel Pelvik Risk Bilgi Skoru (PRBS)'nu tamamlamaları istendi.

Bulgular: Ortalama modifiye PRBS $4,95 \pm 2,5$ idi. Hastaların %26,8'i stres üriner inkontinans (SUI)'a sahipti. Hastaların %14,3'ünde ani sıkışma hissi ve %19,8'inde sık idrara çıkma mevcuttu. Hastaların %11,1'i evre>2 pelvik organ prolapsusu (POP), %7,4'ü hiç pelvik kas egzersizlerini hiç duymadığını ve % hastaların 17,6'sı hiçbir sağlık personeli tarafından pelvik kas egzersizleri hakkında bildirilmemişti. PRBS multipar kadınlarda primiparlardan anlamlı ölçüde yüksekti ($p < 0,0001$). Vajinal doğum PRBS'yi sezaryene göre önemli ölçüde yükseltti ($p = 0,006$) ve sezaryen olmuş olan kadınlar nulliparlara göre anlamlı ölçüde artmış PRBS sahipti ($p < 0,0001$).

Sonuç: Epizyotomi varlığı, menapoz, alt üriner sistem semptomları ve POP PRBS'yi anlamlı olarak arttıran faktörlerdir.

Anahtar Kelimeler: Pelvik taban bozuklukları, pelvik kas egzersizleri, pelvik risk, vajinal doğum

The abstract of this paper has been presented as a poster presentation in 1st UHS Congress of Pelvic Floor Disorders, 5-7 May 2017, Istanbul, Turkey.

ORCID IDs of the authors: M.E. 0000-0002-4525-5125; C.K. 0000-0003-4175-7694; L.Y. 0000-0002-8679-2699.

¹Department of Gynecology and Obstetrics, Health Sciences University Bakırköy Dr. Sadi Konuk Training and Research Hospital, Istanbul, Turkey

²Clinic of Gynecology and Obstetrics, Istanbul Bahçelievler State Hospital, Istanbul, Turkey

³Üsküdar Doğançılar Family Health Center, Istanbul, Turkey

Address for Correspondence/

Yazışma Adresi:

Cihan Kaya, Department of Gynecology and Obstetrics, Health Sciences University Bakırköy Dr. Sadi Konuk Training and Research Hospital, Istanbul, Turkey
E-mail: drchankaya@gmail.com

Received/Geliş Tarihi: 23.10.2017

Accepted/Kabul Tarihi: 22.02.2018

© Copyright 2018 by Available online at
istanbulmedicaljournal.org

© Telif Hakkı 2018 Makale metnine
istanbulmedjergisi.org web sayfasından ulaşılabilir.

Introduction

Pelvic floor disorders affect women in all age groups and cause poor quality of life and economic burden (1). Cross-sectional studies suggest that women who had vaginal birth are more susceptible to urinary incontinence, fecal incontinence, pelvic organ prolapse (POP), and sexual dysfunction than women who had only cesarean deliveries later in life (2, 3). Pelvic floor muscle exercise is defined as the repetitive selective, voluntarily contraction and relaxation of the pelvic muscles. It aims to strengthen the pelvic muscles to support the urethra and increase the urethral sphincteric function (4). It has been established that pelvic floor muscle exercises decrease urinary incontinence in pregnancy, postpartum period, and later in life. It also reduces the episodes of postpartum fecal incontinence and improves sexual dysfunction in the postpartum period (5-7).

The aim of the present study was to investigate the perception of women on the relationship between mode of delivery and pelvic floor disorders and to investigate the knowledge on pelvic floor muscle exercises.

Methods

This was a cross-sectional observation study. Ethics committee approval was received for this study from the Ethics Committee of Health Sciences University Bakırköy Dr. Sadi Konuk Training and Research Hospital (2014/27). Written informed consent was obtained from all of the participants.

Overall, 1316 women who had attended the outpatient gynecology clinic for various complaints were interviewed by an expert gynecologist. Demographic data including age, marital status, education, health insurance, menopausal status, parity, mode of deliveries, birth weights, and lower urinary tract symptoms were included in the interview. The clinical examination data for the presence of episiotomy, stress urinary incontinence (SUI), urgency, frequency, and POP were recorded.

All of the participants were asked to complete a verbal modified Pelvic Risk Knowledge Score (PRKS) questionnaire adapted from Dunbar et al. (8) regarding their knowledge about pelvic floor risks associated with the delivery methods and their awareness about the pelvic muscle exercises (Table 1). For analysis of the awareness of the risk of vaginal delivery (a nine-item questionnaire), addressing this topic were scored along each subject's number of deliveries. For the yes/no questions, a score of 1 was assigned to "Yes" response and 0 to "No" response. Vaginal responses were scored as 1, and cesarean responses as 0. Then, a composite risk score modified PRKS was obtained by adding all of the scores along with 10 items of the number of deliveries. A PRKS of 0 indicates no knowledge, whereas higher scores indicate a higher level of knowledge on a linear scale.

Statistical Analysis

The Number Cruncher Statistical System 2007 statistical software (NCSS, UT, USA) was used for data analysis. Descriptive statistical analysis was expressed as mean±standard deviation. One-way analysis of variance was used for normally distributed data. The Tukey test was used for post hoc analysis of parametric data. A paired sample t-test and the chi-square test were used for comparison of qualitative and quantitative data. A univariate model was created to determine the effect of episiotomy, menopause, SUI, urgency, frequency, and POP on PRKS. A Spearman correlation analysis was performed to determine any correlation between parity and PRKS. A $p < 0.05$ was accepted as statistically significant.

Table 1. Questionnaire items: Pelvic risk knowledge score

1. In your opinion, does a vaginal delivery (natural childbirth) increase your risk of problems controlling your bladder (difficulty holding your water)? Yes/No
2. In your opinion, does a cesarean delivery increase your risk of problems controlling your bladder (difficulty holding your water)? Yes/No
3. In your opinion, which type of delivery could cause more harm to pelvic floor? Vaginal/cesarean
4. In your opinion, does a vaginal delivery (natural childbirth) increase your risk of problems controlling your bowels (leakage of gas and stool)? Yes/No
5. In your opinion, does a vaginal delivery have a negative effect on your sexual function? Yes/No
6. In your opinion, does a cesarean delivery have a negative effect on your sexual function? Yes/No
7. In your opinion, can exercises of the muscles in your pelvic area help lessen the chance of your developing bladder and/or bowel problems later in life? Yes/No
8. Have you ever heard about pelvic muscle exercises? Yes/No
9. Have you ever informed about or suggested to make pelvic muscle exercises to a health care provider? Yes/No

Results

Table 2 shows the demographic and clinical examination findings of the patients. The mean modified PRKS was 4.95 ± 2.5 . Among the patients, 26.8% were diagnosed with SUI, 14.3% had urgency, 19.8% had frequency symptoms, and 11.1% had >stage 2 POP at pelvic examination. Table 3 shows the responses that were given to a nine-item questionnaire by the patients. The responses to questions 8 and 9 were 7.4% and 17.6%, respectively, indicating that awareness of pelvic muscle exercises is really poor

Table 2. The descriptive characteristics and clinical examination findings of the study population

	min-max	mean±SD	
Age	14-81	43.27±12.39	
Parity	0-13	2.22±1.8	
Vaginal birth	0-13	1.95±1.87	
Cesarean birth	0-4	0.28±0.62	
PRKS	0-18	4.95±2.5	
		n	%
Number of deliveries >4000 g	0	1.142	86.8
	1	147	11.2
	≥2	27	2.1
Number of operative delivery	0	1.263	96.0
	≥1	31	2.4
	≥2	22	1.7
Episiotomy	No	780	59.3
	Yes	536	40.7
Marital status	Married	1.079	82.0
	Single	107	8.1
	Divorced	69	5.2
	Widow	61	4.6
Menopause	No	835	63.4
	Yes	481	36.6
Educational status	Illiterate	127	9.7
	Primary school	818	62.2
	High school	238	18.1
Health insurance	University	133	10.1
	None	65	4.9
	Yes	1.251	95.1
Stress urinary incontinence	No	963	73.2
	Yes	353	26.8
Urgency	No	1.128	85.7
	Yes	188	14.3
Frequency	No	1.056	80.2
	Yes	260	19.8
Pelvic organ prolapse (>stage 2)	No	1.170	88.9
	Yes	146	11.1
PRKS: pelvic risk knowledge score; min: minimum; max: maximum; SD: standard deviation			

and health care providers hardly inform their patients about pelvic muscle exercises. For univariate analysis, presence of episiotomy, menopause, SUI, urgency, frequency and POP were the factors that significantly increase the PRKS ($p < 0.0001$) (Table 4). There was a significant correlation between parity and PRKS ($p < 0.0001$). PRKS was higher in nulliparous women than in primiparous women ($p < 0.001$). It was also significantly higher in multiparous women than in primiparous women ($p > 0.0001$) (Table 5). Vaginal birth was also significantly increased in the PRKS with respect to cesarean delivery ($p = 0.006$), but women who had

cesarean deliveries had also significantly higher PRKS than nulliparous women ($p < 0.0001$) (Table 6).

Discussion

Women's attitude on vaginal delivery differs in different parts of the world in relation to traditions and socioeconomic status. The reasons for selecting preferentially vaginal delivery in a Turkish population include fear of surgery, desire of early recovery, and request for having a great number of children (9). In the present study, 49% and 68% of women were unaware that vaginal birth can cause urinary incontinence and fecal incontinence later in life, respectively. Only 37% of women thought that pelvic muscle exercises could help decrease the chance of developing bladder and bowel problems later in life. Among women, 82% did not even hear about pelvic muscle exercises. The present study suggested that there is an important lack of knowledge about the relationship between vaginal birth and pelvic floor disorders. In a previous review, pelvic floor muscle exercises significantly prevent urinary incontinence in late pregnancy and postpartum for the continent women before pregnancy (10). Consistent with these findings, another review of 22 trials by Boyle et al. (6) with 8484 pregnant or postpartum women revealed that continent pregnant women who had intensive antenatal pelvic floor muscle exercises are less likely to report urinary incontinence in late pregnancy and at 6 months of postpartum period. Pelvic muscle exercises are recommended as the first-line management for the prevention of SUI during pregnancy and postpartum period. In addition, the National Institute for Health and Care Excellence suggests pelvic floor muscle exercises to all pregnant women for the prevention of SUI (5). According to the given results to the questions related with pelvic muscle exercises, Turkish women were poorly aware of these exercises, and health care providers were reluctant to inform and teach their patients for this common health burden.

In the present study, we have observed that aging, menopause, having episiotomy, lower urinary tract symptoms, and POP were significantly related with higher PRKS. This result can be explained

Table 3. The responses that were given to a nine-item questionnaire by the patients

		n	%
Question 1	No	648	49.2
	Yes	668	50.8
Question 2	No	1.066	81.0
	Yes	250	19.0
Question 3	Vaginal	562	42.7
	Cesarean	754	57.3
Question 4	No	896	68.1
	Yes	420	31.9
Question 5	No	896	68.1
	Yes	420	31.9
Question 6	No	1.064	80.9
	Yes	252	19.1
Question 7	No	488	37.1
	Yes	828	62.9
Question 8	No	1.084	82.4
	Yes	232	17.6
Question 9	No	1.219	92.6
	Yes	97	7.4

Table 4. The univariate analysis of PRKS and episiotomy, menopause, SUI, urgency, frequency, and POP

		n	PRKS	p
Episiotomy	No	780	4.71±2.74	0.0001
	Yes	536	5.3±2.04	
Menopause	No	835	4.53±2.35	0.0001
	Yes	481	5.68±2.58	
SUI	No	963	4.23±2.12	0.0001
	Yes	353	6.91±2.39	
Urgency	No	1128	4.59±2.29	0.0001
	Yes	188	7.08±2.62	
Frequency	No	1056	4.53±2.27	0.0001
	Yes	260	6.66±2.67	
POP	No	1170	4.66±2.32	0.0001
	Yes	146	7.28±2.63	

PRKS: pelvic risk knowledge score; SUI: stress urinary incontinence; POP: pelvic organ prolapse

Table 5. The correlation analysis between parity and PRKS

	n	PRKS (mean±SD)	p
Nulliparity	249	2.7±1.76	0.0001
Primiparity	167	3.49±1.5 ^α	
≥2 parity	900	5.84±2.29 ^{β,γ}	

^αp=0.001, nulliparity/primiparity; ^βp=0.0001, nulliparity/multiparity; ^γp=0.0001, primiparity/multiparity; PRKS: pelvic risk knowledge score

Table 6. The correlation analysis between mode of delivery and PRKS

		n	PRKS (mean±SD)	p
Mode of delivery	Vaginal birth	808	5.58±2.38	0.006
	C-section	259	5.14±2.22	
	Nulliparity	249	2.70±1.76	0.0001
	C-section	195	4.83±2.18	

PRKS: pelvic risk knowledge score; SD: standard deviation

as Turkish women are not informed about the consequences of vaginal birth, and they only realize this situation when they are symptomatic.

It is a well-known fact that parity increases urinary incontinence in premenopausal women. The first delivery has the most important effect on incontinence, whereas subsequent deliveries have a small but ongoing effect (11, 12). We have found that parity has a significant effect on PRKS, and there is a significant difference between PRKS of primiparous and multiparous women. Although PRKS was significantly lower in patients with a history of cesarean birth than in those with vaginal birth, there was also a significant difference between nulliparous women and participants with a history of cesarean birth, with nulliparous women with lower PRKS. This result can be attributed to the negative impact of pregnancy on pelvic floor muscles and urinary incontinence.

The present study has some limitations. First, although the small sample size can be a limitation of the study, our hospital is located in the largest region of Turkey and covers a population who had migrated from different parts of the country. Second, there is a lack of Turkish validation of PRKS. It can be validated after encouraging results of our results in future studies.

Conclusion

The results of the present study indicate that parity and lower urinary tract symptoms have a significant correlation. Although pelvic muscle exercises are recommended as the first-line management for the prevention of SUI during pregnancy and postpartum period, the knowledge of the Turkish population on this issue is poor, and health care providers should exert more effort to raise awareness in society.

Ethics Committee Approval: Ethics committee approval was received for this study from the Ethics Committee of Health Sciences University Bakırköy Dr. Sadi Konuk Training and Research Hospital (2014/27).

Informed Consent: Written informed consent was obtained from patients and patient's parents who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - M.E., C.K.; Design - M.E., H.C.; Supervision - C.K., L.Y.; Resources - G.U., H.C.; Materials - G.U., H.C.; Data Collection and/or Processing - G.U., E.Ö.; Analysis and/or Interpretation - G.U., E.Ö.; Literature Search - E.Ö., C.K.; Writing Manuscript - L.Y., M.E.; Critical Review - M.E., H.C.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

Etik Komite Onayı: Bu çalışma için etik komite onayı Sağlık Bilimleri Üniversitesi Bakırköy Dr. Sadi Konuk Eğitim ve Araştırma Hastanesi Etik Kurulu'ndan alınmıştır (2014/27).

Hasta Onamı: Yazılı onam çalışmaya katılan hastalardan/hasta ebeveynlerinden alınmıştır.

Hakem Değerlendirmesi: Dış bağımsız.

Yazar Katkıları: Fikir - M.E., C.K.; Tasarım - M.E., H.C.; Denetleme - C.K., L.Y.; Kaynaklar - G.U., H.C.; Malzemeler - G.U., H.C.; Veri Toplanması ve/veya İşlemesi - G.U., E.Ö.; Analiz ve/veya Yorum - G.U., E.Ö.; Literatür Taraması - E.Ö., C.K.; Yazıyı Yazan - L.Y., M.E.; Eleştirel İnceleme - M.E., H.C.

Çıkar Çatışması: Yazarların beyan edecek çıkar çatışması yoktur.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmiştir.

References

1. Memon H, Handa VL. Pelvic floor disorders following vaginal or cesarean delivery. *Curr Opin Obstet Gynecol* 2012; 24: 349-54. [CrossRef]
2. MacLennan AH, Taylor AW, Wilson DH, Wilson D. The prevalence of pelvic floor disorders and their relationship to gender, age, parity and mode of delivery. *BJOG* 2000; 107:1460-70. [CrossRef]
3. Rortveit G, Hannestad YS, Daltveit AK, Hunskaar S. Age- and type-dependent effects of parity on urinary incontinence: the Norwegian EPINCONT study. *Obstet Gynecol* 2001; 98: 1004-10. [CrossRef]
4. Haylen BT, de Ridder D, Freeman RM, Swift SE, Berghmans B, Lee J, et al. An international urogynecological association (IUGA)/international continence society (ICS) joint report on the terminology for female pelvic floor dysfunction. *Int Urogynecol J* 2010; 21: 5-26. [CrossRef]
5. National Institute for Health and Clinical Excellence (NICE). Urinary incontinence in women: management. 2013 Sept (cited 2015 October 30). Available from: URL: www.nice.org.uk/guidance/cg171/resources/urinary-incontinence-in-women-management-35109747194821.
6. Boyle R, Hay-Smith EJ, Cody JD, Mørkved S. Pelvic floor muscle training for prevention and treatment of urinary and faecal incontinence in antenatal and postnatal women. a short version Cochrane review. *NeuroUrol Urodyn* 2014; 33: 269-76. [CrossRef]
7. Mørkved S, Bø K. Effect of pelvic floor muscle training during pregnancy and after childbirth on prevention and treatment of urinary incontinence: a systematic review. *Br J Sports Med* 2013; 48: 299-310. [CrossRef]
8. Dunbar A, Ernst A, Matthews C, Ramakrishnan V. Understanding Vaginal Childbirth: What Do Women Know About the Consequences of Vaginal Childbirth on Pelvic Floor Health? *J Womens Health Phys Therap* 2011; 35: 51-6 [CrossRef]
9. Yıldız Ş, Çaypınar SS, Cengiz H, Dağdeviren H, Kanawati A. Awareness and perceptions of Turkish women towards delivery methods. *J Clin and Exp Invest* 2014; 5: 173-8.
10. Woodley SJ, Boyle R, Cody JD, Mørkved S, Hay-Smith EJC. Pelvic floor muscle training for prevention and treatment of urinary and fecal incontinence in antenatal and postnatal women *Cochrane Database Syst Rev* 2017; 12: CD007471.
11. Leijonhufvud A, Lundholm C, Cnattingius S, Granath F, Andolf E, Altman D. Risks of stress urinary incontinence and pelvic organ prolapse surgery in relation to mode of childbirth. *Am J Obstet Gynecol* 2011; 204: 70. [CrossRef]
12. Handa VL, Blomquist JL, Knoepp LR, Hoskey KA, McDermott KC, Muñoz A. Pelvic Floor Disorders 5-10 years after vaginal or cesarean childbirth. *Obstet Gynecol* 2011; 118: 777-84. [CrossRef]

Cite this article as: Ekin M, Kaya C, Öztürk E, Cengiz H, Uzer G, Yaşar L. The Level of Knowledge of Pelvic Floor Dysfunction After Delivery in Women who Attended to a Tertiary Center. *Istanbul Med J* 2018; 19 (4): 277-80.