

jcrpe-2018-0098.R2

Original Article

DOI: 10.4274/jcrpe.galenos.2019.2018.0098

No Associations Between Serum Lipid Levels or HOMA-IR and Asthma in Children and Adolescents: a NHANES Analysis

Short title: Lipids and Asthma in Children

Min Lu, MD,¹ Beirong Wu MD,¹ Rong Qiao, MD,² Haoxiang Gu,¹ Ying Din, MD,¹ Xiaoyan Dong, MD, PhD¹

¹Division of Pulmonary Medicine, Children's Hospital of Shanghai (Jing An Branch), Shanghai Jiao Tong University, Shanghai 200040

²Outpatient Department, Children's Hospital of Shanghai, Shanghai Jiao Tong University, Shanghai 200062

What is already known on this topic?

- Being overweight in childhood is associated with an increased risk of the development of allergic disease.
- A link has been shown between elevated lipid levels and the development of asthma/wheezing in children and adults.
- Hyperinsulinemia may be associated with the development of asthma.

What this study adds?

- Multivariate analyses found no associations between reduced high-density lipoprotein cholesterol, elevated low-density lipoprotein cholesterol, total cholesterol, and triglycerides and the presence of asthma in children or adolescents.
- Multivariate analyses found no associations between elevated fasting plasma glucose and the presence of asthma in children or adolescents.
- Multivariate analyses found no associations between HOMA-IR and the presence of asthma in children or adolescents.

Correspondence to: Rong Qiao MD, Outpatient Department, Children's Hospital of Shanghai, Shanghai Jiao Tong University, Shanghai, No. 355 Lu Ding Road, Shanghai 200062, P.R. of China

+86 021 62474880

Qiaorong@aliyun.com

24.04.2018

08.02.2019

<https://orcid.org/0000-0001-8150-8911>

Abstract

Objective: Studies have reported inconsistent results of the associations between lipids and insulin resistance (IR) and asthma. The purpose of this study was to examine the associations between abnormal serum lipid levels and homeostatic model assessment-insulin resistance (HOMA-IR) and the presence of current asthma in children and adolescents.

Methods: The United States National Health and Nutrition Examination Survey database from 1999 to 2012

was randomly searched for children (aged 3-11 years) and adolescents (aged 12-19 years) with and without asthma and with complete demographic and clinical data of interest. Logistic regression analyses were performed to examine associations between abnormal serum lipids, glucose, and HOMA-IR and the presence of current asthma.

Results: The data of 11,662 children (3 to 11 years of age) and 12,179 adolescents (12 to 19 years of age) were included in the analysis, including 3,703 participants with asthma and 20,138 participants without asthma. The prevalence of self-reported current asthma was higher among participants aged between 3-11 years (52.9%) than among those aged between 12-19 years (50.7%). Multivariate analyses, after adjusting for sex, race, income-to-poverty ratio, low birth weight, prenatal maternal smoking, tobacco exposure, C-reactive protein level, and body mass index z-score, revealed no associations between elevated fasting plasma glucose, reduced high-density lipoprotein cholesterol, elevated low-density lipoprotein cholesterol, total cholesterol, triglycerides, and HOMA-IR and the presence of current asthma in children or adolescents.

Conclusion: This cross-sectional study found no associations between abnormal serum lipids or HOMA-IR and the presence of current asthma in children or adolescents.

Keywords: Asthma, cholesterol, insulin resistance, lipid, lipoprotein, NHANES, obesity, wheezing

Introduction

Asthma is primarily a disease of childhood, and its increasing prevalence beginning in the 1980s has been referred to as an asthma epidemic (1). It has been estimated that the prevalence of asthma in 1980 was 3.6%, increased to 7.5% in 1995, and further increased to 9.3% in 2010. Since 2010, the overall prevalence of childhood asthma has remained unchanged or has decreased slightly (2).

While a number of risk factors are associated with the development of childhood asthma, the condition has been linked especially to obesity and metabolic syndrome (3-5). Being overweight in childhood has also been associated with an increased risk of the development of allergic disease (6). In addition, increasing attention has been given to the association between hypercholesterolemia and obesity, and obesity with airway hyper-responsiveness, suggesting a potential role of cholesterol and lipid homeostasis in lung physiology and asthma (7-14). Rastogi et al. (15) have also suggested that hyperglycemia and hyperinsulinemia may result in airway hyper-responsiveness.

Results of a number of studies have also linked elevated lipid levels with the development of asthma/wheezing in children and adults (1-5, 16). However, the results are inconsistent with those of other studies that show no association (12, 13, 17), or even a negative association, between elevated lipids and asthma/wheezing (11). The reported associations between asthma and lipid levels and insulin resistance (IR) appear to be independent of body mass index (BMI) (3). As such, dyslipidemia and hyperinsulinemia, precursors to cardiovascular disease and diabetes, may also be associated with the development of asthma and confound its epidemiologic link to obesity (3, 5, 18). The results of most studies, however, have been limited by a cross-sectional study design and a wide range of subjects studied.

Thus, the purpose of the current study was to use a national population-based database to examine the associations between lipid levels and insulin resistance and the presence of current asthma in children and adolescents.

Methods

Data Source

The United States National Health and Nutrition Examination Survey (NHANES) is an ongoing cross-sectional

health survey that represents the non-institutionalized population of the United States. The program uses a complex, multistage design to collect and analyze data representative of different geographic regions. The NHANES data are collected through a combination of interviews and physical examinations of participants by highly trained personnel. The survey is administered by the National Center for Health Statistics of the Centers for Disease Control and Prevention. Further information about the NHANES program is available at the NHANES website: https://www.cdc.gov/Nchs/Nhanes/about_nhanes.htm. Detailed information about NHANES data collection methods is available at https://wwwn.cdc.gov/Nchs/Nhanes/2003-2004/L13_C.htm and https://wwwn.cdc.gov/Nchs/Nhanes/2003-2004/L13AM_C.htm.

The survey protocol and data collection methods were approved by the NHANES Institutional Review Board (IRB), and the NCHS Research Ethics Review Board (ERB) (Protocol#98-12, Protocol#2005-06, and Protocol #2011-17). All of the NHANES data are de-identified and analysis of the data by independent researchers does not require IRB approval or subject informed consent.

Study population

Data from 7 cycles of the NHANES conducted during the period 1999-2012 were used. The data of children (aged 3-11 years) and adolescents (aged 12-19 years) with complete demographic and laboratory data, as well as that of other variables of interest, were included in the analysis. Exclusion criteria were: 1) diagnosis of diabetes mellitus (defined as a self-report of having been told by a doctor or health professional subject had diabetes or sugar diabetes, or currently taking diabetic pills or insulin; 2) pregnant; 3) underweight, defined as body mass index (BMI) <5th percentile for their age and sex (19, 20).

Dependent variables (Y)

The primary outcome of the analysis was the presence of current asthma or wheezing (21). Current asthma was defined as those who reported ever being told that they had asthma and who had an asthma attack in the past year (https://wwwn.cdc.gov/Nchs/Nhanes/2007-2008/MCQ_E.htm). Wheezing was defined as wheezing or whistling in the chest in the past year (https://wwwn.cdc.gov/Nchs/Nhanes/2003-2004/RDQ_C.htm).

Independent variables (X)

The NHANES dataset provided laboratory results of total cholesterol (TC), triglycerides (TG), high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, and fasting plasma glucose (FPG) levels. Of these, TG, LDL cholesterol and FPG were measured only in subsamples (adolescents), while TC and HDL cholesterol were measured in all participants. Homeostatic model assessment-insulin resistance (HOMA-IR) was calculated using the equation: (fasting glucose (mg/dL) × fasting insulin (pmol/L) /405 /6 (22-24) only in subsamples. The cutoff values for abnormal lipid and FPG levels, and HOMA-IR were: ≥170 mg/dL for elevated TC; ≤45 mg/dL for low HDL cholesterol; ≥110 mg/dL for elevated LDL cholesterol; ≥75 and ≥90 mg/dL for elevated TG for participants ≤9 years old and >10 years old, respectively; ≥100 mg/dL for abnormal FPG; ≥3.0 for abnormal HOMA-IR, as reported by the expert panel of the US National Heart, Lung, and Blood Institute (25) and used in prior investigations (15, 26).

Covariates (potential confounders)

Demographic data examined as potential confounders included age, sex, family income-to-poverty ratio, prenatal maternal smoking, birth weight (low birth weight or not), and C-reactive protein (CRP) level. Age- and sex-specific BMI percentiles and BMI Z-scores were determined according to the 2000 Centers for Disease Control (CDC) growth charts using a CDC SAS program www.cdc.gov/nccdphp/dnpao/growthcharts/resources/sas.htm (6).

Tobacco exposure was defined by a “yes” response to the questions: “Have you ever tried cigarette smoking,

even 1 or 2 puffs?” or “Does anyone who lives here smoke cigarettes, cigars, or pipes anywhere inside this home?”

To estimate physical activity before the year 2007, we summed the product of weekly time spent in each activity reported by the participant multiplied by the metabolic equivalent of task (MET) value for that activity yielding a MET-h index. One MET is the energy expenditure of 1 kcal/kg body weight per hour. For cycles after 2007, the physical activity questionnaire was changed. We estimated weekly MET-h for moderate and vigorous activities from questions asking participants about their participation in moderate and vigorous activities, the number of days per week engaged in these activities, and the number of minutes engaged in these activities on a typical day (27).

Statistical Analysis

Differences in categorical variables between participants with and without asthma were determined using the Rao–Scott chi-square test, and differences of continuous variables between groups were examined using the Complex Samples General Linear Model (CSGLM). Demographic data and baseline characteristics are expressed as mean \pm standard error for continuous variables, and unweighted counts (weighted %) for categorical variables. Univariate logistic regression analyses were performed to determine the association between serum lipids, glucose, and HOMA-IR and current asthma. Extended-model approaches were used for covariate adjustment: Model 1 = gender, race, poverty income ratio, low birth weight (children only), prenatal maternal smoking (children only), tobacco exposure, and physical activity (adolescents only); Model 2 = Model 1 + CRP; Model 3 = Model 2 + BMI-z score. Participants with missing data of any covariates were not included in the regression analyses. All analyses included NHANES Medical Examination Center (MEC) sample weight or fasting subsample weight, stratum, and primary sampling units (PSU) per recommendations from the National Center for Health Statistics (NCHS), to address oversampling, non-response, non-coverage, and to provide nationally representative estimates. All statistical assessments were 2-sided and evaluated at the 0.05 level of significance. Statistical analyses were performed using the statistical software package SPSS complex sample module version 22.0 (IBM Corp, Armonk, NY, USA)

Results

A total of 26,158 participants aged between 3 and 19 years were identified in the NHANES 1999-2012 cycle. Participants with diabetes (n=87), who were pregnant (n=115), or who had a BMI z-score less than the 5th percentile (n=2,115) were excluded from the analysis, leaving 23,841 participants as the analytic sample. This final eligible population included 20,138 participants without asthma and 3,703 participants with asthma, as shown in Table 1. The majority of participants were male (50.3% vs. 55.1%, respectively), white (58.3% vs. 58.6%, respectively), with median income-to-poverty ratio (76.1% vs. 75.2%, respectively), normal birth weight (89.4% vs. 87.3%, respectively), and without tobacco exposure (69.7% vs. 65.5%, respectively). The prevalence of current asthma was greater among participants aged between 3-11 years (52.9%) than among those aged between 12-19 years (50.7%). Using NHANES MEC sample weights, the analytic sample size (n = 23,841) was equivalent to a population-based sample size of 65,644,773 participants (55,246,119 without asthma and 10,398,654 with asthma). Significant differences were found in sex, race, low birth weight, prenatal maternal smoking, tobacco exposure, CRP level, and BMI z-score between groups ($P < 0.05$).

As shown in Table 2, TC, HDL, and non-HDL were not associated with current asthma among children aged 3-11. In all multivariate analyses, no association was found between serum lipids and asthma after adjustment for demographic characteristics and smoking (Model 1), addition of CRP level (Model 2), and of BMI z-score (Model 3) (Table 3).

Univariate logistic regression showed that lower HDL (odd ratio (OR)=1.229, 95% confidence interval (CI): 1.063 to 1.421), elevated TG (OR=1.246, 95% CI: 1.013 to 1.533) and abnormal HOMA-IR (OR=1.370, 95%CI: 1.077 to 1.742) were significantly associated with higher risk of asthma in adolescents. However, after adjusting for sex, race, poverty income ratio, tobacco exposure, and physical activity, the association between asthma and HDL (OR=1.189, 95% CI: 0.992 to 1.424), TG (OR=1.161, 95% CI: 0.908 to 1.484), and HOMA-IR (OR=1.243, 95%CI: 0.950 to 1.628) became non-significant (Table 4). Again, no significant associations between serum lipid, glucose, HOMA-IR and asthma were found (Model 2 and Model 3).

Discussion

This study sought to use the NHANES database to examine the relationships between lipids and IR with the presence of current asthma in children and adolescents. Although some associations were found in univariate analysis, after controlling for confounders, multivariate analysis found no associations between lipid levels or IR and asthma in children or adolescents. Consistent with the findings of the present cross-sectional study, two case-control studies reported no associations of asthma with lipids and IR in adults (12, 13).

The potential link between obesity, diabetes, and asthma has been referred to as “metabolic asthma” (3). One hypothesis is that the dysfunction of metabolic pathways that are present in obesity and diabetes exerts a direct and negative influence on the immune system, thereby affecting both adaptive and innate immunity and subsequently increasing the risk of asthma (3). Furthermore, an in vitro study implied that dysregulation of cholesterol transport in human airway smooth muscle cells may be important in the pathogenesis of asthma (14). Yiallourous et al. (16, 18) have performed a series of studies examining relationship between lipids and asthma in children and adolescents. In a cohort of 3,982 children from Cyprus, the authors found that low HDL cholesterol in childhood (11-12 years of age) was associated with the development of asthma in adolescence (age 15-17 years) (18). Utilizing a case-control design, Yiallourous et al. found that adolescent asthma was associated with low serum HDL cholesterol levels independent of HDL levels in childhood (16). Furthermore, in a cohort of children from Cyprus, Yiallourous et al. found that two single nucleotide polymorphisms (SNPs) in different genetic loci were associated with both wheezing and HDL cholesterol levels (28), while the association between these two SNPs and asthma remains to be investigated.

Unlike most studies that did not distinguish lipid particles of different sizes, Scichilone et al. (29) examined the associations between asthma and LDL subclasses in adults in a case-control study, and they found that asthma was associated with smaller LDL particles with proinflammatory property. In addition, Barochia et al. (30) found that serum levels of large HDL_{NMR} particles are positively correlated with forced expiratory volume in 1 second (FEV₁) in adult patients with atopic asthma in a case-control study.

Two cross-sectional studies examined associations between obesity and lipids, respectively, with asthma based on the NHANES database (11, 21). Visness et al. (21) evaluated the association between obesity and atopic and non-atopic asthma in children and adolescents (aged 2-19) using the NHANES database (1999-2006). They found that obesity was significantly associated with current asthma among children and adolescents (OR = 1.68), and the association was stronger in non-atopic (OR = 2.6) than atopic (OR = 1.34) children and adolescents (21). Moreover, Fessler et al. (11) examined 7005 participants ≥ 6 years of age who participated in the NHANES 2005 to 2006 survey, and found that serum total cholesterol and non-HDL cholesterol were negatively associated with asthma. However, the authors noted that the association was chiefly due to the strong relationship with lipid metabolism previously found in Mexican American individuals (11).

The discussion of this topic would not be complete without drawing attention to studies that *did not* find an association between lipids and asthma. Recently, Fang et al. (17) compared the lipid profiles of obese asthmatic

children with those of non-obese asthmatic children. The results showed that hypercholesterolemia and hypertriglyceridemia were not present in any of the asthmatic children, and there were no differences in apo-A1 and apo-B between any of the BMI groups, nor were there differences in LDL levels (17). In a longitudinal study that followed children from birth to 8 years of age, Murray et al. (6) reported that although being overweight was associated with increased risk of allergic disease and wheezing, the strength of the association varied with sex, age, and atopic phenotype.

There are a number of limitations to this study that potentially may have affected the results. Like most studies examining this topic, this was a cross-sectional study and therefore temporal relations and causation cannot be determined. Body composition changes during growth and hormonal expression after puberty may influence the association between adiposity and asthma, particularly among girls. The study population is restricted to non-institutionalized population in the NHANES database, which appears to cause under-representation of severe asthma patients who were hospitalized. Data of puberty were not available in the NHANES database. Due to the lack of certain other data, we were not able to control for other potential confounding factors such as poor asthma control, lung function, and diet. Inaccurate reporting or recall bias may have occurred, because NHANES surveys are based on individual or parents' interviews and questionnaires. On the other hand, data from NHANES are comprehensive and nationally representative, drawing from a large and diverse sample of participants of the population of the United States. Therefore, the findings are likely generalizable to the overall United States population.

Conclusions

The results of this population-based cross-sectional study did not find an association between lipids or IR and the presence of childhood asthma. Further study is necessary to fully understand the associations between lipids and IR and asthma.

Authorship Contribution

Mn Lu: Conception and design, analysis and interpretation of data, drafting of the manuscript, critical revision of the manuscript, final approval of the manuscript

Beirong Wu: Acquisition of data, analysis and interpretation of data, drafting of the manuscript, final approval of the manuscript

Rong Qiao: Conception and design, analysis and interpretation of data, critical revision of the manuscript, final approval of the manuscript

Haoxiang Gu: Acquisition of data, critical revision of the manuscript, final approval of the manuscript

Ying Din: Analysis and interpretation of data, drafting of the manuscript, final approval of the manuscript

Xiaoyan Dong: Acquisition of data, drafting of the manuscript, final approval of the manuscript

References

1. Akinbami LJ, Simon AE, Rossen LM. Changing trends in asthma prevalence among children. *Pediatrics* 2016;137. doi: 10.1542/peds.2015-2354.
2. Centers for Disease Control and Prevention. National Current Asthma Prevalence (2015). Available at: https://www.cdc.gov/asthma/most_recent_data.htm. Accessed January 26, 2018.
3. Perez MK, Piedimonte G. Metabolic asthma: is there a link between obesity, diabetes, and asthma? *Immunol Allergy Clin North Am* 2014;34:777-784.
4. Salome CM, Marks GB. Sex, asthma and obesity: an intimate relationship? *Clin Exp Allergy* 2011;41:6-8.
5. Fenger RV, Gonzalez-Quintela A, Linneberg A, Husemoen LL, Thuesen BH, Aadahl M, Vidal C, Skaaby T, Sainz JC, Calvo E. The relationship of serum triglycerides, serum HDL, and obesity to the risk of wheezing

in 85,555 adults *Respir Med* 2013;107:816-824.

6. Murray C, Canoy D, Buchan I, Woodstock A, Simpson A, Custovic A. Body mass index in young children and allergic disease: gender differences in a longitudinal study. *Clin Exp Allergy* 2010;41:78-85.
7. Al-Shawwa B, Al-Huniti N, Titus G, Abu-Hasan M. Hypercholesterolemia is a potential risk factor for asthma. *J Asthma* 2006;43:231-233.
8. Schäfer T, Ruhdorfer S, Weigl L, Wessner D, Heinrich J, Döring A, Wichmann HE, Ring J. Intake of unsaturated fatty acids and HDL cholesterol levels are associated with manifestations of atopy in adults. *Clin Exp Allergy* 2003;33:1360-1367.
9. Shenoi A, Kumar L, Sarihyan S, Gangully NK. High density lipoprotein cholesterol and total cholesterol in children with asthma and allergic rhinitis. *Acta Paediatr* 1992;81:150-152.
10. Enright PL, Ward BJ, Tracy RP, Lasser EC. Asthma and its association with cardiovascular disease in the elderly. The Cardiovascular Health Study Research Group. *J Asthma* 1996;33:45-53.
11. Fessler MB, Massing MW, Spruell B, Jaramillo R, Draper DW, Madenspacher JH, Arbes SJ, Calatroni A, Zeldin DC. Novel relationship of serum cholesterol with asthma and wheeze in the United States. *J Allergy Clin Immunol* 2009;124:967-974.
12. Picado C, Deulofeu R, Leonart R, Agustí M, Casals E, Quintó L, Mulla J. Lipid and protein metabolism in asthma. Effects of diet and corticosteroid therapy. *Allergy* 1999;54:569-575.
13. Erel F, Gulec M, Kartal O, Caliskaner Z, Ozturk S, Yaman H, Kurt Y, Gocgeldic E, Ors F, Karaayvaz M. Serum leptin levels and lipid profiles in patients with allergic rhinitis and mild asthma. *Allergol Immunopathol (Madr)* 2007;35:232-238.
14. Delvecchio CJ, Bilan P, Nair P, Capone JP. LXR-induced reverse cholesterol transport in human airway smooth muscle is mediated exclusively by ABCA1. *Am J Physiol Lung Cell Mol Physiol* 2008;295:L949-L957.
15. Rastogi D, Fraser S, Oh J, Huber AM, Schulman Y, Bhagtani RH, Khan ZS, Tesfa L, Hall CB, Macian F. Inflammation, metabolic dysregulation, and pulmonary function among obese urban adolescents with asthma. *Am J Respir Crit Care Med* 2015;191:149-160.
16. Yiallourous PK, Savva SC, Kolokotroni O, Dima K, Zerva A, Kouis P, Bousquet J, Middleton N. Asthma: the role of low high-density-lipoprotein cholesterol in childhood and adolescence. *Int Arch Allergy Immunol* 2014;165:91-99.
17. Fang LJ, Huang CS, Liu YC, Su YM, Wan KS. The lipid profile in obese asthmatic children compared to non-obese asthmatic children. *Allergol Immunopathol (Madr)* 2016;44:346-350.
18. Yiallourous PK, Savva SC, Kolokotroni O, Behbod B, Zeniou M, Economou M, Chadjigeorgiou C, Kourides YA, Tornaritis MJ, Lamnisis D, Middleton N, Milton DK. Low serum high-density lipoprotein cholesterol in childhood is associated with adolescent asthma. *Clin Exp Allergy* 2012;42:423-432.
19. Centers for Disease Control. Clinical Growth Charts. National Center for Health Statistics; Hyattsville, MD: 2007.
20. Krebs NF, Himes JH, Jacobson D, Nicklas TA, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. *Pediatrics* 2007;120(Suppl 4):S193-S228.
21. Visness CM, London SJ, Daniels JL, Kaufman JS, Yeatts KB, Siega-Riz AM, Calatroni A, Zeldin DC. Association of childhood obesity with atopic and nonatopic asthma: results from the National Health and Nutrition Examination Survey 1999-2006. *J Asthma* 2010;47:822-829.
22. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in

man. *Diabetologia* 1985;28:412-419.

23. Geloneze B, Repetto EM, Geloneze SR, Tambascia MA, Ermetice MN. The threshold value for insulin resistance (HOMA-IR) in an admixed population IR in the Brazilian Metabolic Syndrome Study. *Diabetes Res Clin Pract* 2006;72:219-220.
24. Sumner AE, Cowie CC. Ethnic differences in the ability of triglyceride levels to identify insulin resistance. *Atherosclerosis* 2008;196:696-703.
25. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents; National Heart, Lung, and Blood Institute. *Pediatrics*. 2011 Dec;128 Suppl 5:S213-56. doi: 10.1542/peds.2009-2107C.
26. Yin J, Li M, Xu L, Wang Y, Cheng H, Zhao X, Mi J. Insulin resistance determined by Homeostasis Model Assessment (HOMA) and associations with metabolic syndrome among Chinese children and teenagers. *Diabetol Metab Syndr* 2013;5:71. doi: 10.1186/1758-5996-5-71.
27. Janssen I, Carson V, Lee IM, Katzmarzyk PT, Blair SN. Years of life gained due to leisure-time physical activity in the U.S. *Am J Prev Med*. 2013;44:23-29.
28. Yiallourous PK, Kouis P, Kolokotroni O, Youhanna S, Savva SC, Dima K, Zerva A, Platt D, Middleton N, Zalloua P. Shared genetic variants between serum levels of high-density lipoprotein cholesterol and wheezing in a cohort of children from Cyprus. *Ital J Pediatr* 2016;42:67. doi: 10.1186/s13052-016-0276-1.
29. Scichilone N, Rizzo M, Benfante A, Catania R, Giglio RV, Nikolic D, Montalto G, Bellia V. Serum low density lipoprotein subclasses in asthma. *Respir Med* 2013;107:1866-1872.
30. Barochia AV, Kaler M, Cuento RA, Gordon EM, Weir NA, Sampson M, Fontana JR, MacDonald S, Moss J, Manganiello V, Remaley AT, Levine SJ. Serum apolipoprotein A-I and large high-density lipoprotein particles are positively correlated with FEV1 in atopic asthma. *Am J Respir Crit Care Med* 2015;191:990-1000.

Table 1. Demographic and basic characteristics of participants aged 3 to 19 years with and without asthma from NHANES 1999-2012 (unweighted n=23,841; weighted n=65,644,773^a)

	Without asthma (n=20,138)	With asthma (n=3,703)	P-value
Sex			
Male	10048 (50.3)	2033 (55.1)	<0.0001*
Female	10090 (49.7)	1670 (44.9)	
Race			
Mexican American	6412 (14.1)	764 (9.0)	
Other Hispanic	1364 (6.5)	302 (7.1)	
White	5425 (58.3)	1051 (58.6)	<0.0001*
Black	5836 (14.0)	1347 (18.3)	
Other	1301 (7.1)	239 (7.1)	
Income-to-poverty ratio [†]			
Not poor	12263 (76.1)	2267 (75.2)	0.452
Poor	6301 (23.9)	1189 (24.8)	
Low birth weight [†]			
No	13380 (89.4)	2326 (87.3)	0.014*
Yes	1776 (10.6)	409 (12.7)	
Prenatal maternal smoking [†]			
No	12955 (83.6)	2222 (80.6)	0.003*
Yes	2015 (16.4)	478 (19.4)	
Tobacco exposure [†]			
No	13668 (69.7)	2338 (65.5)	0.001*
Yes	6345 (30.3)	1354 (34.5)	
Age			
Children	9868 (52.9)	1794 (50.7)	0.061
Adolescents	10270 (47.1)	1909 (49.3)	
CRP [†]	0.16±0.005	0.19±0.014	0.004*
BMI z-score	0.53±0.013	0.74±0.025	<0.0001*
Physical activity MET score [†]	3574.7±109.8	3701.6±199.1	0.607

BMI, body mass index; CRP, C-reactive protein; MET, metabolic equivalent of task.

Data are reported as mean ± standard error, or number (weighted %).

^aMEC sample weighting.

[†]There were missing data in the specific variables.

Table 2. Univariate logistic regression analysis of associations between current asthma and serum lipids, glucose, and HOMA-IR

	Children, 3-11 years (n=11,662)			Adolescents, 12-19 years (n=12,179)		
	Proportion (%)	Prevalence of asthma (%)	Crude OR (95% CI)	Proportion (%)	Prevalence of asthma (%)	Crude OR (95%CI)
TC [†]						
<170 mg/dL	61.7	14.6	Reference	66.9	16.9	Reference
≥170 mg/dL	38.3	16.4	1.153 (0.960, 1.384)	33.1	16.5	0.970 (0.827, 1.138)
HDL [†]						
>45 mg/dL	71.4	14.8	Reference	63.8	15.7	Reference
≤45 mg/dL	28.6	16.4	1.128 (0.946, 1.315)	36.2	18.6	1.229 (1.063, 1.421)
Non-HDL [†]						
<120 mg/dL	66.7	14.5	Reference	68.1	16.5	Reference
≥120 mg/dL	33.3	16.8	1.188 (0.966, 1.461)	31.9	17.3	1.052 (0.885, 1.251)
TG*						
< 90 mg/dL				60.7	16.6	Reference
≥90 mg/dL				39.3	17.1	1.246 (1.013, 1.533)
LDL*						
<110 mg/dL				78.9	16.9	Reference
≥110 mg/dL				21.1	19.6	1.200 (0.933, 1.544)
FPG*						
<100 mg/dL				84.2	17.5	Reference
≥100 mg/dL				15.8	17.0	0.967 (0.726, 1.287)
HOMA-IR*						
≤3				66.4	15.9	Reference
>3				33.6	20.5	1.370 (1.077, 1.742)

FPG, fasting plasma glucose; HDL, high-density lipoprotein, HOMA-IR, homeostatic model assessment-insulin resistance; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglycerides.

*MEC sample weighting.

*Fasting Subsample weighting. (there were around 27% of total participants with weighting data)

Numbers in bold indicated statistical significance ($P < 0.05$).

Uncorrected proof

Table 3. Multivariate logistic regression of the association between serum lipids and asthma in children (n=11,662)

	Model 1 ^a	Model 2 ^b	Model 3 ^c
	aOR (95%CI)	aOR (95%CI)	aOR (95%CI)
TC			
≥170 mg/dL vs. <170 mg/dL	1.145 (0.938, 1.396)	1.130 (0.884, 1.445)	1.109 (0.871, 1.412)
HDL			
≤45 mg/dL vs. >45 mg/dL	1.196 (0.988, 1.447)	1.112 (0.882, 1.402)	1.051 (0.821, 1.347)
Non-HDL			
≥120 mg/dL vs. <120 mg/dL	1.224 (0.986, 1.520)	1.149 (0.873, 1.513)	1.100 (0.839, 1.443)

aOR, adjusted odds ratio; CRP, C-reactive protein; BMI, body mass index.

^a Adjusted for sex, race, income-to-poverty ratio, low birth weight, prenatal maternal smoking and tobacco exposure.

^b Adjusted for sex, race, income-to-poverty ratio, low birth weight, prenatal maternal smoking, tobacco exposure and CRP.

^c Adjusted for sex, race, income-to-poverty ratio, low birth weight, prenatal maternal smoking, tobacco exposure, CRP and BMI z-score.

MEC sample weighting.

Table 4. Multivariate logistic regression of the association between serum lipids, glucose and HOMA-IR and asthma in adolescents (n=12,179)

	Model 1 ^a	Model 2 ^b	Model 3 ^c
	aOR (95%CI)	aOR (95%CI)	aOR (95%CI)
TC [†]			
≥170 mg/dL vs. <170 mg/dL	0.957 (0.801, 1.144)	1.048 (0.864, 1.271)	1.002 (0.830, 1.210)
HDL [†]			
≤45 mg/dL vs. >45 mg/dL	1.189 (0.992, 1.424)	1.179 (0.981, 1.417)	1.065 (0.874, 1.298)
Non-HDL [†]			
≥120 mg/dL vs. <120 mg/dL	1.018 (0.828, 1.252)	1.097 (0.889, 1.355)	1.009 (0.812, 1.253)
TG*			
≥90 mg/dL vs. <90 mg/dL	1.161 (0.908, 1.484)	1.055 (0.818, 1.362)	0.933 (0.710, 1.226)
LDL*			
≥110 mg/dL vs. <120 mg/dL	1.152 (0.860, 1.542)	1.130 (0.828, 1.540)	1.049 (0.765, 1.438)
FPG*			
≥100 mg/dL vs. <100 mg/dL	0.976 (0.683, 1.393)	1.023 (0.711, 1.473)	0.964 (0.676, 1.374)
HOMA-IR*			
>3 vs. ≤3	1.243 (0.950, 1.628)	1.140 (0.861, 1.508)	0.891 (0.662, 1.198)

aOR, adjusted odds ratio; FPG, fasting plasma glucose; HDL, high-density lipoprotein, HOMA-IR, homeostatic model assessment-insulin resistance; LDL, low-density lipoprotein; TC, total cholesterol; TG, triglycerides.

^a Adjusted for sex, race, income-to-poverty ratio, low birth weight, prenatal maternal smoking and tobacco exposure.

^b Adjusted for sex, race, income-to-poverty ratio, low birth weight, prenatal maternal smoking, tobacco exposure and CRP.

^c Adjusted for sex, race, income-to-poverty ratio, low birth weight, prenatal maternal smoking, tobacco exposure, CRP and BMI z-score.

[†]MEC sample weighting.

*Fasting Subsample weighting. (there were around 27% of total participants with weighting data)