



















26. Bilati U, Allémann E, Doelker E. Sonication parameters for the preparation of biodegradable nanocapsules of controlled size by the double emulsion method. *Pharmaceutical development and technology*. 2003;8:1-9.
27. Cun D, Foged C, Yang M, Frøkjær S, Nielsen HM. Preparation and characterization of poly (DL-lactide-co-glycolide) nanoparticles for siRNA delivery. *International journal of pharmaceutics*. 2010;390:70-75.
28. Fonte P, Soares S, Costa A, Andrade JC, Seabra V, Reis S, Sarmiento B. Effect of cryoprotectants on the porosity and stability of insulin-loaded PLGA nanoparticles after freeze-drying. *Biomatter*. 2012;2:329-339.
29. De Chasteigner S, Cavé G, Fessi H, Devissaguet JP, Puisieux F. Freeze-drying of itraconazole-loaded nanosphere suspensions: a feasibility study. *Drug development research*. 1996;38:116-124.
30. Panyam J, Williams D, Dash A, Leslie-Pelecky D, Labhasetwar V. Solid-state solubility influences encapsulation and release of hydrophobic drugs from PLGA/PLA nanoparticles. *J Pharm Sci*. 2004;93:1804-14.
31. Chella N, Narra N, Rama Rao T. Preparation and characterization of liquisolid compacts for improved dissolution of telmisartan. *Journal of drug delivery*. 2014;2014:692793.
32. Kociecka B., Surazynski A., Milyk W., Palka J. The effect of Telmisartan on collagen biosynthesis depends on the status of estrogen activation in breast cancer cells. *Eur J Pharmacol*. 2010;628: 51-56.

Uncorrected proof