



Evaluation of Long-term Intensive Care Patients in a Cardiac Surgery Center

Kalp Cerrahisi Merkezinde Uzayan Yoğun Bakım Hastalarının Değerlendirilmesi

¹ Bedih Balkan¹, ² Hatice Dilek Özcanoglu²

¹Istanbul Provincial Directorate of Health, Mehmet Akif Ersoy Chest and Cardiovascular Diseases Hospital, Clinic of Intensive Care Unit, Istanbul, Turkey

²Istanbul Provincial Directorate of Health, Mehmet Akif Ersoy Chest and Cardiovascular Diseases Hospital, Clinic of Anesthesiology, Istanbul, Turkey

Abstract

Objective: This study was aimed to evaluate the clinical and demographic characteristics of patients in a chronic intensive care unit (ICU) in a cardiac surgery center and to research the effect of these characteristics on scoring systems used.

Method: From 1 September 2014 to 1 September 2019, the study was performed retrospectively with cases admitted to the chronic ICU due to lengthened intensive care requirements after monitoring in the cardiovascular surgery intensive care or coronary ICU. Each case had study forms including information like age, gender, comorbid diseases, acute physiology and chronic health evaluation 2 (APACHE 2) scores, Glasgow Coma scale, the results were statistically assessed.

Results: Two hundred and sixty three patients were enrolled for this study. The mean age of patients was 64.6±6 years, with 61% male and 39% female. Of cases, 46% had coronary bypass surgery (n=122), 25% had valve + aorta dissection (n=67), 19% had acute coronary artery (n=51) and 7% had peripheral artery disease (n=17). The mean follow-up duration for patients was identified as 44.2±96.38 days. Cases developing mortality had significantly higher APACHE 2 scores compared to cases without mortality (p<0.05, 24.9±8.0 vs 20.5±7.2). The most common comorbidities in cases were kidney failure in 41% (n=48). Sixty-nine cases had a percutaneous tracheostomy and 21 cases had a surgical tracheostomy, for a total of 90 cases with a tracheostomy.

The mortality rate of entire monitored patients during this study period was 42.6%.

Conclusion: Very different problems may be encountered in cases observed in the chronic ICU. For these cases, kidney failure is the most

Öz

Amaç: Bu çalışmada kalp cerrahisi merkezinde, kronik yoğun bakım ünitesinde yatan hastaların klinik ve demografik özelliklerinin değerlendirilmesi ve bu özelliklerin kullanılan skorlama sistemlerine olan etkisinin araştırılması amaçlandı.

Yöntem: 1 Eylül 2014-1 Eylül 2019 tarihleri arasında retrospektif olarak kardiyovasküler cerrahi yoğun bakım veya koroner yoğun bakım ünitesinde izlenip yoğun bakım sürecinin uzaması nedeniyle kronik yoğun bakım ünitesine alınan olgularda gerçekleştirildi. Her bir olgunun yaş, cinsiyet, eşlik eden hastalık, Akut Fizyoloji ve Kronik Sağlık Değerlendirmesi 2 (APACHE 2) skorları, Glasgow Koma skalası, bilgileri içeren çalışma formu oluşturuldu. Sonuçlar istatistiksel olarak değerlendirildi.

Bulgular: İki yüz altmış üç hasta çalışmaya alınmıştır. Hastaların ortalama yaşları 64,6±6 yıl olup, %61'i erkek %39'u kadındı. Olguların %46'una koroner bypass (n=122) cerrahisi, %25'ine kapak + aort diseksiyonu (n=67) %19 akut koroner arter sonrası (n=51) %7'si periferik arter hastalığı (n=17) sonrası izlenmişti. Hastaların takip süresi ortalama 44,2± 96,38 gün olarak saptanmıştı. Mortalite gelişen olgularda, mortalite gelişmeyen olgulara göre APACHE 2 skoru anlamlı olarak daha yüksekti (p<0,05 24,9±8,0 vs 20,5±7,2). Olgularda en sık gözlenen komorbidite %41 (n=48) Böbrek yetmezliği idi. Altmış dokuz olguya perkutan trakeostomi ve 21 olguya cerrahi trakeostomi olmak üzere 90 olguya trakeostomi açıldı. Toplam izlenen hastaların %42,6'si (n=112) kaybedildi.

Sonuç: Kronik yoğun bakımda izlenen olgular çok farklı sorunlarla karşılaşılabilir. Bu olgularda böbrek yetmezliği mortaliteyi ve morbiditeyi artıran en önemli risk faktörüdür. Ayrıca hastalarda APACHE 2 skorlaması



Address for Correspondence: Bedih Balkan, İstanbul Provincial Directorate of Health, Mehmet Akif Ersoy Chest and Cardiovascular Diseases Hospital, Intensive Care Unit, İstanbul, Turkey

E-mail: drbedihbalkan21@gmail.com **ORCID:** orcid.org/0000-0003-3510-6991 **Received:** 30.11.2019 **Accepted:** 28.12.2019

Cite this article as: Balkan B, Özcanoglu HD. Evaluation of Long-term Intensive Care Patients in a Cardiac Surgery Center. Bagcilar Med Bull 2020;5(1):13-20

©Copyright 2020 by the Health Sciences University, Bagcilar Training and Research Hospital
Bagcilar Medical Bulletin published by Galenos Publishing House.

Abstract

important risk fact increasing mortality and morbidity. Additionally, APACHE 2 scoring of patients may be helpful for the assessment and prediction of operative mortality and morbidity.

Keywords: Chronic intensive care, cardiac center, APACHE 2, prolonged mechanic ventilation

Öz

operatif mortalite ve morbiditeyi deęerlendirmede ve öngörmeye yardımcı olabilir.

Anahtar kelimeler: Uzayan yoğun bakım hastaları, kalp merkezi, APACHE 2, uzamış mekanik ventilasyon

Introduction

Currently, with the development of technology in cardiac centers, both invasive cardiac procedures and open heart surgery operations are successfully performed for geriatric cases with higher risks. This situation has led to longer intensive care and hospital stays. Studies have stated that nearly 19-45% of cases have lengthened intensive care duration after open-heart surgery (1,2). For lengthened intensive care duration, a variety of risk factors may be listed such as advanced age, chronic obstructive pulmonary disease (COPD), renal failure or dysfunction, atrial fibrillation, low ejection fraction, emergency surgery, previous history of cardiac surgery and inotrope support (3).

As the intensive care duration lengthens for cases, additional problems that may increase mortality and morbidity may develop and support treatments are required to resolve these problems. The most important of these are tracheostomy performed in cases requiring a longer duration of mechanical ventilation support to reduce complications of endotracheal intubation and mechanical ventilation to a minimum and ensure patient comfort, and renal replacement treatments to control kidney failure.

There are many factors affecting mortality in intensive care units (ICU). By noting these factors, a variety of scoring systems were developed to determine the patient's prognosis. The Acute Physiology and Chronic Health Evaluation (APACHE) is one of the most commonly used systems with this aim. The most important advantages of these scoring systems are the creation of common use, reliable database, lowering costs in intensive care, effective use of resources, assisting in clinical decisions and applications and the opportunity for objective assessment. However, some studies have proposed that these scoring systems are insufficient to estimate the prognosis for patients admitted to intensive care for long durations (4).

Prolonged mechanical ventilation (PMV) is a common problem, with a reported incidence between 2.9% and 8.6%

(5) after coronary artery bypass grafting (CABG). The PMV patients were older, sicker, had more complex operations and had higher rates of postoperative morbidity.

The primary aim of our study is to investigate the clinical and demographic characteristics of patients admitted to a chronic intensive care center and report our five-year experience. Our secondary aim to investigate the correlation of patient APACHE 2 and Glasgow Coma scale (GCS) scores with mortality and our final aim is to evaluate the additional support treatments.

Material and Methods

This study was retrospectively performed on cases admitted to the chronic ICU due to lengthened intensive care durations in the cardiovascular surgery intensive care or coronary ICU from 1 September 2014 to 1 September 2019. After receiving Ethics Committee permission from the Ethics Committee of İstanbul Provincial Directorate of Health, Mehmet Akif Ersoy Chest and Cardiovascular Diseases Hospital (decision no: 59, date: 10/9/2019), the patient information for patients admitted to chronic intensive care were investigated from admission files and the hospital electronic records system.

Study forms containing information about the age, gender, main diagnosis, intensive care stay duration, problems developing in intensive care, discharge status, etc. of patients were created. Patients with missing information or who stayed in chronic intensive care for less than twenty-four hours were not included in the study.

Additionally, the APACHE 2 and GCS used during the monitoring and treatment of every patient in intensive care were evaluated.

COPD was defined as those patients with a COPD diagnosis or patients who required bronchodilator treatment before surgery. All patients, except for the emergency cases, were examined by the pulmonologist before the operation as a routine workup.

Chronic renal dysfunction (CRD) denoted the patients who had serum creatinine levels above the normal (0.5-1.3 mg/dL).

The primary clinical endpoint of this study was to evaluate the weaning failure by 7 days.

Was to evaluate the weaning failure by 7 days after surgery. The pulmonary complication was defined as any pulmonary abnormality occurring in the postoperative period that produces identifiable disease or dysfunction that is clinically significant and that adversely affects clinical course such as Postoperative Respiratory Distress syndrome, reintubation, presence of pneumothorax or pulmonary effusion but not, PMV (6).

The neurologic status of the patients was examined daily: the daily neurologic status of the patients were performed neurologic complication was defined as the presence of any cerebrovascular event or transient neurologic dysfunction.

Infectious complications were defined as positive blood, urine, sputum or wound cultures postoperatively, requiring dressings and intravenous antibiotics, requiring revision surgery (like mediastinal infection) or presence of radiographic infiltrates.

The gastrointestinal complication was defined as the presence of 1 or more of the following (7). Hematemesis or melena (with a 42 g decrease in hemoglobin level).

Statistical Analysis

SPSS 22 was used for analyzing data. The distribution of variables was measured with the Kolmogorov-Smirnov test. Descriptive statistics used mean, standard deviation, median, minimum, maximum, frequency and percentage values. Quantitative independent data analysis used the Mann-Whitney U test. Qualitative data analysis used the chi-square test. The effect level was researched with univariate and multivariate logistic regression. $P < 0.05$ was accepted as significant.

Results

Two hundred sixty three patients (61% male, 39% female) were enrolled in the study. The mean age of patients was 64.6 ± 6 years. Of patients, 61% were male and 39% were female. In terms of age, 134 (51%) were under 65 years and 129 (49%) were over 65 years of age. The mean duration of stay in intensive care was 34.6 ± 68 days. Of cases, 46% had coronary bypass surgery ($n=122$), 25% had valve + aorta dissection ($n=67$), 19% had acute coronary artery ($n=51$) and 7% had peripheral artery disease ($n=17$).

The demographic characteristics of the cases are summarized in Table 1.

The mortality rate of entire monitored patients during this study period was 42.6%. When the reasons for mortality of cases are investigated, the most frequently observed comorbidity was kidney failure at 41% ($n=48$). Patients with high APACHE scores and low GCS scores were found and have significantly higher mortality rates.

The mortality status of cases and Influencing factors are summarized in Table 2.

The univariate model observed significant effects of APACHE 2 score, APACHE 2 mortality % rates, GCS score and renal failure for prediction of mortality ($p < 0.05$). When the efficacy of these variables was tested in the multivariate model, APACHE 2 expected mortality rate % and kidney failure were independently significant for prediction of mortality (Table 3). The reasons for which patients were followed up in the ICU (Table 4).

Discussion

There is limited data related to a long duration of admission to intensive care for patients in Turkey. Agencel et al. (8) reported 9.3% of patients admitted to a tertiary ICU stayed for 21 days or longer. Martini et al. (9) stated that surgical critical care patients stayed for a mean of 116 days. However, the ICU duration of stay for patients with respiratory diseases was identified as 24.17 ± 8.38 days (10). In our study, the surviving group of patients was admitted for 27.5 ± 33.4 days, while the patients in the mortality group were monitored in intensive care for mean 44.2 ± 96.38 days (Table 1).

As in the whole world, the elderly population is increasing in Turkey. Recent census counts in our country identified the elderly population rate as 8.3% (11). The elderly population has 4 times higher surgical requirements compared to the remaining population. In the elderly patient group, standard mortality rates for those undergoing operations were identified to be 87% high compared to unoperated peers compared to data in the national mortality information bank (12). Ursavas et al. (10) in a study of a respiratory ICU did not identify a significant difference between the mean ages of surviving and mortal patients. Similarly, in our study, the mortality rates for patients aged over 65 years, comprising 49% of the group admitted to intensive care, were not different compared to those under 65 years of age (Table 2).

According to gender, 60.1% of our patients were male and 39.9% were female. Differences linked to gender have been discussed in previous studies; however, it is uncertain whether gender is a predictive factor for clinical outcomes or not. Research in Sweden revealed 60% of patients in ICU were male; however, disease severity was similar to females (13). Our study shows similarities to this study. In our study, gender was not a factor providing clues about the long duration of stay. It is not considered that gender is associated with high mortality rates (Table 2).

Mortality after cardiac surgery is still a controversial topic in spite of surgical techniques developed and anesthesia management. A variety of studies found mortality after cardiac surgery was from 2.94 to 30.7% (14,15). In our study, mortality was 42.6%. The reason for this may be linked to the inclusion of those admitted to chronic intensive care after cardiovascular surgery and patients with stay

lengthened due to hypoxia or who could not be weaned from mechanical ventilation. Diagnoses for patients admitted to intensive care are various, with some patients having more than one diagnosis. The most common cause in the mortality group was cardiopulmonary arrest occurring due to any factor. In our retrospective study, the patient data not being recorded in detail on admission or cardiopulmonary arrests without explained cause may have led to this general diagnosis being most frequent in the mortality group.

Doerr et al. (16) in a 2801 patient study of APACHE 2 reported it showed good performance in terms of calibration and differentiation statistics. Argyriou et al. (17) reported the APACHE 2 score had a good prediction of outcomes in the cardiac ICU and had comparable differentiative capabilities. Kaniş et al. (18) identified the APACHE 2 score for the surviving group in the ICU was 20.5 ± 7.28 , while it was 24.9 ± 6.22 in the mortality group. They reported the

Table 1. Demographic data

		Minimum-maximum		Median	Av.± SD/n-%	
		23.0	86.0	65.0	64.4	12.6
Age	≤65	-	-	-	134	51.0%
	>65	-	-	-	129	49.0%
Gender	Female	-	-	-	105	39.9%
	Male	-	-	-	158	60.1%
Follow-up duration (day)		1.0	685.0	16.0	34.6	68.4
APACHE 2		7.0	50.0	22.0	22.4	7.9
APACHE mortality (%)		8.0	97.0	42.4	44.1	22.8
GCS		0.0	15.0	11.0	10.5	3.9
Valve + Aorta dissection	(-)	-	-	-	196	74.5%
	(+)	-	-	-	67	25.5%
Bypass	(-)	-	-	-	141	53.6%
	(+)	-	-	-	122	46.4%
Periphery + carotid + abdominal aneurysm	(-)	-	-	-	246	93.5%
	(+)	-	-	-	17	6.5%
Coronary + emergency + COPD	(-)	-	-	-	212	80.6%
	(+)	-	-	-	51	19.4%
Kidney failure	(-)	-	-	-	126	47.9%
	(+)	-	-	-	107	40.7%
Percutaneous tracheostomy	(-)	-	-	-	194	73.8%
	(+)	-	-	-	69	26.2%
Surgical tracheostomy	(-)	-	-	-	240	91.3%
	(+)	-	-	-	23	8.7%
Mortality	(-)	-	-	-	151	57.4%
	(+)	-	-	-	112	42.6%

GCS: Glasgow Coma scale, COPD: Chronic obstructive pulmonary disease, Av: Average, SD: Standard deviation, APACHE: Acute physiology and chronic health evaluation, n: Number

APACHE 2 score was significantly higher for patients in the mortality group. In our study, the mean APACHE 2 scores for patients who died were 24.9±8.0, while it was 20.5±7.2 for surviving patients. Our results are similar to the studies above.

There are a variety of situations limiting the use of another prognostic parameter (19) of the Glasgow Coma scale. Long mechanical ventilation duration and postoperative sedation may be given as examples (20). The values of postoperative scoring systems may be affected by electrolyte and blood glucose imbalance, long mechanical ventilation duration, sedation after open-heart surgery and especially the use of cardiopulmonary bypass (21). As a result, the GCS scoring may not be very effective. In our study, the GCS in the surviving group was 11.4±3.7, while it was 9.43±8 for the mortality group and this was statistically significant (p<0.05).

In the postoperative period (22), the use of intra-aortic balloon pumps and ventricular support devices, and Low Cardiac Output syndrome are significant parameters affecting patient outcomes. Unfortunately, most scoring systems ignore these parameters (23). As a result, mortality rates measured with cardiac postoperative scoring systems may be roughly compatible but insufficient. It may be more appropriate to add a cardiac scoring system to the APACHE 2 and GCS scores.

Mortality due to cardiac reasons in CRF patients is reported from 40-50%. This rate is nearly 20 times the risk for people without renal failure (24). Open heart surgery mortality for chronic renal failure (CRF) patients is higher compared to patients with normal kidney functions (25,26). In our study, the renal failure rate was significantly higher in the mortality group compared to the surviving group (p<0.05) (Table 2).

Table 2. Analysis

		Mortality (-)		Mortality (+)		p		
		Av.± SD/n-%	Median	Av.± SD/n-%	Median			
		64.6±12.1	67.0	64.2	13.2	64.0	0.963 ^m	
Age	≤65	74	49.0%	-	60	53.6%	-	0.464 ^{χ²}
	>65	77	51.0%	-	52	46.4%	-	
Gender	Female	65	43.0%	-	40	35.7%	-	0.230 ^{χ²}
	Male	86	57.0%	-	72	64.3%	-	
Follow-up duration (days)		27.5	33.4	15.0	44.2	96.8	17.5	0.132 ^m
APACHE2		20.5	7.2	21.0	24.9	8.0	25.0	0.000 ^m
APACHE mortality (%)		35.6	19.0	34.0	55.3	22.6	57.0	0.000 ^m
GCS		11.4	3.7	13.0	9.4	3.8	10.0	0.000 ^m
Valve + Aorta dissection	(-)	110	72.8%	-	86	76.8%	-	0.469 ^{χ²}
	(+)	41	27.2%	-	26	23.2%	-	
Bypass	(-)	82	54.3%	-	59	52.7%	-	0.794 ^{χ²}
	(+)	69	45.7%	-	53	47.3%	-	
Peripheral + carotid + abdominal aneurysm	(-)	144	95.4%	-	102	91.1%	-	0.162 ^{χ²}
	(+)	7	4.6%	-	10	8.9%	-	
Coronary + emergency + COPD	(-)	118	78.1%	-	94	83.9%	-	0.241 ^{χ²}
	(+)	33	21.9%	-	18	16.1%	-	
Kidney failure	(-)	91	60.3%	-	35	31.3%	-	0.000 ^{χ²}
	(+)	41	27.2%	-	66	58.9%	-	
Percutaneous tracheostomy	(-)	103	68.2%	-	91	81.3%	-	0.017 ^{χ²}
	(+)	48	31.8%	-	21	18.8%	-	
Surgical tracheostomy	(-)	144	95.4%	-	96	85.7%	-	0.006 ^{χ²}
	(+)	7	4.6%	-	16	14.3%	-	

GCS: Glasgow Coma scale, COPD: Chronic Obstructive Pulmonary disease, Av: Average, SD: Standard deviation, APACHE: Acute physiology and chronic health evaluation, n: Number, ^mMann-Whitney U test/^{χ²}: chi-square test

Table 3. Regression data

	Univariate model				Multivariate model					
	OR	95% CI		p	OR	95% CI		p		
APACHE 2	1.08	1.04	-	1.12	0.000	-	-	-	-	
APACHE mortalite (%)	1.04	1.03	-	1.06	0.000	1.04	1.02	-	1.05	0.000
GCS	0.88	0.82	-	0.94	0.000	-	-	-	-	-
Kidney failure	4.19	2.41	-	7.26	0.000	3.25	1.70	-	6.21	0.000
Percutaneous tracheostomy	0.50	0.28	-	0.89	0.019	0.35	0.17	-	0.73	0.005
Surgical tracheostomy	3.43	1.36	-	8.65	0.009	2.97	1.02	-	8.68	0.046

OR: Odds ratio, CI: Confidence interval, GCS: Glasgow Coma scale, APACHE: Acute physiology and chronic health evaluation

Table 4. Indicators of intensive care

	Number of the patients	Percentage
Chronic renal failure	62	23.50%
Chronic obstructive pulmonary disease	32	12.16%
Hypoxic brain injury	18	6.84%
Sepsis	42	15.96%
Mediastinitis	18	6.84%
Advanced right heart failure	25	9.50%
Stroke + subarachnoid hemorrhage	17	6.46%
Left heart failure + hemodynamic instability	15	5.70%
Gastrointestinal complication	4	1.52%
Acute Respiratory Distress syndrome	22	8.36%
Peripheral arterial disease	8	3.04%

Tracheostomy is a surgical method frequently applied to patients with long term mechanical ventilation, for treatment of upper respiratory tract obstruction, and those with endotracheal intubation in ICU and predicted to remain linked to a mechanical ventilator for a long period (27,28). Percutaneous tracheostomy (PT) is easily performed, has low complication rates and can be performed in a short period at the patient's bedside so it is mostly chosen in recent years and is frequently used in the ICU for cases with elective tracheostomy especially (29,30). During the PT procedure, complications like hemorrhage, hypoxia, hypercapnia, pneumothorax, pneumomediastinum, subdermal emphysema, paratracheal insertion, tracheal wall injury, aspiration, sudden death, and esophagus injury may be observed (29). In our study, 21 (23%) surgical tracheostomy and 69 (77%) PT was performed in 263 patients hospitalized in the ICU in the last 5 years.

In studies, rates for minor hemorrhage linked to the PT procedure are 1.51-5.2% (28,31,32), while rates for major hemorrhage are 0.75-2.6% (28). In our study, after the PT procedure, 10 (14.4%) had minor hemorrhage and

two (1.9%) had a moderate hemorrhage. For the surgical procedure, 10 (76.8%) had minor hemorrhage and five (23.8%) had a moderate hemorrhage. Major hemorrhage was not observed. The reason for the high incidence of minor hemorrhage is considered to be the use of anticoagulant medication by our patients. Only one case developed pneumothorax linked to percutaneous tracheostomy.

Coronary artery bypass surgery comprises a significant portion of everyday cardiac surgery practice; about 90% for many cardiac surgery centers. PMV is still a significant reason for postoperative morbidity and mortality. Apart from the patient-related suffering, these patients cause a significant increase in the workload of postoperative ICU. Postoperative prolonged ventilation was associated with advanced, CRD, and longer perfusion times in CABG patients (33).

Study Limitations

There are some limitations to this study. The first is that our study took place in a single-center, so it may not reflect all chronic intensive care patients in Turkey. The second is that it was a retrospective study. There is a need for multicenter studies to further confirm our findings.

Conclusion

Very different problems may be encountered in cases observed in the chronic ICU. For these cases, kidney failure is the most important risk fact increasing mortality and morbidity. Additionally, APACHE 2 scoring of patients may be helpful for the assessment and prediction of operative mortality and morbidity.

Ethics

Ethics Committee Approval: Ethics Committee permission from the Ethics Committee of İstanbul Provincial Directorate of Health, Mehmet Akif Ersoy Chest and Cardiovascular Diseases Hospital (decision no: 59, date: 10/9/2019).

Informed Consent: Because the study is retrospective study, patient consent was not obtained.

Authorship Contributions

Concept: B.B., H.D.Ö., Design: B.B., H.D.Ö., Data Collection or Processing: B.B., Analysis or Interpretation: H.D.Ö., Literature Search: B.B., H.D.Ö., Writing: B.B.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Van Mastrigt GA, Heijmans J, Severens JL, Fransen EJ, Roekaerts P, Voss G, et al. Short stay intensive care after coronary artery bypass surgery: randomized clinical trial on safety and cost-effectiveness. *Crit Care Med* 2006;34:65-75.
2. De Cocker J, Messaoudi N, Stockman BA, Bossaert LL, Rodrigus IE. Preoperative prediction of intensive care unit stay following cardiac surgery. *Eur J Cardiothorac Surg* 2011;39:60-67.
3. Almashrafi A, Elmontsri M, Aylin P. Systematic review of factors influencing length of stay in ICU after adult cardiac surgery. *BMC Health Serv Res* 2016;16:318.
4. Patel PA, Grant BJB. Application of mortality prediction systems to individual intensive care units. *Intensive Care Med* 1999;25:977-982.
5. Saleh HZ, Shaw M, Al-Rawi O, Yates J, Pullan DM, Chalmers JA, et al. Outcomes and predictors of prolonged ventilation in patients undergoing elective coronary surgery. *Interact Cardiovasc Thorac Surg* 2012;15:51-56.
6. Hulzebos EH, Van Meeteren NL, De Bie RA, Dagnelie PC, Helder PJ. Prediction of postoperative pulmonary complications on the basis of preoperative risk factors in patients who had undergone coronary artery bypass graft surgery. *Phys Ther* 2003;83:8-16.
7. Bolcal C, Iyem H, Sargin M, Mataraci I, Sahin MA, Temizkan V, et al. Gastrointestinal complications after cardiopulmonary bypass: Sixteen years of experience. *Can J Gastroenterol* 2005;19:613-617.
8. Aygencel G, Türkoğlu M. Characteristics, outcomes and cost of prolonged stay ICU patients. *Yoğun Bakım Derg* 2011;3:53-58.
9. Martini V, Lederer AK, Laessle C, Makowicz F, Utzolino S, Fichtner-Feigl S, et al. Clinical Characteristics and Outcomes of Surgical Patients with Intensive Care Unit Lengths of Stay of 90 Days and Greater. *Crit Care Res Pract* 2017;2017:9852017.
10. Ursavas A, Ege E, Yüksel EG, Atabey M, Coşkun F, Yıldız F, et al. Solunumsal yoğun bakım ünitesinde mortaliteyi etkileyen faktörlerin değerlendirilmesi. *Yoğun Bakım Dergisi* 2006;6:43-48.
11. Yaşlı nüfus sayımı, Türkiye İstatistik Kurumu verileri, 2016. Erişim adresi: <http://www.tuik.gov.tr/PreHaberBultenleri.do?id=24644>.
12. Naughton C, Feneck RO. The impact of age on 6-month survival in patients with cardiovascular risk factors undergoing elective non-cardiac surgery. *Int J Clin Pract* 2007;61:768-776.
13. Banck M, Walther S, Karlstrom G, Nolin T, Sjöberg F, Samuelsson C. [There are more male than female patients within intensive care. But it is still unclear whether there are gender inequalities in Swedish intensive care or not]. *Lakartidningen* 2014;111:388-390.
14. Oprea AD, Del Rio JM, Cooter M, Green CL, Karhausen JA, Nailer P, et al. Pre- and postoperative anemia, acute kidney injury, and mortality after coronary artery bypass grafting surgery: a retrospective observational study. *Canadian J Anesth* 2018;65:46-59.
15. Siregar S, Groenwold RHH, de Mol BAJM, Speekenbrink RGH, Versteegh MIM, Bruinsma GJBB, et al. Evaluation of cardiac surgery mortality rates: 30-day mortality or longer follow-up? *Europ J Cardio-Thorac Surg* 2013;44:875-883.
16. Doerr F, Badreldin AM, Heldwein MB, Bossert T, Richter M, Lehmann T, et al. A comparative study of four intensive care outcome prediction models in cardiac surgery patients. *J Cardiothorac Surg* 2011;6:1-8.
17. Argyriou G, Vrettou C, Filippatos G, Sainis G, Nanas S, Routsis C. Comparative evaluation of Acute Physiology and Chronic Health Evaluation II and Sequential Organ Failure Assessment scoring systems in patients admitted to the cardiac intensive care unit. *J Crit Care* 2015;30:752-757.
18. Kamiş A, İşçimen R, Kelebek Girgin N, Yavaşcaoglu B, Kaya FN, Kahveci F, et al. Yoğun bakım hastalarında mikroalbuminürinin prognostik değeri. *Uludağ Üniversitesi Tıp Fakültesi Dergisi* 2008;34:107-114.
19. Marik PE, Varon J. Severity scoring and outcome assessment. Computerized predictive models and scoring systems. *Crit Care Clin* 1999;15:633-646.
20. Rady MY, Ryan T, Starr NJ. Perioperative determinants of morbidity and mortality in elderly patients undergoing cardiac surgery. *Crit Care Med* 1998;26:225-235.
21. Weiss YG, Merin G, Koganov E, Ribo A, Oppenheim-Eden A, Medalion B, et al. Postcardiopulmonary bypass hypoxemia: a prospective study on incidence, risk factors, and clinical significance. *J Cardiothorac Vasc Anesth* 2000;14:506-513.
22. Higgins TL, Estafanous FG, Loop FD, Beck GJ, Lee JC, Starr NJ, et al. ICU admission score for predicting morbidity and mortality risk after coronary artery bypass grafting. *Ann Thorac Surg* 1997;64:1050-1058.
23. Doerr F, Badreldin A, Heldwein M, Bossert T, Richter M, Lehmann T, et al. A comparative study of four intensive care outcome prediction models in cardiac surgery patients. *J Cardiothorac Surg* 2011;6:1-8.
24. Amann K, Tyralla K, Gross ML, Eifert T, Adamczak M, Ritz E. Special characteristics of atherosclerosis in chronic renal failure. *Clin Nephrol* 2003;60:S13-21.
25. Ko W, Kreiger KH, İsom OW. Cardiopulmonary bypass procedures in dialysis patients. *Ann Thorac Surg* 1993;55:677-684.
26. Bhattacharyya N, Cheung AH, Dang CR, Wong LL, Myers SA, Ng RC, et al. Open heart surgery in patients with end-stage renal disease. *Am J Nephrol* 1997;17:435.
27. Pappas S, Maragoudakis P, Vlastarakos P, Assimakopoulos D, Mandrali T, Kandilotos D, et al. Surgical versus percutaneous tracheostomy: an evidence-based approach. *Eur Arch Otorhinolaryngol* 2011;268:323-330.
28. Düger C, İsbir AC, Uysal İÖ, Kol İÖ, Kaygusuz K, Gürsoy S, et al. The evaluation of the complications of surgical and percutaneous tracheostomies in intensive care unit. *Turk J Anaesthesiol Reanim* 2013;41:84-87.

29. Ak K. Perkütan Trakeostomi. Tıp Araştırmaları Dergisi 2016;14:67-73.
30. Totoz T, Türk HŞ, Sayın P, Ünsal O, Çınar S, Oba S. Yoğun bakım ünitemizdeki (YBÜ) perkütan trakeotomi pratiğimiz. Ş.E.E.A.H. Tıp Bülteni 2013;47:11-15.
31. Öncül S, Yılmaz M, Gaygusuz EA, Oysu DA, Esen O, Şimşek T, et al. Our experience in percutaneous tracheostomy which performed by Griggs method: 38 cases. Medical Journal of Kocaeli 2014;2:1-4.
32. Ersoy A, Ali A, Ünlü N, Kara D, Turgut N. Griggs Yöntemi ile Gerçekleştirilen 53 Perkütan Trakeostomi. Okmeydanı Tıp Dergisi 2012;28:134-137.
33. Canver CC, ChandaJ. Intraoperativeand postoperative risk factorsfor respiratory failure after coronary bypass. Ann Thorac Surg 2003;75:853-857; discussion 857-858.