

Yavuz Orak,  
Şehnaz Pelin Tanır,  
Hatice Tuğba Güngör,  
Sondemet Çetin,  
Filiz Orak,  
Fatih Mehmet Yazar,  
Ömer Faruk Boran,  
Adem Doğaner

## Reasons for Hospitalization, Sepsis Development and Mortality Among Syrian Patients in an Intensive Care Unit

### Suriyeli Hastaların Yoğun Bakım Ünitesinde Yatış Nedenleri, Sepsis Gelişimi ve Mortalite

Received/Geliş Tarihi : 12.09.2019  
Accepted/Kabul Tarihi : 15.03.2020

©Copyright 2020 by Turkish Society of Intensive Care  
Turkish Journal of Intensive Care published by Galenos Publishing House.

Yavuz Orak, Ömer Faruk Boran  
Kahramanmaraş Sütçü İmam University Faculty  
Of Medicine, Department of Anesthesiology and  
Reanimation, Kahramanmaraş, Turkey

Şehnaz Pelin Tanır, Hatice Tuğba Güngör,  
Sondemet Çetin  
Kahramanmaraş Necip Fazıl City Hospital, Clinic of  
Anesthesiology and Reanimation, Kahramanmaraş,  
Turkey

Filiz Orak  
Kahramanmaraş Sütçü İmam University Faculty  
of Medicine, Department of Microbiology,  
Kahramanmaraş, Turkey

Fatih Mehmet Yazar  
4Kahramanmaraş Sütçü İmam University Faculty  
of Medicine, Department of General Surgery,  
Kahramanmaraş, Turkey

Adem Doğaner  
Kahramanmaraş Sütçü İmam University Faculty of  
Medicine, Department of Biostatistics and Medical  
Informatics, Kahramanmaraş, Turkey

Yavuz Orak (✉),  
Kahramanmaraş Sütçü İmam University Faculty  
Of Medicine, Department of Anesthesiology and  
Reanimation, Kahramanmaraş, Turkey

E-mail : dryavuzorak@hotmail.com

Phone : +90 532 471 70 56

ORCID ID : orcid.org/0000-0002-8356-6223

**ABSTRACT Objective:** Our aim is to investigate causes of hospitalization of Syrian patients in an intensive care unit (ICU), development of sepsis, relevant reasons of pathogens and mortality rates. **Materials and Methods:** We carried out this study between 2012 and 2016. Patient information was analyzed retrospectively from records and files in the information system.

**Results:** One hundred thirty-nine Syrian patients were hospitalized in an ICU during the study period. The most common ICU diagnoses were respiratory tract infection (29 patients: 20.9%) and trauma (26 patients: 18.7%). Of these patients, 35 were diagnosed with sepsis during their treatment in the ICU. *Acinetobacter baumannii* and *E. coli* were isolated in the culture of the patients with sepsis (17, 12, respectively). *Acinetobacter baumannii* was most common in tracheal cultures and *E. coli* in urine cultures. Additionally, seven patients were diagnosed with sepsis at their first application to the hospital. H1N1 in two, *S. pneumoniae* in three, *H. influenzae* in one, *S. aureus* and aspergilloma in another patient were detected. While 45 of 139 patients died, 28 of 35 patients diagnosed with sepsis died. APACHE II scores, duration of mechanical ventilation, number of days spent in ICU were all higher in the deceased patients than in the surviving patients ( $p<0.001$ ,  $p<0.001$ ,  $p<0.001$ ).

**Conclusion:** Respiratory diseases were the most common causes for Syrian patients' hospitalization in the ICU and developing sepsis. For these patients, sepsis was still an important factor for mortality.

**Keywords:** Syrian patients, sepsis, mortality, intensive care unit

**ÖZ Amaç:** Amacımız; Suriyeli hastaların yoğun bakım (YB) yatış nedenlerini, sepsis gelişimlerini, sepsise neden olan patojenleri ve hastaların ölüm oranlarını araştırmaktır.

**Gereç ve Yöntem:** Çalışmamız 2012-2016 tarihleri arasında YB ünitesinde yapıldı. Hastane bilgi sistemindeki kayıt ve dosyalar geriye dönük olarak analiz edildi.

**Bulgular:** Çalışma süresi içinde 139 Suriyeli hasta YB'ye yatırıldı. En sık YB yatış tanıları solunum yolu enfeksiyonu (29 hasta: %20,9) ve travma (26 hasta: %18,7) idi. Bu hastaların YB tedavileri sırasında 35 hasta sepsis tanısı aldı. Sepsis tanısı alan hastaların kültürlerinin 17'sinde *Acinetobacter baumannii* ve 12'sinde *E. Coli* üredi. *Acinetobacter baumannii* trakea kültürlerinde, *E. Coli* idrar kültürlerinde en yaygındı. Ayrıca, yedi hastaya hastaneye müracaat ettiği ilk anda sepsis tanısı kondu. Bu hastalardan 2'sinde H1N1, 3'ünde *S. pneumoniae*, 1'inde *H. influenzae*, 1'inde *S. aureus* ve aspergilloma saptandı. YB'da yatan 139 hastanın 45'i ölmüş iken sepsis tanısı alan 35 hastanın 28'i öldü. APACHE II skorları, mekanik ventilasyon süresi, YB yatış süresi ölenlerde sağ kalan hastalara göre daha yüksekti ( $p<0,001$ ,  $p<0,001$ ,  $p<0,001$ ).

**Sonuç:** Suriyeli hastalarda en sık YB yatış ve sepsis gelişme nedeni solunum yolu enfeksiyonlarıydı. Bu hastalarda sepsis mortalite için önemli bir faktördü.

**Anahtar Kelimeler:** Suriyeli hastalar, sepsis, mortalite, yoğun bakım ünitesi

## Introduction

More than 43.3 million people worldwide are fleeing war, and there are some political or ethnic pressures regarding taking refugees in another country (1,2). The war in Syria is tragically deadly and devastating enough to affect not only Syria's neighbors; Lebanon, Egypt, Iraq, and Turkey, but also European nations bordering Turkey (3,4). Since March, 2016, Turkey, with 2.7 million Syrians, has been one of the countries who have hosted the most refugees (3-5). There are approximately 300,000 refugees living in camps in the vicinity of Turkish cities near the Syrian border (Şanlıurfa, Gaziantep, Kilis, Hatay, Kahramanmaraş, Adıyaman, Adana, and Osmaniye). Additionally, there are about 2,440,000 refugees living outside the camps, who have spread from the south of Turkey to her central and western cities, like İstanbul, Ankara, and İzmir (4-6).

The disease profiles of refugees, who were at first exposed to traumatic injuries as a result of violent conflicts, then made dangerous journeys to other countries. They would take refugees in different from the disease profiles of the local people, both because of regional differences in some infectious diseases and insufficient living conditions in the nations to which the refugees had migrated (7,8).

In studies conducted in the early stages of the migration, the most important reasons for hospitalization in the intensive care unit (ICU)'s were traumatic pathologies and high-velocity gunshot wounds (9,10). Later, an increase was reported in reasons like respiratory system, cardiac, and tumoral pathologies (11,12). As far as we know, there are no studies to date on the pathogen profiles and sepsis rates of Syrian patients in the ICU's.

The aim of this study was to perform a general evaluation of the indications of hospitalization among Syrian patients in ICU, and to analyze these patients' rates of sepsis development, effective pathogens, and mortality rates and causes.

## Material and Methods

The study was carried out between June 1, 2012, and December 31, 2016, in two separate 14-bed Anesthesiology and Reanimation Intensive Care Units (ARICUs). Approval for the study (05.04.2017/04) was obtained from Kahramanmaraş Sütçü İmam University's Ethics Committee.

Of the 4,912 patients hospitalized in the units, 300 (6.1%) were Syrian. A total of 139 (46.3%) patients whose records

could be accessed were included in the study. Patient information was analyzed retrospectively from the records and patient files in the hospital's information management system.

There were a plethora number of factors for the evaluation stage. Namely, the following were considered for this purpose: patients' ages, gender, diagnoses at hospitalization, comorbid diseases, mechanical ventilation (MV) and number of days on MV, surgeries, diagnoses of sepsis, presence of sepsis originating from intensive care, areas from which cultures were obtained, microorganisms found, antibiotics used, number of days in the ICU, mortality rates, and Acute Physiologic Assessment and Chronic Health Evaluation (APACHE) II scores. For the patients who developed sepsis, the following data were recorded upon admission: the blood glucose level, the Sequential Organ Failure Assessment (SOFA) score and presence or absence of sepsis, the number of days before sepsis set in, the neutrophil-lymphocyte ratio, and the present microorganisms and the parts of the body where they were being produced.

### Definitions

In the study, patients were considered septic when they had SOFA scores of 2 or more and septic pathogens were found, as defined in the international guidelines of the Surviving Sepsis Campaign.

A community acquired infection was detected within the first 48-72 hours of hospitalization or acquired in daily life without significant immune deficiency (11).

### Statistical Analysis

The SPSS 24.0 program (IBM Corporation, Armonk, New York, United States) was used in analysis of the variables. The Shapiro-Wilk test was used to determine the normal distribution of the data. The Mann-Whitney U test was used with the Monte Carlo simulation technique in the comparison of two independent groups with regard to the quantitative data. For comparing categorical variables with one another, Pearson's chi-squared test, Fisher's exact test, and the Fisher-Freeman-Halton test were performed with the Exact and Monte Carlo Simulation techniques. Column ratios were compared to one another and expressed in accordance with the Benjamini-Hochberg adjusted p-values. An odds ratio with a 95% confidence interval was used to show the odds of an outcome occurring in the presence of an exposure as compared to the odds of the outcome occurring in the absence of the exposure.

Quantitative variables are shown in the tables as median (minimum/maximum), and categorical variables as n (%). Variables were examined at a confidence level of 95%, and a p-value was accepted as less than 0.05.

## Results

### General Findings

The mean age of the 139 patients who were included in the study was 46 (1-90), and 90 (647%) of them were male. Diagnoses at the initial hospitalization included 20 patients (14.4%) with gunshot wounds and 29 (20.9%) with respiratory tract infections. The patients were separated into

two groups; survivor and deceased. A total of 45 (32.4%) of the patients died in the ARICU.

APACHE II scores, the length of time connected to MV, and the number of days hospitalized in intensive care were all found to be greater in the deceased patients than in the living patients ( $p<0.001$ ,  $p<0.001$ , and  $p<0.001$ , respectively). Twenty-three (24.5%) of the survivors were trauma patients, and 16 (35.5%) of the deceased were respiratory disease patients. Surgery had been performed on 51 (54.2%) of the surviving patients and on 16 (17%) of the deceased ones ( $p=0.047$ ). The distributions of the patients according to their diagnoses and socio-demographic characteristics are given in Table 1.

**Table 1. Surviving and deceased patients' demographic and clinical characteristics**

		Surviving (n=94)	Deceased (n=45)	Total (n=139)	p	OR (95% CI)
<b>Age</b>	Median (min/max)	43 (1/87)	54 (3/90)	46 (1/90)	0.068	-
<b>Apache II</b>	Median (min/max)	11 (6/29)	29 (9/38)	13 (6/38)	<0.001	-
<b>DMV</b>	Median (min/max)	0 (0/43)	7 (0/270)	1 (0/270)	<0.001	-
<b>DIC</b>	Median (min/max)	3 (1/46)	9 (1/270)	4 (1/270)	<0.001	-
<b>Sex</b>					0.189	
Female	n (%)	36 (38.3)	12 (26.7)	48 (34.5)		-
Male	n (%)	58 (61.7)	33 (73.3)	91 (65.5)		-
<b>Province-referral from</b>						
Patient from other province	n (%)	7 (7.4)	5 (11.1)	12(8.6)	0.588	-
Same-province patient	n (%)	78 (83.0)	34 (75.6)	112 (80.6)		-
Hatay state hospital	n (%)	9 (9.6)	6 (13.3)	15(10.8)		-
<b>Diagnosis</b>						
Gunshot wound	n (%)	11 (11.7)	9 (20.0)	20 (14.4)	0.004	-
Respiratory disease	n (%)	13 (13.8)	16 (35.6)	29 (20.9)		3.4 (1.5-8.01)
Renal disease	n (%)	0 (0.0)	2 (4.4)	2 (1.4)		-
Trauma	n (%)	23 (24.5)	3 (6.7)	26 (18.7)		4.5 (1.3-16.02)
Cardiac disease	n (%)	5 (5.3)	5 (11.1)	10 (7.2)		-
Cancer	n (%)	12 (12.8)	2 (4.4)	14 (10.1)		-
Cerebrovascular disease	n (%)	6 (6.4)	3 (6.7)	9 (6.5)		-
Intoxication	n (%)	6 (6.4)	1 (2.2)	7 (5.0)		-
Gastrointestinal disease	n (%)	10 (10.6)	2 (4.4)	12 (8.6)		-
Other	n (%)	8 (8.5)	2 (4.4)	10 (7.2)		-
<b>MV usage</b>						
No	n (%)	55 (58.5)	2 (4.4)	57 (41.0)	<0.001	-
Yes	n (%)	39 (41.5)	43 (95.6)	82 (59.0)		30.3 (6.9-132.6)

		Surviving (n=94)	Deceased (n=45)	Total (n=139)	p	OR (95% CI)
<b>Surgery</b>					0.047	
No	n (%)	43 (45.7)	29 (64.4)	72 (51.8)		-
Yes	n (%)	51 (54.3)	16 (35.6)	67 (48.2)		2.1 (1.03-4.5)
<b>Anatomical region injured</b>					0.091	
Head	n (%)	6 (11.8)	6 (37.5)	12 (17.9)		-
Thorax and abdomen	n (%)	36 (70.6)	8 (50.0)	44 (65.7)		-
Extremities	n (%)	9 (17.6)	2 (12.5)	11 (16.4)		-

Mann-Whitney U test (Monte Carlo) / Pearson's chi-squared test (Exact/Monte Carlo) / Fisher-Freeman-Halton test (Monte Carlo) / Fisher exact test (Exact) / OddsRatio (95% Confidence Interval) /, min: minimum, max: maximum,  
DMV: Days on mechanical ventilator, DIC: Days in intensive care, MV: Mechanical ventilation, APACHE: Acute physiologic assessment and chronic health evaluation

In 43 patients (30.9%), the most common comorbid diseases after their initial diagnosis was cardiac disease (Table 2).

### Antimicrobial Therapy, Infection Type and Sepsis Findings

Of the living patients, 62 (66%) received single antibiotics while 27 (60%) of the deceased patients received multiple antibiotics ( $p < 0.001$ ). The blood cultures were positive in 8 (8.5%) of the surviving patients and the trachea cultures were positive in 5 (5.3%). In addition, 15 (33.3%) of the deceased patients' blood cultures were positive, as were 18 (40%) of their trachea cultures ( $p < 0.001$  and  $p < 0.001$ , respectively) (Table 3).

While 89 (48.9%) of all the patients did not produce microorganisms, *Acinetobacter baumannii* was found in 17 (9.3%) and *E. Coli* in 15 (8.2%). Of the patients who developed sepsis, 17 (23.6%) produced *Acinetobacter baumannii* and 12 (16.7%) *E. Coli*. *Acinetobacter baumannii* was detected most frequently in trachea cultures and *E. Coli* in urine cultures (Table 4).

The 35 patients with sepsis were divided into two groups: those who survived ( $n=7$ ) and those who did not ( $n=28$ ). The patients' clinical information is given in Table 5.

Ten of the septic patients (28.6%) had diseases of the respiratory system, seven (20%) had gunshot injuries, and five (14.3%) were diagnosed with trauma (Table 6).

$\beta$ -lactam /  $\beta$ -lactamase inhibitor-type antibiotics were used on 65 (28.6%) of the patients, while 2<sup>nd</sup>-, 3<sup>rd</sup>-, and 4<sup>th</sup>- generation cephalosporins were used on 43 (18.9%). Of the patients with sepsis, carbapenems were used on 17 (19.5%), colistin on 16 (18.4%), and tigecycline on 11 (12.6%) patients (Table 7).

	n=163 (%) *X
None	67 (41.1)
Cardiac diseases	43 (26.4)
Diabetes Mellitus	17 (10.4)
Renal diseases	15 (9.2)
Respiratory diseases	10 (6.1)
Cerebrovascular diseases	6 (3.7)
Other	5 (3.1)
Total	163

\*X: One patient has more than one disease

### A Community-acquired Infection

Seven patients were diagnosed with sepsis at the time of hospitalization. Oseltamivir treatment was begun on two patients with initial diagnoses, later confirmed, of H1N1. *S. pneumoniae* was found in three patients, *H. influenzae* in one, and *S. aureus* and aspergilloma in one.

### Discussion

In our study, we investigated the reasons for hospitalization, the sepsis development rates, the pathogens causing sepsis, and the mortality rates and causes for Syrian patients in ICU.

In the early period of the studies relating to the Syrian civil war, patients underwent surgery due to trauma (9,10). In a study evaluating arrivals of both refugees and locals at emergency service units, it was reported that refugees' medical emergencies included high rates of many types

<b>Table 3. Evaluation of cultures taken from surviving and deceased patients</b>						
		Surviving (n=94)	Deceased (n=45)	Total (n=139)	p	OR (95% CI)
<b>Culture taken from</b>						
None taken	n (%)	77 (81.9)	14 (31.1)	91 (65.5)	<0.001	-
Tracheal aspirate	n (%)	2 (2.1)	8 (17.8)	10 (7.2)		22 (4.2-114.6)
Blood	n (%)	5 (5.3)	7 (15.6)	12 (8.6)		7.7 (2.1-27.7)
Urine	n (%)	0 (0.0)	2 (4.4)	2 (1.4)		-
Injury site	n (%)	6 (6.4)	3 (6.7)	9 (6.5)		-
Two or more regions	n (%)	4 (4.3)	11 (24.4)	15 (10.8)		15.1 (4.2-54.3)
<b>Microorganism found</b>						
None	n (%)	74 (78.7)	15 (33.3)	89 (64.0)	<0.001	-
Single microorganism	n (%)	10 (10.6)	15 (33.3)	25 (18.0)		7.7 (2.9-20.4)
Multiple microorganisms	n (%)	10 (10.6)	15 (33.3)	25 (18.0)		7.7 (2.9-20.4)
<b>Antibiotic administered</b>						
None	n (%)	12 (12.8)	1 (2.2)	13 (9.4)	< 0.001	-
Single antibiotic	n (%)	62 (66.0)	17 (37.8)	79 (56.8)		3.2 (0.4-27.1)
Multiple antibiotics	n (%)	20 (21.3)	27 (60.0)	47 (33.8)		16.2 (1.9-135.01)
<b>Urine culture</b>						
Negative	n (%)	89 (94.7)	38 (84.4)	127 (91.4)	0.056	-
Positive	n (%)	5 (5.3)	7 (15.6)	12 (8.6)		-
<b>Blood culture</b>						
Negative	n (%)	86 (91.5)	30 (66.7)	116 (83.5)	<0.001	-
Positive	n (%)	8 (8.5)	15 (33.3)	23 (16.5)		5.4 (2.1-13.9)
<b>Tracheal culture</b>						
Negative	n (%)	89 (94.7)	27 (60.0)	116 (83.5)	<0.001	-
Positive	n (%)	5 (5.3)	18 (40.0)	23 (16.5)		11.9 (4.02-34.9)
<b>Injury site culture</b>						
Negative	n (%)	85 (90.4)	42 (93.3)	127 (91.4)	0.751	-
Positive	n (%)	9 (9.6)	3 (6.7)	12 (8.6)		-
<b>Sepsis</b>						
No	n (%)	86 (91.5)	19 (40.4)	104 (75.5)	<0.001	-
Yes	n (%)	7 (7.5)	28 (59.6)	35 (24.5)		14.7 (5.8-37.5)
Mann-Whitney U test (Monte Carlo) / Pearson's chi-squared test (Exact/Monte Carlo) / Fisher-Freeman-Halton test (Monte Carlo) / Fisher exact test (Exact) / OddsRatio (95% Confidence Interval) / , min: minimum, max: maximum						

of trauma, especially of the head, neck, and extremities, as compared to those rates for local residents (9). Another study showed that the most common surgical site regions were head and neck (52.7%), followed by the thorax and abdomen (27.8%), and multiple-system injuries (13.8%) (10). In another study on trauma, 24.2% of the patients had head and neck trauma, and 15.3% had chest, abdomen and back

trauma (13). Duramaz et al. (14) reported that injuries were more common in lower extremities, upper extremities and axial skeleton. Blunt trauma was significantly higher in upper extremity injuries compared to other types of injuries. In a study conducted in central Europe, the the most common reasons for admission among Syrian patients were surgery (43.3%), medical (36.5%) and psychiatric (15.6%). In addition,

**Table 4. Distribution of microorganisms detected, by culture type**

	All patients n=139	Sepsis patients n=35	Blood culture	Tracheal culture	Urine culture	Injury site culture
Microorganism	n(%)	n (%)	n (%)	n (%)	n (%)	n (%)
None detected	89 (48.9)	-	-	-	-	-
Acinetobacterbaumannii	17 (9.3)	17 (23.6)	5 (17.8)	9 (34.7)	3 (21.4)	3 (42.8)
Streptococcus pneumoniae	3 (1.6)	3 (4.2)	2 (7.1)	0 (0.0)	1 (7.1)	0 (0.0)
Candida fungus	9 (4.9)	8 (11.1)	4 (14.3)	2 (7.7)	2 (14.4)	0 (0.0)
MSSA	6 (3.3)	4 (5.6)	2 (7.1)	1 (3.8)	1 (7.1)	0 (0.0)
CNS	12 (6.5)	5 (6.9)	4 (14.3)	0 (0.0)	1 (7.1)	0 (0.0)
<i>E. Coli</i>	15 (8.2)	12 (16.7)	1 (3.6)	3 (11.5)	6 (42.9)	2 (28.6)
Klebsiella pneumoniae	8 (4.4)	7 (9.7)	0 (0.0)	6 (23.2)	0 (0.0)	1 (14.3)
VRE	1 (0.6)	1 (1.4)	1 (3.6)	0 (0.0)	0 (0.0)	0 (0.0)
Stenotrophomonasmaltophilia	4 (2.2)	4 (5.5)	3 (10.7)	1 (3.8)	0 (0.0)	0 (0.0)
Pseudomonasaeruginosa	7 (3.9)	4 (5.5)	1 (3.6)	2 (7.7)	0 (0.0)	1 (14.3)
Enterococcus	4 (2.2)	3 (4.1)	2 (7.1)	1 (3.8)	0 (0.0)	0 (0.0)
<i>H. influenzae</i>	1 (0.6)	1 (1.4)	0 (0.0)	1 (3.8)	0 (0.0)	0 (0.0)
<i>Aspergillusfumigatus</i>	1 (0.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Enterobacteriaceae	2 (1.1)	1 (1.4)	1 (3.6)	0 (0.0)	0 (0.0)	0 (0.0)
MRSA	1 (0.6)	1 (1.4)	1 (3.6)	0 (0.0)	0 (0.0)	0 (0.0)
Viruses (H1N1)	2 (1.1)	1 (1.4)	1 (3.6)	0 (0.0)	0 (0.0)	0 (0.0)
<b>Toplam</b>	<b>182/100</b>	<b>72/100</b>	<b>28/100</b>	<b>26/100</b>	<b>14/100</b>	<b>7/100</b>
MSSA:, CNS, VRE, MRSA						

the most common acute infectious diseases (43.9%) were respiratory, gastrointestinal and urinary tract infections (15).

In our study, it was seen that the most important causes of hospitalization were pathologies of the respiratory system (20.9%) trauma (18.7 %) and gunshot wound (14.4 %). Besides, 67 (48.2%) of our patients underwent surgical intervention. 44 (65.7%- multiple injuries) of patients had thorax and abdominal injuries. It was further noted that tumoral causes, which were not mentioned in the literature, accounted for as high a rate as 10.1%. In the present study, respiratory failure was found to be the most important cause (35.6%) of mortality. In the diagnostic-based evaluation of the patients, it was observed that the most septic complications were in patients with respiratory failure, and it was thought that this had an effect on the development of mortality (28%). In the correlation analysis we performed, the most important factor affecting mortality was the development of sepsis. In all Syrian patients, the overall mortality rate was 32.37%, while the same rate in Syrian patients with sepsis was 80%.

One of the aims of our study was the evaluation of

the mortality and of the pathogens that cause sepsis development in Syrian patients. In a multicenter international study, the hospital mortality rate of sepsis patients was found to be 47.2 % in Africa and 13.1% in North America (16). Another study showed that mortality rate of severe sepsis was 36.7 % in South/Central America and 44% Eastern Europe (17). In Uganda, hospital mortality associated with sepsis was 43% (18), in Thailand 50% (19). In a study conducted in an ICU in Turkey, the sepsis mortality rate was reported as 87.3%. The most commonly isolated agent in the development of sepsis, according to that study, was Gr (-) bacteria, at 65.9% (20). In our study, it was seen that the sepsis mortality rates of the Syrian patients were similar to those of the local population (80%).

The most common pathogen among sepsis patients was *Acinetobacter baumannii*; the second most common was *E. Coli*. While death from sepsis in high-income countries has been 30%-40% in the last decade and is steadily falling (21,22), the mortality rate in low-income regions of the world has risen as high as 80% (23-26). In a multicenter study conducted in Turkey, the mortality was

		Surviving	Deceased	Total	p
		(n=7)	(n=28)	(n=35)	
<b>Age</b>	<b>Mean ± SD</b>	41.29±19.11	49.93±22.24	48.20±21.67	0.290
<b>Day sepsis developed (+)</b>	<b>Median (min/max)</b>	7 (2/18)	5 (1/75)	5 (1/75)	0.774
<b>N/L Ratio</b>	<b>Median (min/max)</b>	9.63 (5.75/150)	11.50 (0.67/110.34)	11.34 (0.67/150)	1
<b>N/L Ratio %</b>	<b>Median (min/max)</b>	9.65 (6.58/15.68)	11.19 (2.10/47.25)	11.01 (2.10/47.25)	0.749
<b>Blood glucose value at admission</b>	<b>Median (min/max)</b>	145 (83/229)	194 (73/475)	175 (73/475)	0.177
<b>SOFA score</b>	<b>Median (min/max)</b>	5 (2/5)	5 (2/10)	5 (2/10)	0.214
<b>DIC</b>	<b>Median (min/max)</b>	31 (11/46)	17 (2/270)	17 (2/270)	0.121
<b>Sex</b>					1
Female	<b>n (%)</b>	2 (28.6)	8 (28.6)	10 (28.6)	
Male	<b>n (%)</b>	5 (71.4)	20 (71.4)	25 (71.4)	
<b>ARF</b>					0.415
No	<b>n (%)</b>	5 (71.4)	14 (50.0)	19 (54.3)	
Yes	<b>n (%)</b>	2 (28.6)	14 (50.0)	16 (45.7)	
<b>Blood culture</b>					1
Negative	<b>n (%)</b>	3 (42.9)	11 (39.3)	14 (40.0)	
Positive	<b>n (%)</b>	4 (57.1)	17 (60.7)	21 (60.0)	
<b>Tracheal culture</b>					1
Negative	<b>n (%)</b>	3 (42.9)	11 (39.3)	14 (40.0)	
Positive	<b>n (%)</b>	4 (57.1)	17 (60.7)	21 (60.0)	
<b>Urine culture</b>					1
Negative	<b>n (%)</b>	5 (71.4)	20 (71.4)	25 (71.4)	
Positive	<b>n (%)</b>	2 (28.6)	8 (28.6)	10 (28.6)	
<b>Injury site culture</b>					0.256
Negative	<b>n (%)</b>	5 (71.4)	25 (89.3)	30 (85.7)	
Positive	<b>n (%)</b>	2 (28.6)	3 (10.7)	5 (14.3)	

Mann-Whitney U test (Monte Carlo) / independentsamples t test/ Pearson's chi-squared test (Exact/Monte Carlo) / Fisher-Freeman-Halton test (Monte Carlo) / Fisher exact test (Exact) / OddsRatio (95% Confidence Interval) /, min: minimum, max: maximum, SD: Standard deviation, ARF: Acute renal failure, N/L ratio: Neutrophil/Lymphocyte ratio, DIC: Days in intensive care, SOFA: Sequential organ failure assessment

found to be very high among patients with sepsis and septic shock in the ICU (55.7% and 70.4, respectively). The most isolated microorganism was *Acinetobacter* spp (33.7%) and approximately 74.8% of *Acinetobacter* were resistant to carbapenems (27). In our study, the rate of *Acinetobacter* spp in all Syrian patients was 9.3% and 23.6% in patients with sepsis. Carbapenem use rates were 11.9% in all Syrian patients and 19.5% in sepsis patients.

In the study of Turktan et al, 37 patients in an ICU were hospitalized because of infection. The most common

community infection was pneumonia (49%) and urinary infection (16.3%). In eight of them, the infection developed as a result of community-acquired microorganisms, the most important of which was *M. tuberculosis* (11). This situation was thought to have been caused by the difficulty of the refugees' living conditions, and high population density of their living quarters. *E. Coli* was the second most common pathogen reported in that study (11). In our study, although *M. tuberculosis* was not observed as a causative pathogen, it was noted that at the time of admission, seven patients had

Diagnosis	n=35(%)
Gunshot wound	7 (20.0)
Respiratory diseases	10 (28.6)
Renal diseases	2 (5.7)
Trauma	5 (14.3)
Cardiac diseases	5 (14.3)
Cancer	3 (8.6)
Intoxication	1 (2.9)
GI system	1 (2.9)
Other	1 (2.9)
GI:	

Antibiotic	All patients n (%) (X*)	Sepsis patients n (%) (X*)
None used	13 (5.7)	-
$\beta$ -lactam / $\beta$ -lactamase inhibitor	65 (28.6)	12 (13.8)
2 <sup>nd</sup> -, 3 <sup>rd</sup> and 4 <sup>th</sup> -generation cephalosporins	43 (18.9)	2 (2.3)
Carbapenems	27 (11.9)	17 (19.5)
Colistin	15 (6.6)	16 (18.4)
Tigecycline	12 (5.3)	11 (12.6)
Quinolones	11 (4.8)	3 (3.4)
Anti-anaerobic agents	8 (3.5)	3 (3.4)
Linezolid	8 (3.5)	4 (4.6)
Macrolides	7 (3.1)	3 (3.4)
Antifungals	7 (3.1)	7 (8.0)
Sulfonamides	4 (1.8)	4 (4.6)
Oseltamivir	3 (1.3)	2 (2.3)
Aminoglycosides	2 (0.9)	1 (1.1)
Glycopeptides	2 (0.9)	2 (2.3)
X*: One patient used more than one		

developed sepsis as a result of other community-acquired factors. Two of those patients were initially diagnosed with H1N1, and Oseltamivir treatment was begun; the diagnosis was later confirmed. *S. pneumoniae* was detected in three of the remaining patients, *H. influenzae* in one, and *S. aureus* and aspergilloma in the last one.

It has been reported that ICU admission rates are high in migrant populations because such factors as a lack

of prevention and protection in healthcare services, the withdrawal of vaccination programs, low environmental hygiene standards, outdoor living conditions, overcrowding, and exposure to low temperatures have raised the sensitivity to infection (28-33).

Regarding the limitations of our study, it was not possible to evaluate the ICU hospitalization among the local population in the same period, and the number of cases was low. However, its most important advantage is the fact that it is the first study evaluating sepsis among the Syrian patients in an ICU in our country; as such, it can shed light on wider studies to be conducted.

## Conclusion

Syrian patients were admitted to the ICU mostly due to respiratory system disease. Comorbid disease was the most common heart disease. *Acinetobacter baumannii* and *E. Coli* were isolated most frequently in Syrian patients and 2<sup>nd</sup>-, 3<sup>rd</sup>- and 4<sup>th</sup>- generation cephalosporins were used most frequently in septic patients, and carbapenems also were used in these patients. APACHE II scores, the duration of MV, and the number of days hospitalized in ICU were all found to be higher for deceased patients than for the survivors.

## Ethics

**Ethics Committee Approval:** Approval for the study (05.04.2017/04) was obtained from Kahramanmaraş Sütçü İmam University's Ethics Committee.

**Informed Consent:** Patient information was analyzed retrospectively from the records and patient files in the hospital's information management system.

**Peer-review:** Externally peer-reviewed.

## Authorship Contributions

Concept: Y.O., Ş.P.T, H.T.G., S.Ç., F.O., F.M.Y., Ö.F.B., A.D., Design: Y.O., Ş.P.T, H.T.G., S.Ç., F.O., F.M.Y., Ö.F.B., A.D., Data Collection or Processing: Y.O., Ş.P.T, H.T.G., S.Ç., F.O., F.M.Y., Ö.F.B., A.D., Analysis or Interpretation: Y.O., Ş.P.T, H.T.G., S.Ç., F.O., F.M.Y., Ö.F.B., A.D., Literature Search: Y.O., Ş.P.T, S.Ç., F.O., F.M.Y., Ö.F.B., A.D., Writing: Y.O., F.O., F.M.Y., A.D.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study received no financial support.

## References

- Maier T, Schmidt M, Mueller J. Mental health and health care utilization in adult asylum seekers. *Swiss Medical Weekly* 2010;140:w13110.
- Pfortmueller CA, Graf F, Tabarra M, Lindner G, Zimmermann H, Exadaktylos AK. Acute health problems in African refugees: ten years' experience in a Swiss emergency department. *Wiener Klinische Wochenschrift* 2012;124:647-52.
- United Nations High Commissioner for Refugees. UNHCR Global Appeal 2015. Available from: <http://www.unhcr.org/5461e60716.htm>. Accessed June 16, 2016.
- 3RP Regional Refugee & Resilience Plan 2016-2017 in response to the Syria crisis. 2016. Available from: <http://www.3rp.syria.crisis.org/wp/wp-content/uploads/2016/02/Turkey-2016-Regional-Refugee-Resilience-Plan.pdf>. Accessed June 16, 2016.
- AFAD. Republic of Turkey Prime Ministry Disaster and Emergency Management Presidency. Syrian Refugees in Turkey, 2013. Field Survey Results. Available from: [https://www.afad.gov.tr/Dokuman/TR/61-2013123015505-syrian-refugees-in-turkey2013\\_print\\_12.11.2013\\_eng.pdf](https://www.afad.gov.tr/Dokuman/TR/61-2013123015505-syrian-refugees-in-turkey2013_print_12.11.2013_eng.pdf). Accessed June 16, 2016.
- AFAD. Republic of Turkey Prime Ministry Disaster and Emergency Management Presidency. Current Status in AFAD Temporary Protection Centres. 2016. Available from: <http://www.afad.gov.tr/en/lcerik Yazdir.aspx?ID=16&lcerikID=848>. Accessed June 16, 2016.
- Bischoff A, Denhaerynck K, Schneider M, Battegay E. The cost of war and the cost of health care—an epidemiological study of asylum seekers. *Swiss Medical Weekly* 2011;141:w13252.
- Pfortmueller CA, Stotz M, Lindner G, Muller T, Rodondi N, Exadaktylos AK. Multi morbidity in adult asylum seekers: a first overview. *PLoS One* 2013;8:e82671.
- Duzkoylu Y, Basceken SI, Kesilmez EC. Physical Trauma among Refugees: Comparison between Refugees and Local Population Who Were Admitted to Emergency Department-Experience of a State Hospital in Syrian Border District. *J Environ Public Health* 2017;2017:8626275.
- Kocamer ŞB, Dokur M, Uysal E, Çaliker N, Gökçe ON, Deniz İK, et al. Characteristics of the injuries of Syrian refugees sustained during the civil war. *Ulus Travma Acil Cerrahi Derg* 2017;23:199-206.
- Turktan M, Ak O, Erdem H, Ozcengiz D, Hargreaves S, Kaya S, et al. Community acquired Infections among refugees leading to Intensive Care Unit admissions in Turkey. *Int J Infect Dis* 2017;58:111-4.
- van Berlaer G, Bohle Carbonell F, Manantsoa S, de Béthune X, Buyl R, Debacker M, et al. A refugee camp in the centre of Europe: clinical characteristics of asylum seekers arriving in Brussels. *BMJ Open* 2016;24;6:e013963.
- Çelikel A, Karaarslan B, Demirkıran DS, Zeren C, Arslan MM. A series of civilian fatalities during the war in Syria. *Ulus Travma Acil Cerrahi Derg* 2014;20:338-42.
- Duramaz A, Bilgili MG, Bayram B, Ziroglu N, Bayrak A, Avkan MC. Orthopedic trauma surgery and hospital cost analysis in refugees; the effect of the Syrian civil War. *International Orthopaedics* 2017;41:877-84.
- Pfortmueller CA, Schwetlick M, Mueller T, Lehmann B, Exadaktylos AK. Adult Asylum Seekers from the Middle East Including Syria in Central Europe: What Are Their Health Care Problems? *PLoS One* 2016;11:e0148196.
- Vincent J-L, Marshall JC, Namendys-Silva SA, François B, Martin-Loeches I, Lipman J, et al. Assessment of the world wide burden of critical illness: the intensive care over nations (ICON) audit. *Lancet Respir Med* 2014;2:380-6.
- Rhodes A, Phillips G, Beale R, Cecconi M, Chiche JD, De Backer D, et al. The Surviving Sepsis Campaign bundles and outcome: results from the International Multicentre Prevalence Study on Sepsis (the IMPReSS study). *Intensive Care Med* 2015;41:1620-8.
- Jacob ST, Moore CC, Banura P, Pinkerton R, Meya D, Opendi P, et al. Severe sepsis in two Ugandan hospitals: a prospective observational study of management and outcomes in a predominantly HIV-1 infected population. *PLoS One* 2009;4:e7782.
- Khwannimit B, Bhurayanontachai R. The epidemiology of, and risk factors for, mortality from severe sepsis and septic shock in a tertiary-care university hospital setting. *Epidemiol Infect* 2009;137:1333-41.
- Tanriover MD, Guven GS, Sen D, Unal S, Uzun O. Epidemiology and outcome of sepsis in a tertiary-care hospital in a developing country. *Epidemiol Infect* 2006;134:315-22.
- Martin GS, Mannino DM, Eaton S, Moss M. The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 2003;348:1546-54.
- Vincent JL, Sakr Y, Sprung CL, Ranieri VM, Reinhart K, Gerlach H, et al. Sepsis Occurrence in Acutely Ill Patients I Sepsis in European intensive care units: results of the SOAP study. *Crit Care Med* 2006;34:344-53.
- Frikha N, Mebazaa M, Mnif L, El Euch N, Abassi M, Ben Ammar MS [Septic shock in a Tunisian intensive care unit: mortality and predictive factors. 100 cases]. *Tunis Med* 2005;83:320-5.
- Jacob ST, Moore CC, Banura P, Pinkerton R, Meya D, Opendi P, et al. Promoting Resource-Limited Interventions for Sepsis Management in Uganda Study G Severe sepsis in two Ugandan hospitals: a prospective observational study of management and outcomes in a predominantly HIV-1 infected population. *PLoS One* 2009;4:e7782.
- Siddiqui S. Not "surviving sepsis" in the developing countries. *J Indian Med Assoc* 2007;105:221
- Dünser MW, Bataar O, Tsenddorj G, Lundeg G, Torgersen C, Romand JA, et al. Differences in critical care practice between an industrialized and a developing country. *Wien Klin Wochenschr* 2008;120:600-7.
- Baykara N, Akalin H, Arslantaş MK, Hancı V, Çağlayan Ç, Kahveci F, et al. Epidemiology of sepsis in intensive care units in Turkey: a multicenter, point-prevalence study. *Critical Care* 2018;22:93.
- Healing TD, Pelly MD. Refugees and disasters. In: Parry E, Godfrey R, Mabet D, Gill G editors. *Principles of medicine in Africa*. Cambridge University Press: Singapore; 2004. p. 77-86.
- McGready R, Ashley EA, Wuthiekanun V, Tan SO, Pimanpanarak M, Viladpai-Nguen SJ, et al. Arthropod borne disease: the leading cause of fever in pregnancy on the Thai-Burmese border. *PLoS Negl Trop Dis* 2010;4:e888.
- Esteban-Vasallo MD, Dominguez-Berjon MF, Aerny-Perreten N, Astray-Mochales J, Martin-Martinez F, Genova-Maleras R. Pandemic influenza A (H1N1) 2009 in Madrid, Spain: incidence

- and characteristics in immigrant and native population. *Eur J Public Health* 2012;22:792-6.
31. Pourkarim MR, Zandi K, Davani NA, Pourkarim HR, Amini-Bavil- Olyae S. An aberrant high prevalence of hepatitis B infection among Afghans residing in one of the Bushehr refugee camps (Dalakicamp) in the south west of Iran. *Int J Infect Dis* 2008;12:101-2.
  32. Fan CK, Liao CW, Wu MS, Hu NY, Su KE. Prevalence of *Pediculus capitis* infestation among school children of Chinese refugees residing in mountainous areas of northern Thailand. *Kaohsiung J Med Sci* 2004;20:183-7.
  33. Mateen FJ, Carone M, Al-Saedy H, Nyce S, Ghosn J, Mutuerandu T, et al. Medical conditions among Iraq refugees in Jordan: data from the United Nations Refugee Assistance Information System. *Bull World Health Organ* 2012;90:444-51.