

The Effect of Psychodrama on the Empathy and Social Anxiety Level in Adolescents

Psikodramanın Ergenlerde Empati ve Sosyal Kaygı Düzeyine Etkisi

Çağlar Şimşek¹, Ebru Yılmaz Yalçınkaya², Elif Ardiç³, Ejder Akgün Yıldırım⁴

¹Taksim Training and Research Hospital, Patient Rights Unit, Istanbul, Turkey

²University of Health Sciences Turkey, Gaziosmanpaşa Taksim Training and Research Hospital, Clinic of Physical Medicine and Rehabilitation, Istanbul, Turkey

³Marmara University Faculty of Health Sciences, Department of Psychiatric Nursing, Istanbul, Turkey

⁴University of Health Sciences Turkey, Bakırköy Prof. Dr. Mazhar Osman Mental Health and Diseases Training and Research Hospital, Psychotherapy Training and Treatment Center, Istanbul, Turkey

ABSTRACT

Objectives: This study was conducted semi-experimentally to examine the effect of psychodrama on the level of social anxiety and empathy.

Materials and Methods: The sample of the study consisted of nine volunteer students aged 13-17. The study took place between 05 March 2002 and April 2018. Psychodrama sessions consisted of 10 sessions in total, 2 days a week. Empathic Tendency Adolescent scale and Social Anxiety scale were used as data collection tools in the study.

Results: When the total scores obtained from the Social Anxiety scale were examined, a statistically significant difference was found before and after psychodrama ($p=0.029$). When the total scores obtained from the Empathic Tendency scale were analysed, no statistically significant difference was found. However, after psychodrama, there was an increase in the average score of the Empathic Tendency scale.

Conclusion: Psychodrama is an effective method for reducing social anxiety and developing empathic skills in adolescents.

Keywords: Adolescents, social anxiety, empathy

ÖZ

Amaç: Bu çalışma, psikodramanın sosyal kaygı ve empati düzeyine etkisini incelemek amacıyla yarı deneysel olarak yapılmıştır.

Gereç ve Yöntem: Çalışmanın örneklemini, 13-17 yaş aralığında yer alan dokuz gönüllü öğrenci oluşturmuştur. Çalışma 05 Mart - 02 Nisan 2018 tarihleri arasında gerçekleşmiştir. Psikodrama oturumları haftada 2 gün olmak üzere toplamda 10 oturumdan oluşmuştur. Çalışmada veri toplama aracı olarak Empatik Eğilim Ergen ölçeği ve Sosyal Kaygı ölçeği kullanılmıştır.

Bulgular: Sosyal Kaygı ölçeğinden alınan toplam puanlar incelendiğinde, psikodrama öncesi ve sonrası istatistiksel olarak anlamlı fark bulunmuştur ($p=0,029$). Empatik Eğilim ölçeğinden alınan toplam puanlar incelendiğinde ise, istatistiksel olarak anlamlı bir fark yoktur. Ancak psikodrama sonrasında Empatik Eğilim ölçeği puan ortalamasında artış olmuştur.

Sonuç: Psikodrama ergenlerde sosyal kaygının azaltılması ve empatik becerilerin gelişmesinde etkili bir yöntemdir.

Anahtar Kelimeler: Ergenler, sosyal kaygı, empati

Introduction

Social anxiety is the avoidance of social relationships that involve social interactions, which occur in reactionary environments such as talking to others in a meeting or a party, and also felt in special situations such as an examination or assessment and speaking in public.¹ In this case, where keeping away from or avoidance of social environments, the individual shows the avoidance behavior because he/she thinks that he/she will be rejected by others.² Social evaluation anxiety is also a component of social anxiety;

self-assertion theorists argue that the expectation of interpersonal evaluation reveals social anxiety.^{3,4} Socially anxious individuals have doubts about their interpersonal competence and ability to create positive impressions.⁵ Researches have shown that although social anxiety is a common disorder, the age of onset becomes clear with adolescence. It is reported to be one of the most common forms of social distress in adolescents.^{6,7} If social anxiety is not treated, it may lead to discontinuation of education, social withdrawal, depression and alcohol-substance use.⁸ Psychodrama is one of the treatment methods for social anxiety.

Address for Correspondence/Yazışma Adresi: Çağlar Şimşek MD, Taksim Training and Research Hospital, Patient Rights Unit, Istanbul, Turkey

Phone: +90 505 746 32 59 E-mail: caglarsimsek1983@hotmail.com ORCID: orcid.org/0000-0002-7077-7486

Received/Geliş Tarihi: 14.10.2019 Accepted/Kabul Tarihi: 20.05.2020

©Copyright 2020 by the Turkish Association for child and Adolescent Psychiatry / Turkish Journal of Child and Adolescent Mental Health. published by Galenos Publishing House.



Psychodrama is a therapeutic method that aims to create insight and awareness by providing alternative solutions to the problems of individuals, groups and society depending on their emotions, thoughts and some disturbing events in their lives.^{9,10} It is a method of psychotherapeutic and sociotherapeutic action which is mostly applied in groups and is a method of psychotherapy in which individuals rehearse their lives.¹¹ As group therapy may be beneficial in the treatment of psychiatric diseases seen in adolescents, so there is a tendency to create more group activities in them.¹² In addition to the benefit of being in the peer group due to the age group characteristics, psychodrama applications also benefit in the realization of emotions, acquiring new social skills and developing direct self-confrontation and empathy skills in adolescents.^{13,14}

Empathy, which forms the basis of communication, has gained much importance today and has an essential role in interpersonal relations.¹⁵ It is generally regarded as a multidimensional structure that includes cognitive as well as affective processes, including understanding, perceiving and sharing the emotional states of others.¹⁶ It is the ability of the person to see his/her own point of view by putting himself/herself in the place of the other person in terms of emotion, thought and behavior, while respecting his/her point of view and looking at the events from his/her own point of view and acting in common.¹⁷ Dökmen¹⁸ also states that when an individual understands the feelings and ideas of others with empathy and conveys them to the other party, it is also regarded as a component of empathy. If adolescents have an empathic tendency, they will look at the events from different angles and put himself/herself in another's position and resolves the conflict in a more positive way. Resolution of conflicts with empathy in adolescents can have beneficial consequences. Thus, it is important that adolescents have empathic tendency, and they should use it in many areas such as resolution of conflicts.

Psychodrama studies conducted among individuals with social anxiety are limited. Uneri et al.¹⁹ indicated that no psychodrama studies on adolescents had been conducted up to that year. In addition, the situations experienced during adolescence leave important traces in the future lives of individuals. This study aims to determine the effect of psychodrama on the level of empathy and social anxiety in adolescents. The results of the study will contribute to the literature in this field.

Materials and Methods

In this study, a semi-experimental and special-purpose sampling method was used. The study took place between 19 February and 19 March. The sample of the study consisted of 8-13 year-old children hospitalized in the Physical Therapy Clinic of a Training and Research Hospital in the European side of Istanbul who had applied for reading and playing games in this clinic. The study group consisted of 9 volunteer students (4 boys and 5 girls) aged 13-17 years. Preliminary interviews were conducted with the students who applied to the hospital within the scope

of the social responsibility project of a secondary education institution. The students who stated their anxiety of how to interact with paediatric patients as a result of the interview were invited to the psychodrama study. The psychodrama study was carried out by the expert psychiatric nurse who completed the psychodrama training at the Abdülkadir Özbek Psychodrama Institute. During the study, 10 sessions of psychodrama was conducted with the supervision of Abdülkadir Özbek Psychodrama Institute trainer, 2 days a week for 2 hours in the hospital library between March and April 2018. In the study, Ka-Si Empathic Tendency Adolescent scale and Social Anxiety scale for adolescents (SAS-A) were used for data collection. Statistical evaluation was made by applying pre- and post-test.

Inclusion Criteria

- Applying to the hospital within the scope of the social responsibility project
- Being a secondary school student
- Not having a psychiatric diagnosis
- Being a volunteer
- The parents' consent

Exclusion Criteria

- Being a university student
- Being a primary school student

Psychodrama Sessions

A psychodrama session consists of three main phases: warm-up, play stage and group meeting.

Warm-up stage: A psychodrama session begins with the warm-up phase, which aims to prepare the group members and the manager for the role, topic and current situation. Warm-up allows members to increase their spontaneity and break the resistance of the group. At this stage, it is decided how the group will work and who will be the protagonist.²⁰

Play stage: This stage is the action stage that includes the implementation of the decisions made during the warm-up phase. At this stage, the manager applies basic techniques of psychodrama and some auxiliary techniques.

Group meeting: Group meeting is the stage where the group members share th feedback about their role at the end of the game staged in psychodrama. At this stage, the feedback and comments of the manager regarding the process are also very important.⁹ The content of the sessions is as follows:

Session 1: Meeting, Session 2: Exploring emotions, Session 3: What is my anxiety? What it looks like?, Session 4: Confronting anxiety and trust relationship in communication, Session 5: Understanding and accepting emotions, Session 6: What is empathy?, Session 7: Empathic communication and awareness, Session 8: Empathy with disabled people, Session 9: Empathy and trust relationship in communication, Session 10: Closure and farewell.

Data Collection Tools

Sociodemographic Information form

The form, which consists of questions that enquire the demographic information about the volunteers and their family, was prepared by the researchers and filled in by the volunteers themselves.

SAS-A

This scale, which was originally known as the Social Anxiety scale for children-revised, was adapted to Turkish by Aydın et al.²¹. The original scale consisted of a total number of 22 items. Its Turkish version consists of a total of 18 items in addition to 4 items concerning different subjects such as 'I like to read books', and 'I enjoy doing sports' (items 2, 7, 11 and 16). The SAS-A consists of three subscales containing 18 items. In order to determine the validity of the scale, the principal component factor analysis was applied according to the varimax rotation method. As a result of the structural factor analysis, three factors explaining 48% of the total variance were found. These three factors were named as fear of negative evaluation (FNE), social avoidance and feeling of restlessness in general social situations (SARGS) and social avoidance and restlessness in new situations (SARNS). A five-point Likert-type rating (never, rarely, occasionally, usually and always) was used in responses to the items of the scale and expressed with scores ranging from 1 to 5. Participants were asked to mark the intensity of the emotion they were experiencing, and stated in the items. Total score and subscale scores were obtained. The highest score obtained from the scale was 90 and the lowest score was 18. The high scores of the scale reflected high level of social anxiety. In the adaptation study, Aydın et al.²¹ calculated Cronbach's alpha internal consistency coefficients for the whole scale and its subdimensions in order to determine the reliability level of the scale. The reliability coefficients were 0.88 for the whole scale, 0.83 for the FNE, 0.68 for the SARGS and 0.71 for the SARNS. According to the data obtained from the sample group of this research, Cronbach's alpha internal consistency reliability coefficient values were calculated as 0.918 for the whole scale, 0.866 for the FNE, 0.845 for the SARGS and 0.720 for the SARNS subdimensions.

Ka-Si Empathic Tendency scale-adolescent

The Ka-Si Empathic Tendency scale for children and adolescents was developed by Kaya and Siyez²² in 2010 to determine the empathic tendencies of children. It consists of two factors: emotional empathy (10 items) and cognitive empathy (7 items). The Ka-Si Empathic Tendency scale-adolescent form was arranged in four grades: (1) Not at all suitable for me, (2) A slightly suitable for me, (3) Very suitable for me and (4) Totally suitable for me. The lowest and highest scores obtained from the emotional empathy (10 vs 40 pts), the cognitive empathy (7 vs 28 pts) and the empathic tendency (17 vs 68 pts) subdimension scores were indicated. The higher the scores, the more the decrease in empathic tendency. In the studies related to the construct validity of the adolescent form of the Ka-Si Empathic Tendency scale for children and adolescents, first factor analysis was performed one by one for each grade level, then the data of

6-12. Grades were combined and subsequently factor analysis was repeated.

The factor loadings of the 10 items in the emotional empathy subdimension ranged from 0.49 to 0.66 point; their correlations with the total subscale scores ranged from 0.51 to 0.83, and the correlations with scores from the Entire scale 0.47 to 0.70. As a result of the studies conducted for the reliability of the Empathic Tendency scale-adolescent form, Cronbach's alpha coefficient, which showed internal consistency, was 0.87 for the whole scale, 0.82 for emotional empathy, and 0.82 for cognitive empathy subdimensions. The test-retest reliability coefficient of the scale was found to be 0.75 for the whole scale, 0.73 for the emotional empathy and 0.69 for the cognitive empathy subdimensions.²²

Ethical Aspect of the Research

For this study, ethics committee's approval was received from TR Ministry of Health, The Provincial Directorate of Health, Taksim Training and Research Hospital Clinical Research Ethics Committee (no: 82). The institutional permission was obtained from the relevant training and research hospital. Written informed consent was obtained from the children and their parents who agreed to participate in the study. In the study, the principles of providing informed consent, protection of confidentiality and non-harm/utility were taken into consideration.

Statistical Analysis

SPSS 11.5 statistical program was used for data analysis. Data were analysed by descriptive analysis, Kolmogorov-Smirnov test, Mann-Whitney U test and Kruskal-Wallis test.

Limitations of the Study

The limitation of the research was that the research was carried out in a small sample group and cross-sectionally.

Result

Sociodemographic Characteristics

Patients included in our study were in the age of 13 (n=3; 33.3%), 14 (n=2; 22.2%), 15 (n=2; 22.2%), 16 (n=1; 11.1%) and 17 (n=1; 11.1%) years. The mean age of the patients was 14.4±1.42 years. Out of the total, 55.6% were female and 44.4% were male students.

When the total scores of the Empathic Tendency for Adolescents scale were examined, there was no statistically significant difference in the change of score before and after the psychodrama treatment. Although the scores obtained from the Adolescent scale of empathic tendency increased after psychodrama, this change was not statistically significant. Based on the results of the study, the scores tend to increase even if there is no statistically significant change (Table 1). When the total scores of the SAS-A were examined, it was found that there was a statistically significant difference in the change before and after psychodrama (p=0.029*). The social anxiety scale scores of adolescents after psychodrama were

statistically lower than those obtained before psychodrama (Table 2).

Discussion

In this research, which examined the effect of psychodrama on developing adolescents' empathy skills and reducing their social anxiety, statistical analysis and feedback of the members showed that the method was effective. In this study, no statistically significant differences were found in pre-psychodrama and post-psychodrama changes when total scores obtained from empathic tendencies for adolescents were analysed. Although the scores obtained from the Adolescent scale of empathic tendency increased after psychodrama, this change was not statistically significant. This result shows that psychodrama positively contributes to developing empathic skills. The studies in the literature support this study results. In her study with 23 undergraduate students, Dogan²³ stated that the students in the experimental group increased their empathy and counselling skills and self-awareness. In the study by Ozdag²⁴, he determined that psychodrama had a positive effect on students' empathic skills and stated that empathic skill levels of the experimental group increased compared to the control group. In a study, Dogan²⁵ emphasised the role of psychodrama in developing healthy and emotional relationships and stated that the patients had benefited greatly from psychodrama sessions, improved their behavioral and emotional insights and developed empathy skills. In a study conducted by Bairn et al.²⁶ using nine psychodrama methods to improve victim empathy in nine male individuals who had committed sexual crime in England, it was found that victim empathy levels increased significantly after the intervention (over 20 sessions). Ulupinar²⁷ stated in his study that there was an improvement in self-evaluation, awareness, expressing feelings and thoughts, creativity, empathy and problem-solving skills. Kalkan and Ozdel²⁸ found that empathic skills were developed in the study conducted by the elderly in the nursing home using psychodrama techniques. Similarly, Ozcan et al.²⁹ stated in their study that the empathic tendency skills increased in their mean scores after psychodrama practice. In the study by Oflaz et al.³⁰, nurses stated that psychodrama was

helpful in self-recognition, understanding others' perspective, developing their ability to express their own feelings and ideas and connecting with their feelings and ideas. Apart from these, other study results also showed that psychodrama had a positive effect on empathy skills.^{31,32} Unlike these results, Kaner³³ did not find a significant difference in the emotional aspect of empathy in the psychodrama study, but in cognitive dimension.

Empathic tendency is defined as the potential of empathy in an individual's daily life. It constitutes the emotional dimension of empathy and is an innate personality trait. Empathic tendency requires attitude development and personality change rather than a development in skills or cognitive functions. Because of these characteristics, it is very difficult to change the emphatic tendency, and as a result of the research, it is thought that the absence of excessive change in empathic tendency scores may have resulted from these reasons.^{34,35}

Social anxiety is included in the literature under the name of social anxiety. Findings obtained from the study were discussed accordingly. In this study, when the total scores obtained from the Social Anxiety scale were examined, a statistically significant difference was found between before and after psychodrama. There was a decrease in scale scores after psychodrama. According to these results, we can say that psychodrama is an effective method in reducing social anxiety in adolescents. These results are also compatible with the literature. Karatas¹³ in his study, in which adolescents constantly look at their anxiety levels with psychodrama practice, found significant difference between experimental and control groups. Purehsan and Sae³⁶ studied the effect of cognitive behavioral group therapy on reducing social anxiety among university students. It was observed that there was a big difference in the pre-test and post-test values of the experimental group and the study was concluded that cognitive behavioral group therapy was effective in improving the symptoms of social anxiety disorder (SAD). In Turkey, scarce number of studies assessed the relationship between SAD and psychodrama. Dogan³⁷ held 12 sessions of psychodrama on 21 graduate students and found that psychodrama

Table 1. The level of empathy in adolescents before and after psychodrama sessions

	Before psychodrama		After psychodrama		Difference		p
	X ± SD	Med (min-max)	X ± SD	Med (min-max)	X ± SD	Med (min-max)	
Empathic Tendency scale	51.56±4.67	52 (46-58)	56.67±10.21	61 (38-65)	-5.11±7.69	-8 (-14-10)	0.081 (t=-1.994)

Mann-Whitney U, Kruskal-Wallis test, SD: Standard deviation, min= Minimum, max: Maksimum

Table 2. Social anxiety level of the adolescents before and after psychodrama

	Before psychodrama		After psychodrama		Difference		p
	X ± SD	Med (min-max)	X ± SD	Med (min-max)	X ± SD	Med (min-max)	
Social Anxiety scale	61.67±16.81	59 (36-86)	48.78±8.12	49 (35-63)	12.89±14.52	11 (6-33)	0.029* (t=-2.663)

*p<0.05, Mann-Whitney U, Kruskal-Wallis test, SD: Standard deviation, min= Minimum, max: Maksimum

was effective in reducing anxiety levels of students. Although Uneri et al.¹⁹ examined the use of psychodrama in adolescents with SAD, since the objectives of their study were investigation of characteristics of psychodrama group treatment applications in adolescents, the strengths and weaknesses of the applications and the creation of psychodrama group model for adolescents, any statistical data could not be derived from this study for comparison. When we look at the literature, it is difficult to come across a study where social anxiety or SAD and psychodrama were dealt in combination. In the study, Dehnavi et al.³⁸ found that there was a significant decrease in the social anxiety symptoms among the adolescents in the experimental group compared to the control group in which male adolescents measured the effectiveness of psychodrama in reducing SAD. Akinsola and Udoka³⁹ found that psychodrama was effective in reducing anxiety levels in children. Oguzhanoglu et al.⁴⁰ stated that there was a decrease in social anxiety scores after the combined use of drug and psychodramatic group therapy. Gorkem and Tuccar⁴¹ found a decrease in anxiety levels of university students after 13 sessions of psychodrama.

In the study where Abeditehrani et al.⁴² used cognitive behavioral therapy and psychodrama together, they found a decrease in the symptoms of social anxiety. In another study where the effect of psychodrama on reducing anxiety in male schizophrenia patients was investigated, it was observed that psychodrama reduced anxiety symptoms in them.⁴³

The results of the study by Pratama et al.⁴⁴ are quite similar to the results of this study. Pratama et al.⁴⁴ found that psychodrama and sociodrama techniques are effective in developing emotional intelligence in their group counselling study. Their study states that individuals trust themselves to speak in front of others and have empathy to care more about the situation of others.

Conclusion

As a result of group therapy using psychodrama technique with adolescents, a significant decrease in the social anxiety levels of the participants and increase in their empathy scores were found when they were examined one by one. It was thought that the change in the empathic approach could be measured better by increasing the number of group members, and more comprehensive results could be provided with a registration method compatible with the qualitative analysis. Psychodrama is a psychotherapy method that exerts beneficial effects in solving problems peculiar to adolescence, concerns of success, social anxiety, problem solving in general and development of empathic skills. Since in our country inadequate number of studies have been performed in this field, primarily qualified relevant studies should be performed.

Ethics

Ethics Committee Approval: For this study, ethics committee's approval was received from TR Ministry of Health, The Provincial Directorate of Health, Taksim Training and Research Hospital Clinical Research Ethics Committee (no: 82).

Informed Consent: Written informed consent was obtained from the children and their parents who agreed to participate in the study.

Peer-review: Internally peer-reviewed.

Authorship Contributions

Concept: Ç.Ş., E.Y.Y., E.A., E.A.Y., Design: Ç.Ş., E.Y.Y., E.A., E.A.Y., Data Collection or Processing: Ç.Ş., Analysis or Interpretation: Ç.Ş., E.A., Literature Search: Ç.Ş., E.A., Writing: Ç.Ş., E.A.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Wittchen HU. The many faces of social anxiety disorder. *International Clinical Psychopharmacology*. 2000;(15 Suppl 1):7-12.
2. Teachman BA, Allen JP. Development of Social anxiety: Social Interaction Predictors of Implicit and Explicit Fear of Negative Evaluation. *Journal of Abnormal Child Psychology*. 2007;35:63-78.
3. Kashdan TB, Herbert JD. Social anxiety disorder in childhood and adolescence: current status and future directions. *Clin Child Fam Psychol Rev*. 2001;4:37-61.
4. La Greca AM. Friend of foes? Peer influences on anxiety among children and adolescents. In: Silverman WK, Treffers. *Anxiety disorders in children and adolescents; research, assessment, and intervention*. San Diego-California; Academic press Inc; 2001:159-186.
5. Jackson T. Protective self-presentation, sources of socialization, and loneliness among Australian adolescents and young adults. *Personality and Individual Differences*. 2007;43:1552-1562.
6. Rosen LM, Sheridan AM, Sambrook AK, Dennison JM, Jenness JM, Askren KM, Meltzoff NA, McLaughlin AK. Salience network response to changes in emotional expressions of others is heightened during early adolescence: relevance for social functioning. *Developmental Science*. 2018;21:e12571.
7. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005;62:617-627.
8. Evren C. Social anxiety disorder and alcohol use disorders. *Current Approaches in Psychiatry*. 2010;2:473-515.
9. Dökmen U. *Sociometry and psychodrama*. Istanbul: Sistem Publishing (4th ed). 2003;1-122.
10. Ozbek A, Leutz G. *Stage interaction in psychodrama group psychotherapy*. Ankara: Abdülkadir Ozbek Psychodrama Institute Publications; 2003.
11. Kellermann PF. *Focus on psychodrama: The therapeutic aspects of psychodrama*. London: Jessica Kingsley Publishers Ltd. 1992.
12. Hoag JM, Burlingame MG. Evaluating the effectiveness of child and adolescent group treatment: A meta-analytic review. *J Clin Child Psychol*. 1997;26:234-246.
13. Karatas Z, Gökçakan Z. Investigation of the effect of group practices using psychodrama techniques on reducing aggression in adolescents. *Turkish Journal of Psychiatry*. 2009;20:357-366.
14. Gökler B, Gökler I. *Group psychotherapy and psychodrama in children and adolescents*. Aysev AA, Taner IY, eds. *Child and Adolescent Mental Health and Diseases*. Istanbul; Golden Print; 2007:893-915.

15. Cakır V. Interactive TV and Interactive Drama. New Communication Environments and Interaction International Conference / 1-3 November 2006. https://www.researchgate.net/publication/292982267_ETKILESIMLI_TV_VE_ETKILESIMLI_DRAMA
16. Decety J, Ackson PL. The functional architecture of human empathy. *Behavioral and Cognitive Neuroscience Reviews*. 2004;3:71-100.
17. Tarhan N. *Community Psychology: From Social Schizophrenia to Social Empathy*, Istanbul; Timas Publishing; 2010.
18. Dökmen U. *Communication conflicts and empathy in art and daily life*. (5th ed). Istanbul; Remzi Bookstore; 2009:1-392.
19. Uneri OS, Akgün EY, Tanıdır C, Aytemiz T. The Use of Psychodrama in Adolescents with Social Anxiety Disorder. *Anatolian Journal of Psychiatry*. 2016;17:347-353.
20. Altınay D. *Psychodrama group psychotherapy handbook*. Istanbul; Sistem Publishing; 2003.
21. Aydın A, Sütcü TS, Sorias O. Examining the validity and reliability of the Social Anxiety Scale for Adolescents. *Journal of Child and Youth Mental Health*. 2007;14:79-89.
22. Kaya A, Siyez DM. KA-Sİ Empathic Tendency Scale for Children and Adolescents: Development, validity and reliability study. *Education and Science Journal*. 2010;35:110-125.
23. Dogan T. The effects of the psychodrama in instilling empathy and self-awareness: A pilot study. *PsyCh Journal*. 2018;7:227-238.
24. Ozdag S. Effects of Psychodrama Groups on Nursing School Students' Self-Esteem, Assertive Behavior, Empathic Tendency and Empathic Skill Levels. Unpublished Doctoral dissertation. Hacettepe University Institute of Health Sciences; Ankara; 1999:132.
25. Dogan T. The Role of Psychodrama in Developing Healthy Emotional Relations: A Case Study. *Turkish Psychological Counseling and Guidance Journal*. 2012;4:49-60.
26. Bairn C, Allam J, Eames T, Dunford S, Hunt S. The use of psychodrama to enhance victim empathy in sex offenders: An evaluation, *Journal of Sexual Aggression*. 1999;4:4-14.
27. Ulupınar S. The effect of psychodrama practice on nursing students' problem solving skills. *Anatolian Journal of Psychiatry*. 2014;15:55-62.
28. Kalkan NO, Ozdel O. A psychodrama study on old-age, nursing home and life journeys. *Turkish Psychiatry Journal*. 2005;16:124-132.
29. Ozcan KN, Bilgin H, Eracan N. The Use of Expressive Methods for Developing Empathic Skills. *Issues in Mental Health Nursing*. 2011;32:131-136.
30. Oflaz F, Meric M, Yuksel C, Ozcan CT. Psychodrama: An innovative way of improving self-awareness of nurses. *Journal of Psychiatric and Mental Health Nursing*. 2011;18:569-575.
31. Sangappa SB, Tekian A. Communication skills course in an Indian undergraduate dental curriculum: A randomized controlled trial. *Journal of Dental Education*. 2013;77:1092-1098.
32. Kesten KS. Role-play using SBAR technique to improve observed communication skills in senior nursing students. *The Journal of Nursing Education*. 2011;50:79-87.
33. Kaner S. Effect of psychodrama and reality therapy on the self concept and empathy levels of the young offenders. *Psychiatry Psychology Psychopharmacology Journal*. 1993;1:165-172.
34. Tutuk A, Al D, Dogan S. Investigation of Communication Skills and Empathy Levels of Nursing Students. *Cumhuriyet University School of Nursing Journal*. 2002;6:36-41.
35. Mete S, Gercek E. Investigation of Empathic Tendency and Skills of Nursing Students Studying with Problem Based Learning Management. *Cumhuriyet University School of Nursing Journal*. 2005;9:1-7.
36. Purehsan S, Sae O. Effectiveness Of Cognitive-Behavioral Group Therapy (CBGT) On Reduction Of Social Phobia". *Procedia Social and Behavioral Sciences*. 2010;1694-1697.
37. Dogan T. The effects of psychodrama on young adults' attachment styles. *The Arts in Psychotherapy*. 2010;37:112-119.
38. Dehnavi S, Hadadi Z, Jelveh M, Safaryazdi Z, Panahi P. The Effectiveness of Psychodrama in the Reduction of Social Anxiety Disorder among Male Adolescents. *The Neuroscience Journal of Shefaye Khatam*. 2014;2:52-53.
39. Akinsola FE, Udoka AP. Parental Influence on Social Anxiety in Children and Adolescents: Its Assessment and Management Using Psychodrama. *Psychology*. 2013;4:246-253.
40. Oguzhanoglu KN, Bayraktutan M, Varma SG, Ugurlu TT. Sosyal anksiyete bozukluğunda aleksitimi ve sempatik deri yanıtları üzerine ilaç ve psikodrama grup terapisinin etkileri. *Klinik Psikiyatri Dergisi*. 2019;22:452-462.
41. Görkem A, Tüccar L. The effect of psychodrama on university students' anxiety levels. *Turkish Studies*. 2018;13:599-620.
42. Abeditehrani H, Dijka C, Toghchi SM, Arntz A. Integrating Cognitive Behavioral Group Therapy and Psychodrama for Social Anxiety Disorder: An Intervention Description and an Uncontrolled Pilot Trial. *Clinical Psychology in Europe*. 2020;2:1-21.
43. Tehrani AH. Effectiveness of Psychodrama on Reduction of Anxiety among Male Schizophrenic Patients. *The Neuroscience Journal of Shefaye Khatam*. 2014;2:64.
44. Pratama SY, Wibowo EM, Awalya M. Group Counseling with Psychodrama and Sociodrama Techniques to Improve Emotional Intelligence. 2019;8:79-85.