

# Combining Spreader Grafts with Suture Suspension for Management of Narrow Internal Nasal Valve Angles

## *Dar İnternal Nazal Valv Açılarının Tedavisinde Spreader Greftlerle Sütür Süspansiyonunun Kombinasyonu*

Original Investigation  
Özgün Araştırma

Ahmed Ismail , Wael Hussein , Samy Elwany 

Department of Otorhinolaryngology, Alexandria University Alexandria School of Medicine, Alexandria, Egypt

### Abstract

**Objective:** The present study aimed to evaluate the long-term results of combining spreader grafts with lateral suspension techniques for management of narrow internal nasal valves.

**Methods:** This prospective study included 79 patients who underwent surgical interference for dysfunctional nasal valve. The surgical intervention was a combination of two commonly used techniques for management of internal nasal valve dysfunction: spreader graft insertion and lateral suspension of the upper lateral cartilages. The patients were assessed using the nasal obstruction symptom evaluation (NOSE) tool, computed tomography (CT) scans, and endoscopic examinations.

**Results:** All patients reported initial improvement and satisfaction in nasal breathing. Seventy-two patients

(91%) reported sustained relief of nasal breathing for at least 3 years postoperatively. The mean NOSE scores decreased from 69.8 to 20.65 postoperatively. The internal nasal valve angle, as measured endoscopically, increased from 7.23° to 13.05° on the right side and from 6.75° to 11.98° on the left side; the increase was statistically significant. The angle, as measured on axial CT scans, increased from 8.11° to 14.15° on the right side and from 6.98° to 12.13° on the left side; this increase, too, was statistically significant.

**Conclusion:** Combining spreader grafts with lateral suspension sutures provide good, stable, and sustained repair of dysfunctional narrowed internal nasal valves.

**Keywords:** Narrow nasal valve, nasal spreader graft, lateral cartilage suspension

### Öz

**Amaç:** Bu çalışmada dar internal nazal valv tedavisinde spreader greftlerin lateral süspansiyon teknikleriyle kombine edilmesinin uzun vadeli sonuçlarının değerlendirilmesi amaçlanmıştır.

**Yöntemler:** Bu ileriye dönük çalışma, disfonksiyonel nazal valv yönelik cerrahi uygulanan 79 hastayı kapsadı. Cerrahi uygulamada, internal nazal valv disfonksiyonunun tedavisi amaçlı yaygın olarak kullanılan iki teknik olan spreader greft uygulaması ile üst lateral kırıkdağın lateral süspansiyonu kombine edildi. Hastalar, burun tıkanıklığı semptom değerlendirmesi (NOSE) aracı, bilgisayarlı tomografi (BT) ve endoskopik muayene ile değerlendirildi.

**Bulgular:** Tüm hastalar başlangıçta burun solunumunda gelişme ve memnuniyet bildirdiler. Yetmiş iki

hasta (%91) postoperatif en az 3 yıllık sürede burun solunumundaki rahatlamının sürdüğünü belirtti. Ameliyat sonrası ortalama NOSE skorları 69.8'den 20.65'e düştü. Endoskopik olarak ölçülen iç nazal valv açısı sağ tarafta 7.23°den 13.05°e, sol tarafta 6.75°den 11.98°e yükseldi; artış istatistiksel olarak anlamlıydı. Aksiyel BT incelemelerinde ölçülen açı, sağ tarafta 8.11°den 14.15°e, sol tarafta 6.98°den 12.13°e yükseldi. Bu artış da istatistiksel olarak anlamlıydı.

**Sonuç:** Spreader greftleri lateral süspansiyon dikişleri ile birlikte uygulamak, fonksiyonel daralmanın olduğu internal nazal valvlerin iyi, stabil ve uzun süreli olarak onarılmasını sağlar.

**Anahtar kelimeler:** Dar nazal valv, nazal spreader greft, lateral kırıkdağ süspansiyonu



**ORCID IDs of the authors:**  
A.I. 0000-0001-8898-7222;  
W.H. 0000-0002-2571-7900;  
S.E. 0000-0002-1951-1842.

**Cite this article as:** Ismail A, Hussein W, Elwany S. Combining Spreader Grafts with Suture Suspension for Management of Narrow Internal Nasal Valve Angles. Turk Arch Otorhinolaryngol 2018; 56: 25-9

**Corresponding Author/Sorumlu Yazar:**  
Samy Elwany; samy.elwany@alexmed.edu.eg

**Received Date/Geliş Tarihi:** 27.10.2017

**Accepted Date/Kabul Tarihi:** 17.01.2018

© Copyright 2018 by Official Journal of the Turkish Society of Otorhinolaryngology and Head and Neck Surgery Available online at [www.turkarchotolaryngol.net](http://www.turkarchotolaryngol.net)

© Telif Hakkı 2018 Türk Kulak Burun Boğaz ve Baş Boyun Cerrahisi Derneği Makale metnine [www.turkarchotolaryngol.net](http://www.turkarchotolaryngol.net) web sayfasından ulaşılabilir.

DOI: 10.5152/tao.2018.2996

### Introduction

Proper breathing is one of the most important constituents and features of health and well-being. Several factors contribute to nasal airflow sufficiency. The internal nasal valve is a major factor affecting the amount and nature of nasal breathing; its dysfunction is commonly overlooked by physicians. A narrow internal nasal valve angle is a surgical challenge

for otolaryngologists and plastic surgeons. Several procedures have been described to manage narrow internal nasal valve angles with variable results. These procedures include spreader grafts, batten grafts, and lateral suspension of the upper lateral cartilages (1-5).

Being a functional problem and owing to the multistructural nature of the nasal frame, a precise ob-

jective measurement of nasal airflow is still a clinical challenge. Most studies used subjective assessment of nasal breathing to determine the success or failure of surgical procedures (6-8). Objective tools such as CT scan and nasal endoscopy have also been described for assessment of the internal nasal valve (9, 10).

The present study aimed to evaluate the long-term results of combining spreader graft with lateral suspension techniques for the management of narrow nasal valve regions. Subjective and objective tools were used in the evaluation process.

## Methods

This prospective study included 79 patients who underwent surgical correction of the narrow internal nasal valve region. Patients with significant hypertrophy of inferior turbinates were excluded from the study. All patients were operated upon and followed-up in a tertiary care university hospital from March 2009 to August 2014. Informed consent was obtained from all patients, and the study was approved by the Ethics Committee of the Alexandria University School of Medicine (11/18 - 10.02.2017).

Diagnosis was made via standard ear, nose, and throat examination, anterior rhinoscopy, and endoscopic nasal examinations. Narrowing of the nasal valve was confirmed using the Cottle's test and internal tenting test.

Subjective assessment was performed using the nasal obstruction symptom evaluation (NOSE) score (11). Baseline NOSE scores were obtained at a preoperative visit, and follow-up NOSE scores were obtained.

A 0° 4-mm rigid endoscope (Storz Endoscopic Systems, Germany) was used for endoscopic examination of the patient in the sitting position. The internal nasal valve was identified at the first point of junction between the upper lateral cartilage and the nasal septum. Usually, not more than 1 cm of the endoscope is introduced into the nose. A series of photographs were captured, and the most appropriate one was selected for measurements (Figure 1).

All patients underwent standard CT evaluation with a slice thickness of 0.6 mm in bone algorithm. The internal nasal valve angle was measured on axial cuts using the OsiriX viewer (Pixmeo SARL, Switzerland) (Figure 2).

## Surgical technique

The surgical intervention was a combination of two commonly used techniques for management of internal nasal valve dysfunction: spreader graft insertion and lateral suspension of the upper lateral cartilages.

An open rhinoplasty approach was used to expose the nasal vault in all patients (Figure 3). Bilateral mucoperichondrial septal flaps were elevated. Bilateral subperichondrial tunnels were then created under the upper lateral cartilages, and the upper lateral cartilages were carefully disarticulated from the septum.




Figure 1. a, b. (a) Preoperative (Pre) and (b) postoperative (Post) endoscopic measurements of the internal nasal valve angle. NS: nasal septum; ULC: upper lateral cartilage




Figure 2. a, b. (a) Preoperative (Pre) and (b) postoperative (Post) radiologic measurements of the internal nasal valve angle. NS: nasal septum; ULC: upper lateral cartilage




Figure 3. The nasal dorsum is exposed, and the nasal septum (NS) is separated from both the upper lateral cartilages (ULCs) through open rhinoplasty approach.

A well-sized septal cartilage graft was harvested to be used as spreader graft. In patients with previous septal surgery where septal cartilage harvest was not attainable, a conchal cartilage harvest was performed (12). Any necessary rhinoplastic procedures were subsequently performed.

## Spreader graft insertion

A spreader graft (2 cm×4 mm) was inserted on either sides of the nasal septal dorsum and secured in place with 5-0 Polydioxanone (PDS) sutures. The medial edge of the upper lateral car-




Figure 4. The spreader grafts (SG) are inserted and secured in place between the nasal septum (NS) and the upper lateral cartilages (ULCs).




Figure 5. Lateral suspension sutures are placed at a distance equal to half the width of the upper lateral cartilage (ULC). The curved arrows indicate the direction of suturing. The sutures are secured in the midline over the nasal bridge.



Figure 6. a, b. A male patient (a) with a crooked nose and nasal hump complaining of marked nasal obstruction, particularly on the right side. The same patient (b) 12 months after surgical intervention using a combined bilateral spreader graft insertion with lateral suspension flaring sutures to support both the upper lateral cartilages. Nasal breathing was improved after surgery.

tilages was then sutured to the spreader grafts and to each other in the midline, using the same type of sutures, creating a slightly wider nasal septal angle (Figure 4).

Flaring sutures were then performed using the lateral suspension technique (5). The upper lateral cartilages were bilaterally sutured at a distance equal to half the width of the cartilage over the nasal bridge using a 5-0 PDS suture (Figure 5).

Endoscopic examination of the nasal cavity was then performed to ensure adequate widening of the internal nasal valve angle. Mattress suturing of the nasal septal mucosa eliminating sub-perichondrial dead space, was performed using 5-0 polyglactin sutures; the columellar incision was finally closed using 5-0 polypropylene sutures.

The patients were followed-up weekly for a month, then monthly for 3 months, and every 6 months for up to 3 years. All measurements were repeated during the last follow up visit.

#### Statistical analysis

Statistical analyses were performed using the XLSTAT (Add-insoft, NY, USA). The Student's t-test was used to compare the preoperative and postoperative means.  $p < 0.05$  was considered statistically significant.

#### Results

The study included 79 patients with diagnosed narrowing of the internal nasal valve. The patients comprised 56 males (age, 18-54 years; mean age,  $31 \pm 3$  years) and 23 females (age, 18-42 years; mean age,  $28 \pm 4$  years).

Previous nasal surgery included nine septorhinoplasty procedures (11.5%), three septoplasty operations (3.8%), and five turbinate surgeries (6.4%).

All patients underwent internal nasal valve repair using a combination of spreader graft insertion and lateral suspension technique. Sixty-two patients underwent additional rhinoplastic procedures. Seventeen patients required harvesting of the auricular cartilage.

The average duration for the surgical procedure was 90 min ( $\pm 19$  min). The operative duration was longer in patients who had previous rhinoplastic procedures ( $105 \text{ min} \pm 23 \text{ min}$ ).

All patients reported initial improvement and satisfaction in nasal breathing. Seventy-two patients (91%) reported sustained relief of nasal breathing for at least 3 years postoperatively. Figure 6-8 show three patients at the end of their follow-up period. All seven patients who reported inadequate improvement of nasal obstruction had at least one previous nasal surgical procedure; five (6.4%) had septorhinoplasty procedures, and two (2.3%) had septoplasty procedures.

The results of the NOSE scale and the endoscopic and CT scan measurements of the internal valve angle are shown in Table 1.



Figure 7. a, b. A female patient (a) with a deviated nasal axis and droopy tip complaining of nasal obstruction, particularly on the left side. The same patient (b) 16 months after surgical intervention using a combined bilateral spreader graft insertion with lateral suspension flaring sutures. Nasal breathing was improved after surgery.



Figure 8. a, b. A female patient (a) with a nasal deformity complaining of marked nasal obstruction on both sides. The same patient (b) 32 months after surgical intervention using a combined bilateral spreader graft insertion with lateral suspension flaring sutures.

Table 1. Results of the NOSE scale and the endoscopic and computed tomography scan measurements of the internal valve angle

	Preoperative	Postoperative	p
NOSE* scale	58-89	17-24	0.012
Range (mean±SD)	(69.8±8.7)	(20.65±4.8)	
Endoscopic measurement (right)	5.93°-8.46°	11.34°-14.83°	0.032
Range (mean±SD)	(7.23°±1.05°)	(13.05°±0.91°)	
Endoscopic measurement (left)	4.99°-8.01°	10.16°-12.98°	0.019
Range (mean±SD)	(6.75°±1.34°)	(11.89°±1.16°)	
CT scan measurement (right)	6.83°-9.02°	12.79°-14.13°	0.020
Range (mean±SD)	(8.11°±1.14°)	(14.15°±1.17°)	
CT scan measurement (left)	5.01°-8.59°	10.46°-13.08°	0.015
Range (mean±SD)	(6.98°±1.03°)	(12.13°±1.24°)	

\*NOSE: nasal obstruction symptom evaluation score; CT: computed tomography; SD: standard deviation

The preoperative NOSE scale ranged from 58 to 89 with a mean of  $69.8 \pm 8.7$ . The mean postoperative NOSE scale measurement, during the last follow up visit, was  $20.65 (\pm 4.8)$  and was significantly better than the preoperative score ( $p=0.012$ ).

The internal nasal valve angle, as measured endoscopically, increased from  $7.23^\circ$  to  $13.05^\circ$ , on average, after surgery on the right side and from  $6.75^\circ$  to  $11.98^\circ$  on the left side, and this increase was statistically significant ( $p=0.023$  and  $0.019$  respectively). The angle, as measured on axial CT scans, on average increased from  $8.11^\circ$  to  $14.15^\circ$  after surgery on the right side and from  $6.98^\circ$  to  $12.13^\circ$  on the left side, and this increase, too, was statistically significant ( $p=0.020$ ,  $0.015$  respectively).

No significant complications were reported in the series. Prolonged postoperative edema was recorded in five patients (6.4%). No wound infection, keloid formation, septal hematoma/perforation, or complications related to the auricle were recorded in any of the patients.

## Discussion

Nasal valve dysfunction is a commonly neglected cause of chronic nasal obstruction that continues to be overlooked by several rhinologists (7,12). Kasperbauer and Kern (13) stated that knowledge regarding the structure and function of the nasal valve region is required by those who conduct nasal surgeries. The internal nasal valve is frequently overlooked by surgeons and not uncommonly compromised by nasal surgery, particularly rhinoplasty. Several techniques have been described to restore a functional patent internal nasal valve, and many times more than one technique is needed to repair dysfunctional nasal valves.

In the present study, we reevaluated a combination of the spreader graft technique, which widens the narrow nasal valve angle, with a suture suspension technique, which allows for more stable support of the upper lateral cartilages. Meyer et al. (14) described another suture technique to support the upper lateral cartilages.

Spreader grafts are best placed through an external approach. The advantage of this approach is that it allows precise secure placement of the graft in the internal nasal valve area (15). In addition, we used needles to secure the spreader grafts in position before applying the sutures, utilizing the idea proposed by Park et al. (16). Combining the suture technique and cartilage graft is useful in externally creating a symmetrical nose while internally obtaining a bisymmetrical cross-sectional nasal valve area.

The internal tenting test was positive in patients indicating a weak upper lateral cartilage. Postoperatively, all patients initially reported improved nasal breathing, including patients who were negative for the internal tenting test. These patients also showed sustained improvement in nasal breathing for the whole study period. This shows that even patients with no clinical evidence of a weak upper lateral cartilage would benefit from the lateral suspension technique.

Seven patients with at least one previous nasal surgical procedure showed less sustainability of their initial nasal breathing

improvement (five with previous rhinoplasty operation and two with previous septoplasty operations). Failure of improvement in nasal breathing may be attributed to excessive fibrosis following previous surgical interventions.

Previous studies have shown that either spreader grafts or lateral suspension technique alone may sometimes fail to improve nasal breathing in patients with dysfunctional nasal valves (17). Because most patients with nasal valve dysfunction actually have a combination of both narrow nasal valve angles together with weak upper lateral cartilages, tackling one side of the problem would not practically solve the breathing problem on a long-term basis. Even patients with clinically normal strength of the upper lateral cartilage may benefit from the upper lateral cartilage supporting sutures.

The main limitations of this study are the probable difficulties in selecting patients who are most suitable for the described procedure, as well as the difficulties in properly placing the sutures. Moreover, in subsequent studies, we will elaborate more on the standardization of the endoscopic and radiological measurements of the internal nasal valve angle; however, this study shows that combining the spreader grafts technique with lateral suspension of the upper lateral cartilages would ensure better and longer lasting results for the management of dysfunctional nasal valve region than any of the two techniques alone.

## Conclusion

Adding suspension sutures of the upper lateral cartilages to the spreader grafts technique provides more efficient and stable repair of the internal nasal valve even in patients with normal strength of the upper lateral cartilages. The described technique is simple and does not interfere with any other rhinoplastic procedures.

**Ethics Committee Approval:** Ethics committee approval was received for this study from the ethics committee of Alexandria University Alexandria School of Medicine (11/18 - 10.02.2017).

**Informed Consent:** Written informed consent was obtained from patients who participated in this study.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Concept - A.I., W.H., S.E.; Design - A.I., S.E.; Supervision - A.I., W.H., S.E.; Resource - A.I., W.H., S.E.; Materials - A.I., W.H., S.E.; Data Collection and/or Processing - A.I., S.E.; Analysis and/or Interpretation - A.I., W.H., S.E.; Literature Search - A.I., W.H., S.E.; Writing - A.I., W.H., S.E.; Critical Reviews - A.I., S.E.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study has received no financial support.

**Etik Komite Onayı:** Bu çalışma için etik komite onayı Alexandria Üniversitesi Tıp Fakültesi Etik Komitesi'nden alınmıştır (11/18 - 10.02.2017).

**Hasta Onamı:** Yazılı hasta onamı bu çalışmaya katılan hastalardan alınmıştır.

**Hakem Değerlendirmesi:** Dış bağımsız.

**Yazar Katkıları:** Fikir - A.I., W.H., S.E.; Tasarım - A.I., S.E.; Denetleme - A.I., W.H., S.E.; Kaynaklar - A.I., W.H., S.E.; Gereçler - A.I., W.H., S.E.; Veri Toplanması ve/veya İşlemesi - A.I., S.E.; Analiz ve/veya Yorum - A.I., W.H., S.E.; Literatür Taraması - A.I., W.H., S.E.; Yazıyı Yazan - A.I., W.H., S.E.; Eleştirel İnceleme - A.I., S.E.

**Çıkar Çatışması:** Yazarlar çıkar çatışması bildirmemişlerdir.

**Finansal Destek:** Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

## References

1. Elwany S, Thabet H. Obstruction of the nasal valve. *J Laryngol Otol* 1996; 110: 221-4. [\[CrossRef\]](#)
2. Clark JM, Cook TA. The "butterfly" graft in functional secondary rhinoplasty. *Laryngoscope* 2002; 112: 1917-25. [\[CrossRef\]](#)
3. André RF, Vuyk HD. Nasal valve surgery; our experience with the valve suspension technique. *Rhinology* 2008; 46: 66-9.
4. Rizvi S, Gauthier M. Lateralizing the collapsed nasal valves simplified: 10-Year Survey of a Simple Concealed Suture Technique. *Laryngoscope* 2011; 121: 558-61. [\[CrossRef\]](#)
5. Scuito S, Bernardeschi D. Upper lateral cartilage suspension over dorsal grafts: a treatment for internal nasal valve dynamic incompetence. *Facial Plast Surg* 1999; 15: 309-16. [\[CrossRef\]](#)
6. Constantinides M, Galli SK, Miller PJ. A simple and reliable method of pt evaluation in the surgical treatment of nasal obstruction. *Ear Nose Throat* 2002; 81: 734-7.
7. Schlosser RJ, Park SP. Surgery for the dysfunctional nasal valve cadaveric analysis and clinical outcomes. *Arch Facial Plast Surg* 1999; 1: 105-10. [\[CrossRef\]](#)
8. Mıman MC, Deliktaş H, Ozturan O, Toplu Y, Akarçay M. Internal nasal valve: revisited with objective facts. *Otolaryngol Head Neck Surg* 2006; 134: 41-7. [\[CrossRef\]](#)
9. Poetker DM, Rhee JS, Mocan BO, Michel MA. Computed tomography technique for evaluation of the nasal valve. *Arch Facial Plast Surg* 2004; 6: 240-3. [\[CrossRef\]](#)
10. Beriat K, Karadag D. The value of computed tomography in evaluation of internal nasal valve angle. *Med J Trakya Univ* 2010; 27: 270-4.
11. Stewart MG, Witsell DL, Smith TL, Weaver EM, Yueh B, Hanley MT. Development and validation of the nasal obstruction symptom evaluation (NOSE) scale. *Otolaryngol Head Neck Surg* 2004; 130: 157-63. [\[CrossRef\]](#)
12. Cervelli V, Spallone D, Bottini J, Silvi E, Gentile P, Curcio B, Pascali M. Alar batten cartilage graft: Treatment of internal and external nasal valve collapse. *Aesth Plast Surg* 2009; 33: 625-34. [\[CrossRef\]](#)
13. Kasperbauer JL, Kern EB. Nasal valve physiology. Implications in nasal surgery. *Otolaryngol Clin North Am* 1987; 20: 699-719.
14. Meyer R, Jovanovic B, Derder S. All about nasal valve collapse. *Aesthetic Plast Surg* 1996; 20: 141-51. [\[CrossRef\]](#)
15. André RF, Paun SH, Vuyk HD. Endonasal spreader graft placement as treatment for internal nasal valve insufficiency: no need to divide the upper lateral cartilages from the septum. *Arch Facial Plast Surg* 2004; 6: 36-40. [\[CrossRef\]](#)
16. Park C, Kim ES, Roh TS. Needle fixation technique in cartilage grafting. *Ann Plast Surg* 1999; 43: 331-3. [\[CrossRef\]](#)
17. Piriã T, Tikanto J. Unilateral and bilateral effects of nasal septum surgery demonstrated with acoustic rhinometry, rhinomanometry, and subjective assessment. *Am J Rhinol* 2001; 15: 127-33. [\[CrossRef\]](#)