

Rare Pathology of Sinonasal Tract; Eosinophilic Angiocentric Fibrosis

Sinonazal Bölgenin Nadir Görülen Patolojisi; Eozinofilik Anjiosentrik Fibrozis

Case Report
Olgu Sunumu

Bahar Kayahan, Cavid Cabbarzade, Serdar Özer, Rıza Önder Günaydın, Ömer Taşkın Yücel
Department of Otolaryngology, Faculty of Medicine, Hacettepe University, Ankara, Turkey

Abstract

Eosinophilic angiocentric fibrosis (EAF) is a relatively rare, benign and slowly progressive fibroinflammatory disorder, which mainly involves the upper respiratory tract. EAF was first described by Holmes and Panje in 1983. EAF is typically seen in young to middle-aged females and it has a slowly progressive nature. As well as the sinonasal region the nasal septum is mainly the most common localization of EAF, involvement of the larynx and orbita was also described in literature. Symptoms vary with the involved area but nasal obstruction, epistaxis and epiphora are the most common

complaints. The etiology of EAF is not clear and it is diagnosed by specific histopathological features. Histologically, the lesion is characterized by a perivascular, eosinophil-rich inflammatory infiltrate and progressive fibrosis which appears as a whirling “onion-skin” pattern. In this paper; a case of sinonasal EAF is reported with the clinical course, radiological and histological features, etiology and treatment of these lesions based on updated literature.

Key Words: Eosinophilic angiocentric, fibrosis, sinonasal

Özet

Eozinofilik anjiosentrik fibrozis (EAF); nadir görülen, üst solunum yolunu tutan benign ve yavaş ilerleme gösteren fibroinflamatuvar hastalıktır. İlk olarak Holmes ve Panje tarafından 1983 yılında tanımlanmıştır. EAF tipik olarak genç-orta yaşlı kadınlarda görülür; progresyonu oldukça yavaştır. Sinonazal bölge ve özellikle nasal septum en sık görülen tutulum yeri olmakla birlikte, larinks ve orbita tutulumu olduğunu belirten literatür bilgileri mevcuttur. Semptomları yerleşim yerine göre değişmekle birlikte sıklıkla burun tıkanıklığı, epistaksis ve epifora gibi belirtiler

izlenir. EAF etiolojisi bilinmemektedir ve spesifik histopatolojik özellikler ile tanı alır. Histolojik olarak lezyon, eozinofilden zengin perivasküler inflamasyon ve progresif “soğan kabuğu” paternine sahip fibrozis ile karakterizedir. Bu yazıda bir sinonazal bölge EAF olgusu klinik özellikleri, radyolojik ve histolojik bulguları, etyoloji ve tedavi özellikleri ile güncel literatür bilgileri eşliğinde sunulmuştur.

Anahtar Kelimeler: Eozinofilik anjiosentrik, fibrosis, sinonazal

Introduction

Eosinophilic angiocentric fibrosis (EAF) is a rare, benign and slowly progressive fibroinflammatory disorder (1, 2). EAF was first described by Holmes and Panje in 1983 as “intranasal granulome faciale” and in 1985 Roberts and McCann (2, 3) reported 3 similar intranasal cases with detailed histopathological behavior. Although EAF has a clear clinical and histological appearance, its pathogenesis is still controversial (2, 4). EAF typically presents in young to middle-aged females with a submucosal inflammatory, fibrosing tumor-like lesion causing slowly progressive upper airway symptoms. Histologically, the lesion is characterized by a perivascular, eosinophil-rich inflammatory infiltrate and progressive fibrosis which appears as a whirling “onion-skin” pattern (2-4). We report a case of sinonasal EAF and review the clinical course, radiological and histological features, etiology and treatment modalities based on updated literature.

Case Presentation

A fifty-five-year old female patient was admitted to our clinic with a history of swelling and pain in the right premaxillary and alar region and also some attacks of epistaxis from the right nasal passage. On her physical examination; a submucosal pinkish purple mass was detected which was located inferolaterally of the inferior turbinate. The mass was cystic-like on palpation. On her magnetic resonance imaging (MRI), a soft tissue mass was detected originating from the lateral portion of the inferior turbinate and lying laterally to the premaxillary region (Figure 1). Transnasal endoscopic excision of the mass was performed. On pathological examination, dense angiocentric “onion-skin” like fibrosis foci and eosinophil rich inflammation were detected, so it was diagnosed as “eosinophilic angiocentric fibrosis” (Figure 2a, b). She was not given any medical therapy after surgery. During postoperative follow-ups, an MRI was performed on the 3rd month, when there was



This study was presented as a poster at the 9th Turkish Rhinology Congress, 23-26 May 2013, Antalya, Turkey

Bu çalışma, 9. Türk Rinoloji Kongresi'nde poster olarak sunulmuştur, 23-26 Mayıs 2013, Antalya, Türkiye

Address for Correspondence/Yazışma Adresi:
Bahar Kayahan, Department of Otolaryngology, Hacettepe University, Faculty of Medicine, Ankara, Turkey
Phone: +90 312 305 42 24
E-mail: baharkayahan@hotmail.com
Received Date/Geliş Tarihi: 14.08.2013
Accepted Date/Kabul Tarihi: 13.12.2013
Available Online Date/Çevrimiçi Yayın Tarihi: 15.01.2014

© Copyright 2014 by Official Journal of the Turkish Society of Otorhinolaryngology and Head and Neck Surgery Available online at www.turkarchotolaryngol.net
© Telif Hakkı 2014 Türk Kulak Burun Boğaz ve Baş Boyun Cerrahisi Derneği Makale metnine www.turkarchotolaryngol.net web sayfasından ulaşılabilir.
DOI:10.5152/tao.2014.1387

no sign of recurrence of the intranasal mass and also the residual lesion on premaxilla, which could not be resected, was observed as stable (Figure 3a-c). The patient was informed about the rarity of the diagnosis and the conflicts of the treatment options and she gave an informed consent for this case report.

Discussion

Eosinophilic angiocentric fibrosis is an uncommon inflammatory fibrosing lesion of the upper airway tract (2-5). The sinonasal tract and mainly the nasal septum is the most common location for the EAF, laryngeal and orbital involvements are also described in the literature (1, 2, 4). It has been claimed that the slow progression and nonspecific symptoms account for its delayed presentation, and diagnosis is usually postponed for an average of approximately 5 years (2, 5). Symptoms may vary according to the involved site but are frequently seen as nasal obstruction, pain, epistaxis and epiphora (2, 5, 6). On physical examination, the EAF lesion is usually observed as a pink-purple mucosal surface with submucosal thickening without any specific appearance.

Pathogenesis of EAF is not well understood. Allergic etiology is proposed to be responsible due to predominance of eosinophil rich inflammation (2, 3, 7, 8). However, only some cases have a history of allergy in the literature. Also, EAF has been found to be refractory to steroids and the other anti-allergic therapies (7, 8).



Figure 1. Preoperative MRI showing the lesion
MRI: magnetic resonance imaging





Figure 2. a, b. Histopathologic staining showing the 'onion-skin' appearance of the lesion

Trauma was also thought to be a predisposing factor but this was not supported by cases in the literature (2). There has been some evidence that trauma may cause acceleration of disease progression. After surgical resection, EAF showed rapid progression in some cases (2, 8-10). Our case did not have any symptoms of allergy and she did not have prior nasal trauma or surgery either.

Imaging modalities, both magnetic resonance imaging (MRI) and computerized tomography (CT) are nonspecific and typically reveal a well circumscribed submucosal soft tissue density mass (2). On MRI, EAF is generally isointense on T1-weighted images with moderate heterogeneous enhancement after contrast administration. This heterogeneous contrast involvement is seen because of different stages of inflammation and fibrosis in the same lesion. Most lesions show hypointense signal intensity on T2-weighted images owing to fibrosis (1). On non-enhanced CT, EAF is usually seen as a homogenous, isodense soft tissue mass, rarely with calcifications (1). As lesions progress, they may cause adjacent bone remodelling such as thinning, slight absorption and sclerosis, but not an infiltrative pattern due to its benign nature (1, 2). On our patient's CT images, a homogenous soft tissue mass was observed which did not cause any bony destruction.

The histology of EAF is pathognomonic and characterized by perivascular inflammatory cell infiltration with progressive fibrosis around small vessels leading to the typical "onion-skin" pattern (3, 4). Eosinophils are the predominant cells with some degrees of plasma cells and lymphocytes of early inflammatory lesions. In the course of EAF, fibrosis becomes more visible around the microvasculature (2). The main differential diagnosis includes lesions with prominent eosinophilic infiltrates (2-4). Granuloma faciale, an inflammatory vascular reaction, presents clinically as papules localized almost on the face. Histologically, there is a polymorphous inflammatory infiltrate and it consists of neutrophils and eosinophils in the dermis (3, 4). Both histological similarities and coexistence in some cases suggest that EAF is a submucosal variant of granuloma faciale affecting mainly the upper respiratory tract, but there is no consensus about this (2, 8, 10). Absence of geographic necrosis, necrotizing vasculitis and granulomatous inflammation excludes Wegener's granulomatosis (WG) and Churg-Strauss



Figure 3. a-c. Postoperative MRI showing that there was not any recurrence of the disease
MRI: magnetic resonance imaging

syndrome (CS). Also, EAF lacks systemic vasculitis which is a participant of CS and mucosa over the EAF lesion does not show necrosis which is usually seen in WG lesions (4, 10, 11). Blood test positivity for c-ANCA and p-ANCA supports the diagnosis of WG and CS, respectively (4, 11). Both markers were found to be negative in our case.

Treatment of EAF has been a challenge as approximately 70% of patients have persistent or recurrent disease during follow up (2, 5). Medical therapy is found to be ineffective but may cause symptomatic relief (2, 8). Intranasal, intralesional or systemic corticosteroids, antihistamines, dapsone, hydroxychloroquine, azathiopurine and tamoxiphen are trial therapies in the literature but none of them were found to be as effective as surgery (2, 4, 5, 8). Surgical excision of the lesion should be complete resection as far as possible (2, 7). Nevertheless, only 30% of the cases do not have recurrence after surgery, most of the patients need multiple resections (2-4). Some authors suggest medical therapy after surgery but there is not enough evidence that this could prevent any recurrence (2, 4).

Conclusion

EAF is a benign, progressive fibrosing disease with an elusive etiology. It has a nonspecific clinical presentation but a typical histological appearance that leads to diagnosis. Treatment remains controversial but as more cases are reported it is likely that a clear understanding of etiology and management will result in better outcomes.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has received no financial support.

Peer-review: Externally peer-reviewed.

Informed Consent: Written informed consent was obtained from patients who participated in this study.

Author Contributions: Concept - B.K., C.C.; Design - B.K., C.C.; Supervision - S.Ö., Ö.G., Ö.T.Y.; Funding - B.K., C.C.; Materials - B.K., C.C.; Data Collection and/or Processing - B.K., C.C.; Analysis and/or Interpretation - S.Ö., Ö.G., Ö.T.Y.; Literature Review - B.K., C.C.; Writing - B.K., C.C.; Critical Review - S.Ö., Ö.G., Ö.T.Y.; Other- B.K., C.C.

Çıkar Çatışması: Yazarlar herhangi bir çıkar çatışması bildirmemişlerdir.

Finansal Destek: Yazarlar bu çalışma için finansal destek almadıklarını beyan etmişlerdir.

Hakem Değerlendirmesi: Dış bağımsız.

Hasta Onamı: Yazılı hasta onamı bu çalışmaya katılan hastalardan alınmıştır.

Yazar Katkıları: Fikir - B.K., C.C.; Tasarım - B.K., C.C.; Denetleme - S.Ö., Ö.G., Ö.T.Y.; Kaynaklar - B.K., C.C.; Malzemeler - B.K., C.C.; Veri toplanması ve/veya işlemesi - B.K., C.C.; Analiz ve/veya Yorum - S.Ö., Ö.G., Ö.T.Y.; Literatür taraması - B.K., C.C.; Yazıyı Yazan - B.K., C.C.; Eleştirel İnceleme - S.Ö., Ö.G., Ö.T.Y.; Diğer - B.K., C.C.

Kaynaklar

1. Yang BT, Wang YZ, Wang XY, Wang ZC. Nasal Cavity Eosinophilic Angiocentric Fibrosis: CT and MR Imaging Findings, *AJNR Am J Neuroradiol* 2011; 32: 2149-53. [\[CrossRef\]](#)
2. Sunde J, Alexander KA, Reddy VV, Woodworth BA. Intranasal eosinophilic angiocentric fibrosis: a case report and review, *Head Neck Pathol* 2010 4: 246-8. [\[CrossRef\]](#)
3. Onder S, Sungur A. Eosinophilic angiocentric fibrosis: an unusual entity of the sinonasal tract, *Arch Pathol Lab Med* 2004; 128: 90-1.
4. Jain R, Robblee JV, O'Sullivan-Mejia E, Lea J, Heller A, Faquin WC, et al. Sinonasal eosinophilic angiocentric fibrosis: A report of four cases and review of literature, *Head Neck Pathol* 2008; 2: 309-15. [\[CrossRef\]](#)
5. Kosarac O, Luna MA, Ro JY, Ayala AG. Eosinophilic angiocentric fibrosis of the sinonasal tract, *Ann Diagn Pathol* 2008; 12: 267-70. [\[CrossRef\]](#)
6. Valenzuela AA, Whitehead KJ, Brown I, Sullivan TJ. Eosinophilic angiocentric fibrosis: an unusual entity producing complete lacrimal duct obstruction, *Orbit* 2006; 25: 159-61. [\[CrossRef\]](#)
7. Watanabe N, Moriwaki K. Atypical eosinophilic angiocentric fibrosis on nasal septum, *Auris Nasus Larynx* 2006; 33: 355-8. [\[CrossRef\]](#)
8. Paun S, Lund VJ, Gallimore A. Nasal fibrosis: long-term follow up of four cases of eosinophilic angiocentric fibrosis, *J Laryngol Otol* 2005; 119: 119-24. [\[CrossRef\]](#)
9. Tabae A, Zadeh MH, Proytcheva M, LaBruna A. Eosinophilic angiocentric fibrosis, *J Laryngol Otol* 2003; 117: 410-3. [\[CrossRef\]](#)
10. Burns BV, Roberts PF, De Carpentier J, Zarod AP. Eosinophilic angiocentric fibrosis affecting the nasal cavity: A mucosal variant of the skin lesion granuloma faciale, *J Laryngol Otol* 2001; 115: 223-6. [\[CrossRef\]](#)
11. Slovik Y, Putterman M, Nash M, Sion-Vardy N. Eosinophilic angiocentric fibrosis of the sinonasal tract in a male patient with chronic bowel inflammation, *Am J Rhinol* 2006; 20: 91-4.