



© Osman Uzundere,
© Cem Kıvılcım Kaçar,
© Meral Erdal Erbatur,
© Mehmet Salih Gül,
© Mesut Akgündüz,
© Zeki Korhan,
© Ömer Can

Factors Affecting the Mortality of Patients in Critical Condition with Coronavirus Disease-2019 in the Intensive Care Unit

Koronavirüs Hastalığı-2019 Nedeniyle Yoğun Bakım Ünitesinde Yatan Kritik Hastalarda Mortaliteye Etki Eden Faktörler

Received/Geliş Tarihi : 27.01.2021
Accepted/Kabul Tarihi : 23.03.2021

© Copyright 2021 by Turkish Society of Intensive Care
Turkish Journal of Intensive Care published by Galenos Publishing House.

Osman Uzundere, Cem Kıvılcım Kaçar, Meral Erdal Erbatur, Mehmet Salih Gül, Mesut Akgündüz, Zeki Korhan, Ömer Can

University of Health Sciences Turkey, Diyarbakır Gazi Yaşargil Training and Research Hospital, Clinic of Anesthesiology and Reanimation, Diyarbakır, Turkey

Osman Uzundere MD, (✉),
University of Health Sciences Turkey, Diyarbakır Gazi Yaşargil Training and Research Hospital, Clinic of Anesthesiology and Reanimation, Diyarbakır, Turkey

E-mail : osmanuzundere@hotmail.com

Phone : +90 412 258 00 60

ORCID ID : orcid.org/0000-0002-5968-4561

ABSTRACT Objective: This study aimed to determine the factors affecting the mortality of patients in critical condition with coronavirus disease-2019 (COVID-19) in the intensive care unit (ICU).

Materials and Methods: We included a total of 445 patients who are admitted in the ICU due to COVID-19. Patients were divided into two groups-those who survived and those who died during the ICU follow-up-and their demographic, clinical and laboratory characteristics were compared. Factors affecting mortality were also determined.

Results: Older age, high Kidney Disease: Improving Global Outcome (KDIGO) stage and Sequential Organ Failure Assessment (SOFA) scores at first admission to the ICU, high neutrophil/lymphocyte ratio, high D-dimer levels, low bicarbonate (HCO_3) values and high lactate dehydrogenase (LDH) and creatinine levels were determined as independent risk factors for mortality in patients in critical condition with COVID-19 admitted in the ICU. Particularly, a substantial relationship was observed between the KDIGO stage and mortality during the ICU admission.

Conclusion: Age, KDIGO stage and SOFA scores at first admission, neutrophil/lymphocyte ratio and D-dimer, HCO_3 , LDH and creatinine levels were independent risk factors for mortality in patients in critical condition with COVID-19 admitted in the ICU.

Keywords: Coronavirus disease 2019, ICU, kidney disease: improving global outcome, mortality, sequential organ failure assessment score

ÖZ Amaç: Bu çalışmada yoğun bakım ünitesinde (YBÜ) yatan kritik koronavirüs hastalığı-2019 (COVID-19) hastalarını demografik, klinik ve laboratuvar özellikleri açısından karşılaştırıp mortaliteye etkili olan faktörleri saptamayı amaçladık.

Gereç ve Yöntem: Çalışmaya YBÜ'de COVID-19 nedeniyle yatan 445 hasta dahil edildi. Hastalar YBÜ'de takipleri sırasında mortalite gelişmeyenler ve mortalite gelişenler olarak iki gruba ayrılıp demografik, klinik ve laboratuvar özellikleri açısından karşılaştırıldı ve mortaliteye etki eden faktörler saptanmaya çalışıldı.

Bulgular: İleri yaş, YBÜ'ye ilk yatıştaki yüksek Böbrek Hastalıkları: Küresel Sonuçların İyileştirilmesi (KDIGO) evresi ve Sıralı Organ Yetmezliği Değerlendirmesi (SOFA) skorları, yüksek nötrofil lenfosit oranı, yüksek D-dimer düzeyleri düşük bikarbonat (HCO_3) değerleri, yüksek laktat dehidrogenaz (LDH) düzeyleri ve yüksek kreatinin düzeyleri YBÜ'de yatan kritik COVID-19 hastalarında mortalite için bağımsız risk faktörleri olarak saptandı. Özellikle YBÜ'ye başvuru esnasındaki KDIGO evresiyle mortalite arasındaki ilişki dikkat çekiciydi.

Sonuç: YBÜ'de yatan kritik COVID-19 hastalarında yaş, ilk yatıştaki KDIGO ve SOFA skorları, nötrofil lenfosit oranı, D-dimer, HCO_3 , LDH ve kreatinin mortalite için bağımsız risk faktörleridir.

Anahtar Kelimeler: Koronavirüs hastalığı-2019, YBÜ, böbrek hastalıkları: küresel sonuçların iyileştirilmesi, mortalite, sıralı organ yetmezliği değerlendirilmesi

Introduction

Based on the data published by the World Health Organization (WHO) on December 12, 2020, the coronavirus disease-2019 (COVID-19) pandemic, in which 69.5 million individuals were infected and 1,582,674 individuals have died, continues to be an issue worldwide (1). Although vaccination studies, which have recently accelerated, are a hope, approximately 15% of the patients with COVID-19 develop critical illnesses requiring oxygen support. In approximately 5% of the patients, respiratory failure secondary to acute respiratory distress syndrome (ARDS) as well as numerous complications including sepsis and septic shock, thromboembolism, renal failure, and cardiac damage, may further develop into a critical illness (2). The disease mortality can be extremely high, particularly due to the complications that may develop in the critical patient group. In the initial publications, it was stated that in-hospital mortality due to severe acute respiratory syndrome coronavirus 2 was approximately 28% (3). Moreover, it was emphasized that mortality was higher in critically ill patients (76%) hospitalized in the intensive care unit (ICU) (4).

When the publications on mortality in patients with COVID-19 were examined, there were reportedly numerous factors that could affect the clinical course and patient mortality (3,5-12). Among these factors, patient characteristics, male sex, advanced age, obesity, smoking, and comorbid diseases (particularly diabetes and hypertension) as well as Acute Physiology and Chronic Health Evaluation-II (APACHE-II) and Sequential Organ Failure Assessment (SOFA) score and several laboratory values are reportedly associated with mortality in patients with COVID-19 (4,12-14).

Currently, no effective treatment has been discovered for managing the COVID-19 epidemic, which has affected the world to a substantial extent (15). Determining the factors that affect mortality remains an important concern in terms of decreasing mortality due to the disease. In the current study that was planned with considering this notion, we aimed to perform a comparative assessment of critical patients with COVID-19 who were followed up in ICUs in our region since the beginning of the pandemic in terms of demographic, clinical, and laboratory characteristics and to determine the factors that affect mortality in this patient group.

Materials and Methods

Study Design, Population, and Data

Critical patients hospitalized due to COVID-19 in ICU of the University of Health Sciences Turkey, Diyarbakır Gazi Yaşargil Training and Research of Hospital between March 22 and September 1, 2020, were included in this study. The study was approved by the Clinical Research Ethics Committee of University of Health Sciences Turkey, Gazi Yaşargil Training and Research of Hospital (decision no: 550, date:11.09.2020). The trial was registered with clinicaltrials.gov (NCT04659876). This retrospective cohort study was conducted in accordance with the 2008 Declaration of Helsinki criteria.

Critical patients diagnosed with COVID-19 on the dates specified, followed up in ICU, aged >18 years, in serious need of oxygen support according to WHO (2) and the temporary guidelines of T.C. Science Board of the Ministry of Health [presence of fever, muscle/joint pain, cough, and sore throat; tachypnea (30 breaths/min) or dyspnea; use of extra respiratory muscles; SpO₂ level below of ≤90% in room air; bilateral diffuse pneumonia symptom detected on chest radiography or computerized tomography (CT); and PaO₂/FiO₂ ratio of <300], and developed or had complications including severe pneumonia, ARDS, sepsis/septic shock, and acute renal failure were included in the study (16). Patients with COVID-19 aged <18 years with mild-to-moderate symptoms, no respiratory distress, and no signs of diffuse pneumonia on chest X-ray or CT as well as ICU patients excepted from COVID-19 diagnosis were excluded from the study. In addition, patients whose complete data could not be accessed from the hospital system or the patient file records were excluded. When the patients were admitted to ICU for the first time, their clinical conditions were evaluated with APACHE-II and SOFA scores, and the degree of renal failure was evaluated using the Kidney Disease: Improving Global Outcomes (KDIGO) classification (17).

Age; sex; comorbidity; ABO and Rh blood groups; APACHE-II and SOFA scores and KDIGO stage during admission to ICU; hemogram parameters [white blood cell (WBC), neutrophil, lymphocyte, neutrophil/lymphocyte (N/L) ratio, hemoglobin, hematocrit, and platelet count]; blood gas values [pH, partial oxygen pressure (PO₂), partial carbon dioxide pressure (PCO₂), bicarbonate (HCO₃), and lactate]; coagulation parameters [prothrombin time (PTZ)

and D-dimer]; blood biochemistry results [creatinine kinase (CK), lactate dehydrogenase (LDH), C-reactive protein (CRP), urea, creatinine, alanine aminotransferase, aspartate aminotransferase (AST), total bilirubin, direct bilirubin, and indirect bilirubin]; and procalcitonin (PCT) and ferritin levels of the patients were recorded. Moreover, the length of stay in ICU and whether the patient died or survived were recorded. Patient data were rechecked for erroneous information before the last data entry and entered into a computerized database.

Patients were divided into two groups-those who survived (survivors) and those who died (non-survivors) during ICU follow-up. Both groups were compared in terms of clinical characteristics; APACHE-II and SOFA scores and KDIGO stage; and laboratory values at the first admission to ICU. We attempted to determine the factors that affect mortality in critically ill patients hospitalized in ICU with COVID-19 diagnosis.

Statistical Analysis

SPSS 16.0 software for Windows (SPSS Inc., Chicago, IL, USA) was used for statistical analysis. Continuous data were expressed as means (SD or minimum-maximum), and categorical data were expressed as frequencies with percentages. Comparison of categorical data in the groups was performed using chi-square and Fisher's Exact test, and the results were presented as n%. Kolmogorov-Smirnov test was used to determine whether the numerical data fit the normality distribution. Data

conforming to the normality distribution were evaluated using Student's t-test, and Mann-Whitney U and Kruskal-Wallis tests were employed to compare data that did not fit the normality distribution. Binary logistic regression was performed for the risk factors that were found to be significant in the univariate analysis. Odds ratio (OR) with 95% confidence interval (CI) was used to report the association between mortality and exposure to the risk factors. In all comparisons, p value of <0.05 was considered significant.

Results

Overall, data of 474 patients were accessed in the study. According to the exclusion criteria, 29 patients were excluded, and the study was completed with 445 patients. The mean [standard deviation (SD)] age of the patients was 68.5 (15.1) years; 232 (52.1%) patients were male and 213 (47.9%) were female. Of the patients included in the study, 338 (76%) had at least one comorbid disease. The most common comorbid diseases were hypertension (40.2%) and diabetes (28.5%). Further, 296 patients died during their follow-up period in ICU and mortality was 66.5%. The mean (SD) length of stay of patients in ICU was 11.2 (10.7) days. The demographic, clinical, and laboratory characteristics of patients are detailed in Table 1.

Table 1. Demographic, clinical, and laboratory characteristics of patients hospitalized in the intensive care unit due to COVID-19

Characteristic	All patients (n=445) mean (min-max)	Survivors (n=149) mean (min-max)	Non-survivors (n=296) mean (min-max)	p-value
Age (year)	68.5 (18-100)	62.7 (22-95)	71.4 (18-100)	<0.001
Sex				
Female	213 (47.9%)	83 (55.7%)	130 (43.9%)	0.019
Male	232 (52.1%)	66 (44.3)	166 (56.1%)	
Blood group				
A	225 (50.6%)	74 (49.7%)	151 (51%)	0.48
B	71 (16%)	21 (14.1%)	50 (16.9%)	
AB	33 (7.4%)	9 (6%)	24 (8.1%)	
O	116 (26.1%)	45 (30.2%)	71 (24%)	
Rh factor				
Negative	61 (13.7%)	21 (12.7%)	42 (14.2%)	0.87
Positive	384 (86.3%)	130 (87.3)	254 (85.8%)	

Table 1. Continued				
Characteristic	All patients (n=445) mean (min-max)	Survivors (n=149) mean (min-max)	Non-survivors (n=296) mean (min-max)	p-value
Comorbidities				
No	107 (24%)	42 (28.2%)	65 (22%)	0.14
Yes	338 (76%)	107 (71.8%)	231 (78%)	
Diabetes	127 (28.5%)	38 (25.5%)	89 (30.1%)	0.31
Hypertension	179 (40.2%)	64 (43%)	115 (38.3%)	0.53
COPD	45 (10.1)	19 (12.8%)	26 (8.8%)	0.19
CKD	34 (7.6%)	12 (8.1%)	22 (7.4%)	0.81
CVD	66 (14.8%)	19 (12.8%)	47 (15.9%)	0.38
KDIGO score				
0	189 (42.4%)	113 (75.8%)	76 (25.6%)	<0.001
1	83 (18.7%)	17 (11.4%)	66 (22.3%)	
2	79 (17.8%)	12 (8.1)	67 (22.6%)	
3	94 (21.1%)	7 (4.7)	87 (29.4%)	
APACHE-II score	16.61 (2-49)	13.1 (2-33)	18.3 (2-49)	<0.001
SOFA score	4.34 (1-17)	3.3 (1-12)	4.8 (1-17)	<0.001
Laboratory				
White blood cells ($\times 10^3/\mu\text{L}$)	11.33 (1.13-57.4)	10.6 (2.95-42.7)	11.6 (1.13-57.4)	0.032
Neutrophil ($\times 10^3/\mu\text{L}$)	9.51 (0.66-37.5)	8.75 (1.3-37.5)	9.9 (0.66-34.4)	0.004
Lymphocyte ($\times 10^3/\mu\text{L}$)	0.98 (0.14-3.59)	1.1 (0.19-3.59)	0.93 (0.14-3.5)	<0.001
Neutrophil/lymphocyte ratio	12.8 (0.12-87.14)	9.4 (0.33-60.04)	14.5 (0.12-87.14)	<0.001
Hemoglobin (g/dL)	12.8 (5.6-19.2)	13.06 (5.9-17)	12.6 (5.6-19.2)	0.015
Hematocrit (%)	40.61 (17.8-61.6)	41.2 (20.2-55.9)	40.2 (17.8-61.6)	0.039
Platelet ($\times 10^3/\mu\text{L}$)	242.5 (30-671)	253.4 (84-671)	237.03 (30-628)	0.048
Prothrombin time (s)	13.83 (9.7-34.6)	13.3 (9.9-22.5)	14.1 (9.7-54.9)	0.011
D-dimer (ng/mL)	2019.2 (8.4-44498)	954.5 (75-16948)	2564.4 (8.4-44498)	<0.001
pH	7.36 (6.82-7.55)	7.38 (6.91-7.54)	7.36 (6.82-7.55)	0.01
PO ₂ (mmHg)	41.37 (13.5-206)	42.14 (17.9-162)	40.3 (13.5-198)	0.76
PCO ₂ (mmHg)	38.61 (20-115)	39.1 (16.9-108)	38.2 (20-115)	0.12
HCO ₃ (mmol/L)	21.78 (5.3-32.1)	23.02 (5.9-31.5)	21.16 (5.3-32.1)	<0.001
Lactate (mmol/L)	2.76 (0.6-26)	2.2 (0.6-8.2)	3.04 (0.7-26)	<0.001
Lactate dehydrogenase (U/L)	514.9 (99-4500)	406.7 (139-1079)	569.4 (99-4500)	<0.001
Creatine kinase (IU/L)	314.05 (0.32-14952)	204.4 (11-2949)	369.4 (0.32-14952)	<0.001
C-reactive protein (mg/L)	141.2 (2-350)	120.4 (2-350)	151.7 (2-350)	<0.001
Blood urea nitrogen (mg/dL)	64.1 (8-280)	47.6 (8-267)	72.5 (13-280)	<0.001
Creatinine (mg/dL)	1.55 (0.36-21.8)	1.28 (0.44-10.4)	1.68 (0.36-21.8)	<0.001
ALT (U/L)	42.6 (6-1254)	32.2 (6-442)	47.9 (6-1254)	0.13
AST (U/L)	67.9 (7-3444)	40.8 (9-518)	81.5 (7-3444)	<0.001
Total bilirubin (mg/dL)	0.73 (0.12-6.8)	0.68 (0.14-3.69)	0.75 (0.12-6.8)	0.20
Direct bilirubin (mg/dL)	0.38 (0.1-4.7)	0.34 (0.1-2.31)	0.4 (0.1-4.7)	0.02

Table 1. Continued

Characteristic	All patients (n=445) mean (min-max)	Survivors (n=149) mean (min-max)	Non-survivors (n=296) mean (min-max)	p-value
Indirect bilirubin (mg/dL)	0.33 (0.01-2)	0.33 (0.03-1.86)	0.33 (0.01-2)	0.93
Procalcitonin (ng/mL)	3.19 (0.02-100)	1.39 (0.02-62.8)	4.13 (0.03-100)	<0.001
Ferritin (µg/L)	854.8 (5.86-2000)	673.8 (5.86-2000)	951.5 (16.6-2000)	<0.001
Length of stay in the intensive care unit (day)	11.2 (1-91)	13.02 (1-91)	10.3 (1-79)	0.004

COPD: Chronic obstructive pulmonary disease, CKD: chronic kidney disease, CVD: cardiovascular disease, KDIGO: Kidney Disease: Improving Global Outcomes, APACHE-II: Acute Physiology and Chronic Health Evaluation-II, SOFA: Sequential Organ Failure Assessment, PO₂: partial oxygen pressure, PCO₂: partial carbon dioxide pressure, HCO₃: bicarbonate, ALT: alanine aminotransferase, AST: aspartate aminotransferase, min: minimum, max: maximum, COVID-19: coronavirus disease-2019

Univariate Analysis

Patients were divided into two groups-those who survived (survivors =149, 33.5%) and those who died (non-survivors =296, 66.5%) during ICU follow-up-and compared. In terms of demographic and clinical characteristics, the mean patient age of the non-survivor group was higher (71.4 vs. 62.7 years; $p<0.001$). Mortality was higher than survival in male patients (56.1% vs. 43.9%; $p=0.019$). Patients with KDIGO stage 1, 2, and 3 showed higher mortality than expected ($p<0.001$). Further, patients who died were found to have higher APACHE-II and SOFA scores ($p<0.001$; $p<0.001$) (Table 1).

On comparing both groups in terms of laboratory values at the first admission to ICU, the non-survivors showed a significant higher N/L ratio, WBC, neutrophil, PTZ, D-dimer, lactate, LDH, CK, CRP, urea, creatinine, AST, direct bilirubin, PCT, and ferritin values and lower lymphocyte, hemoglobin, hematocrit, platelet, pH, and HCO₃ values. Details and significance values of the comparison between both groups are shown in Table 1.

Risk Factors for Mortality in ICU Patients with COVID-19

Results of the binary logistic regression are shown in Table 2. Advanced age (OR: 1.03; 95% CI: 1.008-1.055), KDIGO stage 1 (OR: 5.23; 95% CI: 2.490-10.97), KDIGO stage 2 (OR: 7.07; 95% CI: 2.9-17.24), KDIGO stage 3 (OR: 33.98; 95% CI: 8.860-130.3), high SOFA score (OR: 1.194; 95% CI: 1.007-1.416), high N/L ratio (OR: 1.069; 95% CI: 1.006-1.137), high D-dimer levels (OR: 1.000; 95% CI: 1.0-1.001), low HCO₃ values (OR: 0.888; 95% CI: 0.802-0.983), high LDH levels (OR: 1.004; 95% CI: 1.002-1.006), and elevated creatinine levels (OR: 0.499; 95% CI: 0.368-0.676)

were identified as independent risk factors for mortality in critical COVID-19 patients hospitalized in ICU.

Discussion

In the present study that evaluated the factors affecting mortality in critical patients with COVID-19 followed up in ICU, the mortality was determined to be 66.5%. Moreover, advanced age; high KDIGO stage and SOFA scores at the first admission to ICU; N/L ratio; D-dimer, LDH, and creatinine levels; and low HCO₃ value were determined as independent risk factors affecting mortality in this critical patient group.

Most studies conducted on patients with COVID-19 have emphasized that advanced age is an independent risk factor for mortality (6,7,9,11,18-20). With increasing age, compared with young individuals, stronger host innate responses to viral infections, decreased type 1 interferon expression, age-related defects in T and B cell functions, and excessive type 2 cytokine production result in deficient response to viral infections and prolonged proinflammatory responses, which are considered as the causes of increased mortality risk in older aged patients with COVID-19 (3). In the present study, mean patient age of the non-survivors was 71.4 (18-100) years, and similar to previous studies, advanced age was determined as an independent risk factor for COVID-19.

Numerous studies have reported that male patients with COVID-19 exhibit a more severe disease, and the mortality risk in the male sex is higher (9,14,20,21). The high mortality in males has been attributed to higher chronic comorbidities, such as cardiovascular disease, hypertension, and lung disease, and smoking rate (20). In the present study, the patient group with a mortal course of the disease exhibited a predominance of male population, in accordance

with the literature. However, as a result of the logistic regression analysis, it was determined that male gender is not an independent risk factor for critical COVID-19 patients hospitalized in the ICU. This unexpected result contradicts the studies in the literature. This result may be due to the single-center nature of our study and the limited number of patients.

On literature review, another risk factor that can affect the clinical course of the disease in patients with COVID-

19 is the presence of comorbidities. In the meta-analysis by Martins-Filho et al. (6), the authors emphasized that the presence of comorbidities in patients with COVID-19 resulted in a 1.6 times increase in the in-hospital mortality. In a study conducted by COVID-ICU Group (7) wherein they examined 4,244 critical patients with COVID-19, the presence of a history of diabetes mellitus led to a 1.51 times increase in the 90-day mortality. By contrast, in the present study, the presence of comorbidities did not affect mortality in critical

Table 2. Risk factors for mortality in critical patients with COVID-19 in the intensive care unit

Characteristic	Mean (SD)	OR	95% CI OR	p-value
Age (year)	68.5 (15.1)	1.03	1.008-1.055	0.009
Male (%)	232 (52.1)	1.13	0.570-2.230	0.7
KDIGO score (%)				
1	83 (18.7%)	5.23	2.490-10.97	<0.001
2	79 (17.8%)	7.07	2.900-17.24	<0.001
3	94 (21.1%)	33.98	8.860-130.3	<0.001
APACHE-II score	16.6 (7.23)	1.023	0.960-1.090	0.49
SOFA score	4.34 (2.59)	1.194	1.007-1.416	0.041
Laboratory				
White blood cells ($\times 10^3/\mu\text{L}$)	11.33 (6.42)	0.995	0.918-1.079	0.91
Neutrophil ($\times 10^3/\mu\text{L}$)	9.51 (5.19)	0.950	0.831-1.087	0.45
Lymphocyte ($\times 10^3/\mu\text{L}$)	0.98 (0.57)	1.284	0.593-2.780	0.52
Neutrophil/lymphocyte ratio	12.8 (11.36)	1.069	1.006-1.137	0.031
Hemoglobin (g/dL)	12.8 (2.05)	0.694	0.373-1.292	0.25
Hematocrit (%)	40.61 (6.12)	1.062	0.867-1.300	0.56
Platelet ($\times 10^3/\mu\text{L}$)	242.5 (100.3)	0.998	0.995-1.001	0.27
Prothrombin time (s)	13.83 (2.76)	1.052	0.925-1.198	0.44
D-dimer (ng/mL)	2019.2 (4887.1)	1.000	1.000-1.001	0.007
pH	7.36 (0.1)	5.381	0.052-554.0	0.47
HCO ₃ (mmol/L)	21.78 (4.34)	0.888	0.802-0.983	0.02
Lactate (mmol/L)	2.76 (2.35)	1.010	0.813-1.254	0.92
Lactate dehydrogenase (U/L)	514.9 (413.07)	1.004	1.002-1.006	<0.001
Creatine kinase (IU/L)	314.05 (841.9)	1.000	1.000-1.001	0.17
C-reactive protein (mg/L)	141.2 (89.01)	1.003	0.999-1.006	0.10
Blood urea nitrogen (mg/dL)	64.1 (48.2)	1.000	0.991-1.010	0.91
Creatinine (mg/dL)	1.55 (1.76)	0.499	0.368-0.676	<0.001
AST (U/L)	67.9 (214.3)	1.000	0.993-1.007	0.95
Direct bilirubin (mg/dL)	0.38 (0.36)	0.619	0.288-1.332	0.22
Procalcitonin (ng/mL)	3.19 (10.08)	1.026	0.978-1.077	0.29
Ferritin ($\mu\text{g/L}$)	854.8 (635.2)	1.000	1.000-1.001	0.79

KDIGO: Kidney Disease: Improving Global Outcomes, APACHE-II: Acute Physiology and Chronic Health Evaluation-II, SOFA: Sequential Organ Failure Assessment, HCO₃: bicarbonate, AST: aspartate aminotransferase, SD: standard deviation, OR: odds ratio, CI: Confidence interval

patients with COVID-19 hospitalized in ICU. We believe that this result can be attributed to the patient population of our study. The mean patient age in this study was considerably high [68.5 (15.1) years], and most patients (76%) reported at least one comorbid disease. We believe that these factors led to the result observed.

APACHE-II and SOFA are scoring systems that are frequently used during the follow-up examination of critically ill patients for assessing disease severity and mortality (22). Some studies have stated that these scoring systems can be used to determine the disease course in patients with COVID-19 and that high APACHE-II and SOFA scores are associated with poor prognosis and mortality (4,22,23). In the present study, it was observed that non-survivor patients showed higher APACHE-II and SOFA scores during their first admission to ICU. However, only a high SOFA score was determined as a risk factor for mortality in critical patients with COVID-19 in ICU.

One of the important organs that is affected besides the respiratory system in patients with COVID-19 is the kidneys. Although renal manifestations specific to COVID-19 have not clearly been defined, acute renal damage may reportedly develop in 0.5-29% of the patients with COVID-19 and the incidence of acute renal damage is higher in patients experiencing severe disease or death (11,23-25). In the present study, patients with a mortal course showed higher KDIGO stages on the first day of admission to ICU. In addition, it was found that with the increase in the KDIGO stage of the patient, there was an increase in the mortality risk. These results indicate that a high KDIGO stage at the time of the first admission to ICU is an independent risk factor for mortality. Urgent application of appropriate treatments to patients with high KDIGO stage at admission will contribute to a substantial reduction in mortality risk.

Characteristic laboratory findings observed in critical patients with COVID-19 are reportedly low lymphocyte, albumin, and PaO₂ levels and high WBC, neutrophil, LDH, CRP, urea, creatinine, PTZ, activated partial thromboplastin time, ferritin, and PCT levels (12,26). Linli et al. (27) evaluated 192 critical patients with COVID-19 and stated that abnormal CRP, WBC, AST, and pH values were associated with high mortality and that CRP values should be closely monitored in these patients. Cummings et al. (11), evaluating 257 critically ill patients, stated that high D-dimer levels were an independent risk factor for in-hospital mortality. In

the present study, high WBC, neutrophil, N/L ratio, PTZ, D-dimer, lactate, LDH, CK, CRP, urea, creatinine, AST, direct bilirubin, PCT, and ferritin levels as well as low lymphocyte, hemoglobin, hematocrit, platelet, pH, and HCO₃ levels were detected. High N/L ratio, D-dimer, LDH, and creatinine values and low HCO₃ value were identified as independent risk factors for mortality in critical patients with COVID-19. Careful monitoring of these values in critical patients with COVID-19 hospitalized in ICU may act as a caution sign for mortality.

The most important limitation of this study was that it is retrospective and single centered. Conducting studies on critical patients with COVID-19 hospitalized in ICU with multicenter and large patient series across the country or the world will provide more precise information. Another study limitation is that the parameters such as obesity and regional and ethnic differences mentioned in some studies were not included. There was no information about these features in the data we obtained.

Conclusion

As a result of present study, it has been determined that the demographic characteristics of critical COVID-19 patients hospitalized in the ICU, as well as the clinical situation at the first admission to the ICU and some laboratory values are independent risk factors for mortality. In particular, the relationship between high KDIGO stages and mortality at the first admission to the ICU was noteworthy. We believe that monitoring these factors during the follow-up period of critical patients with COVID-19 in ICU can help predict the clinical course of the disease and reduce mortality.

Ethics

Ethics Committee Approval: The study was approved by the Clinical Research Ethics Committee of University of Health Sciences Turkey, Gazi Yaşargil Training and Research of Hospital (decision no: 550, date: 11.09.2020).

Informed Consent: This retrospective cohort study was conducted in accordance with the 2008 Declaration of Helsinki criteria.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: O.U., C.K.K., M.E.E., M.S.G., M.A., Z.K., Ö.C., Concept: O.U., C.K.K., M.E.E., M.S.G., Z.K., Ö.C., Design: O.U., C.K.K., M.E.E., M.A., Data

Collection and Process: O.U., C.K.K., M.E.E., M.S.G., M.A., Z.K., Ö.C., Analysis or Interpretation: O.U., C.K.K., M.E.E., M.A., Ö.C., Literature Search: O.U., C.K.K., M.E.E., M.S.G., M.A., Z.K., Ö.C., Writing: O.U., C.K.K.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. WHO Coronavirus Disease (COVID-19) Dashboard. [cited 2020 Dec 12]. Available from URL: <https://covid19.who.int/>
2. WHO Clinical management of COVID-19 [cited Oct 25 2020]. Available from URL: <https://www.who.int/publications/i/item/clinical-management-of-covid-19>.
3. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054-62.
4. Luo M, Cao S, Wei L, Zhao X, Gao F, Li S, et al. Intubation, mortality, and risk factors in critically ill Covid-19 patients: A pilot study. *J Clin Anesth* 2020;67:110039.
5. Zhang J, Wang X, Jia X, Li J, Hu K, Chen G, et al. Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect* 2020;26:767-72.
6. Martins-Filho PR, Tavares CSS, Santos VS. Factors associated with mortality in patients with COVID-19. A quantitative evidence synthesis of clinical and laboratory data. *Eur J Intern Med* 2020;76:97-9.
7. COVID-ICU Group on behalf of the REVA Network and the COVID-ICU Investigators. Clinical characteristics and day-90 outcomes of 4244 critically ill adults with COVID-19: a prospective cohort study. *Intensive Care Med* 2021;47:60-73.
8. Du Y, Lv Y, Zha W, Zhou N, Hong X. Association of body mass index (BMI) with critical COVID-19 and in-hospital mortality: A dose-response meta-analysis. *Metabolism* 2021;117:154373.
9. Yu C, Lei Q, Li W, Wang X, Liu W, Fan X, et al. Clinical Characteristics, Associated Factors, and Predicting COVID-19 Mortality Risk: A Retrospective Study in Wuhan, China. *Am J Prev Med* 2020;59:168-75.
10. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ* 2020;368:m1091.
11. Cummings MJ, Baldwin MR, Abrams D, Jacobson SD, Meyer BJ, Balough EM, et al. Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet* 2020;395:1763-70.
12. Skevaki C, Fragkou PC, Cheng C, Xie M, Renz H. Laboratory characteristics of patients infected with the novel SARS-CoV-2 virus. *J Infect* 2020;81:205-12.
13. Mehraeen E, Karimi A, Barzegary A, Vahedi F, Afsahi AM, Dadras O, et al. Predictors of mortality in patients with COVID-19-a systematic review. *Eur J Integr Med* 2020;40:101226.
14. Parra-Bracamonte GM, Lopez-Villalobos N, Parra-Bracamonte FE. Clinical characteristics and risk factors for mortality of patients with COVID-19 in a large data set from Mexico. *Ann Epidemiol* 2020;52:93-8.e2.
15. Sanders JM, Monogue ML, Jodlowski TZ, Cutrell JB. Pharmacologic Treatments for Coronavirus Disease 2019 (COVID-19): A Review. *JAMA* 2020;323:1824-36.
16. T.R. Ministry of Health COVID-19 information page. The management of Severe Pneumonia, ARDS, Sepsis and Septic Shock [cited Dec 30 2020]. Available from: <https://covid19.saglik.gov.tr/>
17. Kellum JA, Lameire N, Aspelin P, Barsoum RS, Burdmann EA, Goldstein SL, et al. Kidney disease: improving global outcomes (KDIGO) acute kidney injury work group. KDIGO clinical practice guideline for acute kidney injury [Internet]. Nature Publishing Group. Vol. 2. p. 1-138; 2012. *Kidney International Supplements* [cited Dec 21 2020]. Available from URL: <https://experts.umn.edu/en/publications/kidney-disease-improving-global-outcomes-kdigo-acute-kidney-injur>.
18. Leung C. Risk factors for predicting mortality in elderly patients with COVID-19: A review of clinical data in China. *Mech Ageing Dev* 2020;188:111255.
19. Gallo Marin B, Aghagoli G, Lavine K, Yang L, Siff EJ, Chiang SS, et al. Predictors of COVID-19 severity: A literature review. *Rev Med Virol* 2021;31:1-10.
20. Albitar O, Ballouze R, Ooi JP, Sheikh Ghadzi SM. Risk factors for mortality among COVID-19 patients. *Diabetes Res Clin Pract* 2020;166:108293.
21. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y, et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol* 2020;146:110-8.
22. Zou X, Li S, Fang M, Hu M, Bian Y, Ling J, et al. Acute Physiology and Chronic Health Evaluation II Score as a Predictor of Hospital Mortality in Patients of Coronavirus Disease 2019. *Crit Care Med* 2020;48:e657-65.
23. Yao Q, Wang P, Wang X, Qie G, Meng M, Tong X, et al. A retrospective study of risk factors for severe acute respiratory syndrome coronavirus 2 infections in hospitalized adult patients. *Pol Arch Intern Med* 2020;130:390-9.
24. Kunutsor SK, Laukkanen JA. Renal complications in COVID-19: a systematic review and meta-analysis. *Ann Med* 2020;52:345-53.
25. Nimkar A, Naaraayan A, Hasan A, Pant S, Durdevic M, Suarez CN, et al. Incidence and Risk Factors for Acute Kidney Injury and Its Effect on Mortality in Patients Hospitalized From COVID-19. *Mayo Clin Proc Innov Qual Outcomes* 2020;4:687-95.
26. Lu L, Zhong W, Bian Z, Li Z, Zhang K, Liang B, et al. A comparison of mortality-related risk factors of COVID-19, SARS, and MERS: A systematic review and meta-analysis. *J Infect* 2020;81:e18-e25.
27. Linli Z, Chen Y, Tian G, Guo S, Fei Y. Identifying and quantifying robust risk factors for mortality in critically ill patients with COVID-19 using quantile regression. *Am J Emerg Med* 2021;45:345-51.