

The Effect of the Fracture Types on the Activity Daily Living and Mortality in Geriatric Patients

Yüksel Uğur Yaradılmış¹, Alparslan Kılıç¹, Ali Teoman Evren¹, Tolgahan Kuru², Mahmut Özdemir³, Murat Altay¹

¹University of Health Sciences Turkey, Keçiören Health Practice and Research Center, Department of Orthopaedics and Traumatology, Ankara, Turkey

²Çanakkale Onsekiz Mart University Faculty of Medicine, Department of Orthopaedics and Traumatology, Çanakkale, Turkey

³VM Medikal Park Pendik Hospital, Clinic of Orthopaedics and Traumatology, Ankara, Turkey

Abstract

Objective: Knowledge of the frequency of geriatric fracture types and independency life rates will provide insights on its prevention in the community. Our study aimed to determine the effect of fracture types on loss of activity daily living (ADL) and mortality with one-year follow-up.

Materials and Methods: This study included a total of 522 patients (>65 years of age) with extremity and axial skeletal fractures. Fractures were classified according to location into 17 types. Incidence of the fracture types, hospitalization and operation rates in geriatric fractures were defined. Barthel and Katz indexes were evaluated for Basic activity daily living scale (BADLs) at the final control. The relationship between fracture types and mortality and decrease of BADLs was determined.

Results: The mean age of the patients was 76.24 ± 7.6 (range: 65–96) and the female-to-male ratio was 7/3. Four fracture types which constitute 60% of the fractures were noted: hip fracture (29.5%), vertebra fracture (7%), distal radius fracture (16%), and proximal humerus (8%). Mortality was noted in 21% of the patients. Mortality ratio and BADLs were statistically different according to the fracture types. The hip, vertebra, and femur fractures had lower Barthel index compared to other fractures.

Conclusion: High functional dependency and mortality were found in the patients with geriatric fracture, especially in the hip, femur and vertebra fractures. The patients have upper extremity fractures can reach the highest value of BADLs at one year follow up.

Keywords: Geriatric fractures, Activity Daily living, Katz index, Barthel index, Mortality

Introduction

In the elderly population, an increase in bone porosity and a decrease in bone tissue proportion are expected (1). In addition, falling is a health problem with a multifactorial etiology, especially over the age of 65 (2). Fractures are more common in older ages, because of lower bone quality and higher frequency of falls. Over the age of 65, the incidence of fractures is four times higher (3). With a worldwide aging population, the importance of the prevention and management of osteoporotic fragility fractures has emphasized over time.

Hip fractures, distal radius fractures, proximal humerus fractures, and vertebral fractures are the most common four types of geriatric fractures (4–6). While most of the fractures are seen commonly in males, the incidence of fractures in older ages is two times higher in females than in males (7). Another prominent feature in elderly fractures is that while the upper extremity fractures are often treated conservatively, the lower extremity fractures, especially the hip fractures, require hospitalization and surgical intervention (8).

Geriatric fractures are commonly evaluated with health-related quality of life (QoL) instruments, such as Short Form 36 of the

Address for Correspondence: Yüksel Uğur Yaradılmış, University of Health Sciences Turkey, Keçiören Health Practice and Research Center, Department of Orthopaedics and Traumatology, Ankara, Turkey

Phone: +90 532 769 78 02 **E-mail:** ugur_yaradilmis@outlook.com **ORCID:** orcid.org/0000-0002-7606-5690

Received: 05.08.2021 **Accepted:** 28.12.2021

Cite this article as: Yaradılmış YU, Kılıç A, Evren AT, Kuru T, Özdemir M, Altay M. The Effect of the Fracture Types on the Activity Daily Living and Mortality in Geriatric Patients. Eur J Geriatr Gerontol

Medical Outcomes Study (SF-36), Short Form 12 of the Medical Outcomes Study (SF-12), and EuroQol (EQ-5D) SF-12 [9, 10]. An important deficiency in literature regarding geriatric fractures is the fact that the relationship between fracture type and Activity Daily living (ADL) and mortality in geriatric patients is yet not to be investigated. Another point is the inadequacy of literature regarding the epidemiological study of fractures. Geriatric fractures have been examined with its subtypes and hip fractures subtypes was shown to emphasize their high mortality and morbidity (7). Alarkawi et al. made attention to the adequate research on fractures besides hip and vertebra fractures (11). This elderly group experience many comorbidities and major problems, such as osteoporosis and sarcopenia, which are difficult to change (12, 13). Perhaps it is important that they can continue their independent life and these patients should be evaluated with Activity Daily living.

Knowledge of the frequency of geriatric fracture types and independency life rates will provide insights on its prevention in the community as well as aid the determination of effective treatment strategies.

Our study aimed to determine the epidemiological characteristics of geriatric fractures and the effect of fracture types on loss of daily life activities and mortality with one-year follow-up.

Materials and Methods

Patients

In this study, a total of 535 patients (over 65 years of age) who were diagnosed with fractures in Orthopedics and Traumatology Clinic between January and December 2019 were retrospectively analyzed. Multiple fractures (n = 8) and periprosthetic fractures (n = 5) were excluded. A total of 522 patients with extremity and axial skeletal fractures were included in this study (Figure 1). Fractures were classified into 17 types as to their locations




Figure 1. Flow chart of patients enrolled in the study

(Table 1). Incidence of the fracture types among geriatric patients was defined. Epidemiological data of these patients, including age, gender, and comorbidities.

Hospitalization and Operation

General and extremity examinations were performed in the emergency department. Hospitalization was applied to patients who had surgical indications according to the fracture site and who were vitally unstable or expected to have instabile fracture. Patients who did not require surgical intervention and were vitally stable were discharged with their specific recommendations. Surgical treatment was done after hospitalization in line with the anesthesia risk ratio and patient approvals. Mechanical and medical deep venous thrombosis prophylaxis was administered immediately after hospitalization. Hospitalization and operation ratio of the fractures were determined.

Mortality

Mortality evaluated in 522 included patients. The exitus patients' data were collected from the death reporting system record. Besides fracture types, other variables relationship between mortality was determined.

Table 1. Classification of cetabula ceta according to the fractures locations

Fracture types	Subtypes for some fracture
Hip fractures	Intertrochanteric, femoral neck, subtrochanteric fractures
Vertebra fractures	Lombar, thoracal and cervical fractures
Humerus proximal fractures	
Distal radius fractures	
Ankle fractures	All malleol fractures
Foot fractures	Metatars, phalanx and tarsal fractures
Hand fractures	Metakarp, phalanx and carpal fractures
Pelvis and cetabular fractures	
Tibial plateau fractures	
Patella fractures	
Tibial fractures	Tibial saft and distal tibia fractures
Femural fractures	Femural saft and distal femur fractures
Humerus fractures	Humeral saft and distal humerus fractures
Radius fractures	Radius saft and proximal radius fractures
Ulna fractures	Ulnar saft and proximal ulna fractures
Clavicula fractures	
Scapula fractures	

Activity Daily living (ADL)

Basic ADL scale (BADLs) was applied to the patients at the final control (12–24 months). Barthel and Katz indexes were evaluated for Basic ADLs (14, 15). The Katz index measures six items: bathing, dressing, toileting, transferring (bed to chair and back), maintaining continence and feeding. Each of items was evaluated with one point, with a maximum of 6 points being obtainable. Functional independent patient is indicated by a Katz index of 6, while a functional dependent patient is indicated by a Katz index < 6. validity and reliability of the Turkish version of Katz index evaluated in healthy volunteers after 65 years of age Arik et al. (16). The Barthel Index comprises 10 items (feeding, bathing, grooming, dressing, bowel control, bladder control, toilet use, transfers, mobility on level surfaces and stairs) with a maximum of 100 points being obtainable. Validity and reliability of the Turkish version of Barthel index was assessed by Kucukdeveci et al. (17). A total of 114 exitus patients, patients with Katz index <6 before fracture (n = 8), and missing patients (n = 84) were excluded for BADLs evaluation. BADLs was performed for 312 patients (Figure 1). The relationship between fracture types and decrease of BADLs was determined.

Statistics

Data obtained in the study were analyzed statistically using SPSS v.22 software, and at a confidence interval of 95%. Qualitative data were stated as frequency distribution and quantitative data were stated as mean, minimum and maximum values. The χ^2 test was applied

to categorical data and the Student's t-test to quantitative data. Normality of data distribution was tested with the Shapiro–Wilk test. Demographic values were evaluated with the Mann Whitney U-test. Mortality, Barthel and Katz index of the fractures were evaluated with the Kruskal Wallis. The relationship between fractures types and mortality, and Katz index evaluated with the Mann Whitney U-test and Chi-square test. A value of $p < 0.05$ was accepted as statistically significant.

Results

Mean age of the patients was 76.24 ± 7.6 (range: 65–96) and female-to-male ratio was 7/3. There were 53% lower extremity fractures, 35% upper extremity fractures, 7% vertebra fractures and 5% pelvis–acetabular fractures. The demographic data of the patients were presented in Table 2. While 37.5% of the patients were hospitalized, 32.5% of the patients were operated. Mortality ratio was 21%.

Four fracture types, which constitute 60% of the fractures were noted: hip fractures (29.5%), vertebra fractures (7%), distal radius fractures (16%) and proximal humerus fractures (8%). Frequency of the geriatric fractures are presented in Figure 2.

Table 2. Demographic data of the patients

	n=522	Frequency
Age	76.24±7.6 (65-96)	
65-75	247	45.5%
75-85	196	36%
>85	100	18.5%
Gender		
Female/male	384/159	
Follow-up	16±1 (12-24)	
Side		
Right/left	266/237	
Vertebra	36	
Subtype		
Upper extremity	190	35%
Lower extremity	290	53%
Vertebra	37	7%
Pelvis-acetabulum	25	5%
Season		
Summer	147	29%
Spring	157	27%
Winter	123	23%
Autumn	116	21%




Figure 2. Frequency of fractures types

The hospitalized patients with hip fractures were 72.5% and other fracture types hospitalized are given in Figure 3. Only 6% of the patients with upper extremity fractures underwent surgery. The mean follow-up period of the patients was 16 (range: 12–24) months and the mortality ratio was 21%. The mortality ratios in the fracture types respectively; hip %46 (n=72), pelvis-acetabular %34 (n=9), vertebra %20 (n=8), tibia plateau %34 (n=4), ankle %14 (n=6), proximal humerus %25 (n=9) and distal radius fractures %6 (n=4). There was a statistical difference between mortality and fracture types ($p < 0.05$). Mortality was affected by variables such as age, comorbidity, and operation (Table 4).

Table 3. Basic activity of daily life of fracture types

	Frequency (n=512)	Barthel (n=312)	Katz (n=312)
Total		95±11.4	5.5±1
P value		0.005*	0.004*
Hip fractures	29.5%	90.7±14.1	5±1.3
Vertebra fractures	7%	88.7±19.5	5.1±1.4
Distal radius fractures	16%	98±1.3	5.9±0.3
Proximal humerus fractures	8%	93±11.7	5.3±1
Pelvis-cetabular fractures	5%	98±4.4	5.8±0.4
Tibial plateau fractures	3%	94±8.9	5.4±0.9
Patella fractures	1%	100	6
Ankle fractures	8.5%	100	6
Hand fractures	6.3%	98.5	5.8
Foot fractures	7.9%	100	6
Tibial fractures	1%	90	5
Femural fractures	1%	90	5
Humerus fractures	2%	100	6
Radius fractures	1%	100	6
Ulna fractures	1%	100	6
Clavicula fractures	1%	100	6
Scapula fractures	1%	100	6

*: p<0.05 and there is a significant statistical difference




Figure 3. Hospitalization and operation of the fracture types
 While 75% of the patients had a Katz index of 6 point, 25% of the patients were not completely functionally independent. Regarding the Barthel index values, 70% of the patients had 100 points. There was a statistical difference between BADLs and

fracture types (Table 3). Barthel index was lower in the patients with hip, vertebra, and femoral fractures than other fractures. Katz index was affected by variables such as age, comorbidity, and operation (Table 5).

Table 4. Relationship between demographic values and mortality

	Exitus	Live	p
	114	408	
Age	80.7±7.6	74.7±6.9	≤0.001*
Sex (F/M)	41/16	112/50	0.820
Comorbidity number	2.03±1	1.64±1.1	0.002*

*: p<0.05 and there is a significant statistical difference

Table 5. Relationship between demographic values and basic activity of daily life

	Katz index<6	Katz index=6	p
	74	238	
Age	78.5±7	73.5±6.5	≤0.001*
Sex (F/M)	25/12	82/37	0.878
Comorbidity number	1.48±1	2.2±1.2	≤0.001*

*: p<0.05 and there is a significant statistical difference

Discussion

Aging is associated with a higher fracture risk and diminished capacity of bone to heal. Bone fractures are common in the elderly, with residual lifetime fracture risk in a person aged 60 years reported to be %29 in males and 56% in females (18). Regeneration of bone defects often presents significant challenges, particularly in these patients with decreased tissue regeneration capacity (19). Many cases of delayed union and non-union are idiopathic in nature, several reports have suggested that these complications are more common in the elderly (20). Geriatric fractures can lead to work absence, decreased productivity, disability, impaired quality of life, health loss, and high health-care costs and are a major burden to individuals, families, societies, and health-care systems (21). Our study aimed to determine the frequency of geriatric fracture types and effect of fracture types on loss of activity daily living (ADL) and mortality with one-year follow-up .

In studies regarding aged-related fractures, hip fractures, vertebral fractures, and distal radius fractures are common fractures. The incidence of fractures is affected by geographic, ethnic, and socio-economic factors (22). While hip fractures are 11 times more incident in European countries, vertebral fractures were 3 times more incident in Scandinavian countries (23). Geriatric cases were most commonly located at the femur (43.4%), followed by radius (11.8%), humerus (10.6%),

and lumbar vertebra (3%). Fractures in geriatric cases are most commonly seen in women (61%), among those treated surgically (67%), and during winter (32.9%). In incident studies, diagnostic determination of vertebral fractures is difficult, since they are ignored over time and since no further examinations are performed (8). In our study, the four most common fractures were respectively: hip fractures (%29.5), distal radius fractures (%16), proksimal humerus fractures (%8) and vertebra fractures (%7). In addition, it was seen 2 times more frequently in women. Geriatric fractures are more common in summer and spring, but less common in winter and autumn in our study.

Aged-related fractures have never been evaluated with BADL. Katz and Barthel indexes are often used for BADL. Barthel Index is one of the most widely-used tools for assessment of functional independence (14). Kucukdeveci et al. in patients with stroke and spinal cord injury (17). Katz index is a shorter assessment and may be easier for patients (15). Arik et al. determined after 65 years of age the Katz index as 4.7 ± 1.6 and the Barthel Index as 86 ± 25 (16). In our study were found Katz index as 5.5 ± 1 and Barthel Index as 95 ± 11.4 for geriatric fractures. In line with these values, it is important to note that the rate of complete independence was low in the patients >75 years old. In our study, mild dependence (range: 62–90) was observed in vertebral, hip, and femur fractures according to the Barthel index, while complete independence (range: 91–99) was observed in other fractures. In a prospective evaluation of hip fractures, low ADL values was observed and the mortality rate of these patients was determined (24). In another study, ADL and Charlson comorbidity scores were used during follow-up to determine re-hospitalization after hip surgery (25).

Fractures are more mortal among the elderly than in normal population (11). Hip and vertebral fractures cause more morbidity and mortality than other fractures (26, 27). There is limited information about the effect of several fractures on mortality, except for vertebra and hip fractures. Alarkawi et al., in their study, mentioned the significant effect of proximal, non-hip, non-vertebra fractures, and subsequent fractures on mortality, excluding hip fractures (11). Although mortality is most common in lower extremity fractures; resulted in mortality in 46% of patients with hip fractures and 34% of patients with pelvis fractures in our study. On the other hand, 20% mortality was observed in vertebral fractures and it was not higher than expected. Apart from the type of fracture, variables such as advanced age and comorbidity, also increase mortality. Hip fractures have been frequently investigated in the literature because they are both common and mortal in elderly. Mortality has been found in many studies to correlate with age and comorbid conditions such as chronic liver, kidney, or cardiovascular diseases and pneumonia and dementia in hip fractures (28, 29). The other factors associated with mortality were cognitive impairment, delirium, living with caregiver, smoking and poor function before fracture (29, 30). When

evaluating mortality for elderly patients, it is necessary to consider a lot of risk factors.

Study Limitations

This study has several limitations, which include collection of data from a single center and its retrospective design. This study is innovative due to its inclusion of all fracture types and the questioning ADLs; however, the addition of multi-center studies with more patients in the future will increase the reliability and generalizability of the data. Another limitation of this study is the lack of data on status of patients in terms of osteoporosis.

Conclusion

Hip fractures in the geriatric forties constitute a major burden in Turkish population. High functional dependency and mortality are expected due to patients with geriatric fractures, especially the hip, femur, and vertebra fractures. The patients have upper extremity fractures can reach a high rate of full BADLs at one year follow up.

Ethics

Ethics Committee Approval: The study was approved by the Review Board and Commission of Keçiören Health Practice and Research Hospital in Turkey (approval date: 04.03.2021 number: 73-929).

Informed Consent: Written informed consent was obtained from participants.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Schuit SCE, van der Klift M, Weel AEAM, de Laet CEDH, Burger H, Seman E, et al. Fracture incidence and association with bone mineral density in elderly men and women: The Rotterdam Study. *Bone* 2004; 34:195-202.
- Yu WY, Hwang HF, Hu MH, Chen CY, Lin MR. Effects of fall injury type and discharge placement on mortality, hospitalization, falls, and ADL changes among older people in Taiwan. *Accident Analysis & Prevention*, 2013;50, 887-894.
- Gosch M, Kammerlander C, Neuerburg C. Osteoporose – Epidemiologie und Versorgungsqualität [Osteoporosis-epidemiology and quality of care]. *Z Gerontol Geriatr*. 2019;52(5):408-413.
- Gül D, Akpancar S. Fractures in Geriatric Cases. *Geriatrisk Bilimler Dergisi*, 2019;2(1), 14-19.
- De Laet CE, Pols HA. Fractures in the elderly: epidemiology and demography. *Baillieres Best Pract Res Clin Endocrinol Metab*. 2000;14(2):171-9.
- Baidwan NK, Naranje SM. Epidemiology and recent trends of geriatric fractures presenting to the emergency department for United States population from year 2004-2014. *Public Health*. 2017;142:64-69.
- Alexiou KI, Roushias A, Varitimidis SE, Malizos KN. Quality of life and psychological consequences in elderly patients after a hip fracture: a review. *Clin Interv Aging*. 2018;24;13:143-150.

8. Clynes MA, Harvey NC, Curtis EM, Fuggle NR, Dennison EM, Cooper C. The epidemiology of osteoporosis. *Br Med Bull.* 2020;15;133(1):105–117.
9. Hallberg I, Rosenqvist AM, Kartous L, Löfman O, Wahlström O, Toss G. Health-related quality of life after osteoporotic fractures. *Osteoporos Int.* 2004;15(10):834–41.
10. Lips P, van Schoor NM. Quality of life in patients with osteoporosis. *Osteoporos Int.* 2005 16(5):447–55.
11. Alarkawi D, Bliuc D, Tran T, Ahmed LA, Emaus N, Bjørnerem A, Jørgensen L, Christoffersen T, Eisman JA, Center JR. Impact of osteoporotic fracture type and subsequent fracture on mortality: the Tromsø Study. *Osteoporos Int.* 2020;31(1):119–130.
12. Liu J, Curtis EM, Cooper C, Harvey NC. State of the art in osteoporosis risk assessment and treatment. *J Endocrinol Invest.* 2019;42(10):1149–1164.
13. Messina C, Maffi G, Vitale JA, Olivieri FM, Guglielmi G, Sconfienza LM. Diagnostic imaging of osteoporosis and sarcopenia: a narrative review. *Quant Imaging Med Surg.* 2018 ;8(1):86–99.
14. Mahoney FI, Barthel DW. Functional evaluation: the Barthel index. *Maryland State Medical Journal.* 1965;14, 61–65.
15. Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of illness in the aged. The index of ADL: a standardized measure of biological and psychosocial function. *JAMA.* 1963;185, 914–919.
16. Arik G, Varan HD, Yavuz BB, Karabulut E, Kara O, Kilic MK et.al. Validation of Katz index of independence in activities of daily living in Turkish older adults. *Arch Gerontol Geriatr.* 2015;61(3):344–50.
17. Kucukdeveci AA, Yavuzer G, Tennant A, Suldur N, Sonel B, Arasil T. Adaptation of the modified Barthel index for use in physical medicine and rehabilitation in Turkey. *Scandinavian Journal of Rehabilitation Medicine.* 2000;32, 87– 92.
18. Jones G, Nguyen T, Sambrook PN, Kelly PJ, Gilbert C, Eisman JA. Symptomatic fracture incidence in elderly men and women: the Dubbo Osteoporosis Epidemiology Study (DOES). *Osteoporosis International.* 1994;4(5), 277–282.
19. Gruber R, Koch H, Doll BA, Tegtmeier F, Einhorn TA, Hollinger JO. Fracture healing in the elderly patient. *Exp Gerontol.* 2006 Nov;41(11):1080–93.
20. Foulke BA, Kendal AR, Murray DW, Pandit H. Fracture healing in the elderly: A review. *Maturitas.* 92, 2016;49–55.
21. Borgström F, Karlsson L, Ortsäter G, et. al. International Osteoporosis Foundation. Fragility fractures in Europe: burden, management and opportunities. *Arch Osteoporos.* 2020 Apr 19;15(1):59.
22. Fuggle NR, Curtis EM, Ward KA, Harvey NC, Dennison EM, Cooper C. Fracture prediction, imaging and screening in osteoporosis. *Nat Rev Endocrinol.* 2019;15:535–547.
23. Kanis JA, Oden A, McCloskey EV, Johansson H, Wahl DA, Cooper CA. systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporos Int.* 2012;23:2239–56.
24. Aarden JJ, van der Esch M, Engelbert RHH, van der Schaaf M, de Rooij SE, Buurman BM. Hip Fractures in Older Patients: Trajectories of Disability after Surgery. *J Nutr Health Aging.* 2017;21(7):837–842.
25. da Casa C, Pablos-Hernández C, González-Ramírez A, Julián-Enriquez JM, Blanco JF. Geriatric scores can predict long-term survival rate after hip fracture surgery. *BMC Geriatr.* 2019;19(1):205.
26. Dennison E, Cole Z, Cooper C. Diagnosis and epidemiology of osteoporosis. *Curr Opin Rheumatol* 2005;17:456–461.
27. Melton LJ 3rd, Achenbach SJ, Atkinson EJ, Thorneau TM, Amin S. Long-term mortality following fractures at different skeletal sites: a population-based cohort study. *Osteoporos Int* 2013;24(5):1689– 1696 12.
28. Yong EL, Ganesan G, Kramer MS, Howe TS, Koh JSB, Thu WP, et. al. Risk Factors and Trends Associated With Mortality Among Adults With Hip Fracture in Singapore. *JAMA Netw Open.* 2020 Feb 5;3(2):e1919706.
29. Chang W, Lv H, Feng C, Yuwen P, Wei N, Chen W, et. al. Preventable risk factors of mortality after hip fracture surgery: Systematic review and meta-analysis. *Int J Surg.* 2018 Apr;52:320–328.
30. Liu Y, Wang Z, Xiao W. Risk factors for acetabula in elderly patients with hip fractures: a meta-analysis of 18 studies. *Aging Clin Exp Res.* 2018;30(4):323–330.