

The Effect of Different Anesthesia Techniques on Strabismus Surgery: A Prospective Observational Study

Farklı Anestezi Tekniklerinin Şaşılık Cerrahisi Üzerine Etkisi: Prospektif Gözlemsel Çalışma

● Bülent Barış Güven¹, ● Mehmet Mutlu², ● Seray Türkmen²

¹University of Health Sciences Turkey Hamidiye Faculty of Medicine; İstanbul Sultan 2. Abdülhamid Han Training and Research Hospital, Clinic of Anesthesiology and Reanimation, İstanbul, Turkey

²University of Health Sciences Turkey, Prof. Dr. Cemil Taşcıoğlu City Hospital, Clinic of Anesthesiology and Reanimation, İstanbul, Turkey

ABSTRACT

Background: Strabismus correction surgery is the most common eye operation in children. This present study aims to evaluate the effects of orotracheal intubation using a neuromuscular blocking agent (NMA) in strabismus surgery anesthesia and laryngeal mask airway (LMA) application using total intravenous anesthesia without using NMA on the deviation angles measured following strabismus surgery.

Materials and Methods: ASA I-II group patients with concomitant horizontal strabismus, aged under 14, scheduled for strabismus surgery under general anesthesia were prospectively examined for the study. The study did not include those with restrictive type strabismus, vertical deviation, intraoperative vertical muscle group treatments, anisometropia, and corrected visual acuity less than 0.8. According to the order of arrival, either OTE was conducted with NMA, or LMA (LMA-group) was used without NMA for general anesthesia in patients.

Results: The study included 49 patients, divided into LMA (n=24) and OTE (n=25). The patients' mean age was 8.04±3.63 years. The distribution was similar between groups, with 63% (n=31) having esotropia and 37% (n=18) having exotropia. Angles of deviation were measured to be above 10 prism diopter in 1 patient in the LMA group and two patients in the OTE group at the end of postoperative 1st month. There was no statistically significant difference between the groups regarding decreased postoperative deviation angles and preoperative-postoperative deviation angle measurements.

Conclusion: The tone changes in the eye muscles using NMAs in strabismus surgery anesthesia caused no changes in the postoperative deviation angles or the surgical result.

Keywords: Anesthesia, surgery, strabismus

ÖZ

Amaç: Strabismus düzeltme operasyonları çocuklarda en sık yapılan göz operasyonlarından. Bu çalışmada amacımız şaşılık cerrahisi anesteziinde nöromusküler blokör ajan (NMA) kullanılarak yapılan orotrakeal entübasyon (OTE) uygulaması ile NMA kullanılmadan total intravenöz anestezi kullanılarak yapılan laringeal maske airway (LMA) uygulamasının şaşılık cerrahisi sonrası ölçülen kayma açılarına etkisini değerlendirmektir.

Gereç ve Yöntemler: Genel anestezi altında şaşılık cerrahisi planlanan concomitant horizontal yönde şaşılığı bulunan, ASA I-II grubu, 14 yaş altındaki hastalar çalışma için prospektif olarak değerlendirilmeye alındı. Restriktif tip strabismus olanlar, vertikal yönde kayması olanlar veya intraoperatif vertikal kas gruplarına da müdahale edilenler, anizometropisi bulunanlar ve düzeltilmiş görme keskinliği 0,8'in altında olanlar çalışma dışı bırakıldı. Hastalara genel anestezi için geliş sırasına göre ya NMA uygulanarak OTE yapıldı ya da NMA uygulanmadan LMA (LMA-grubu) kullanıldı.

Bulgular: LMA-grubu (n=24) ve OTE-grubu (n=25) olmak üzere toplam 49 hasta çalışmaya alındı. Hastaların yaş ortalaması 8,04±3,63 yıl idi. Hastalıkların %63'ünde (n=31) esotropia varken, %37'sinde (n=18) eksotropia vardı ve gruplar arasındaki dağılımı benzerdi. Postoperatif birinci ayın sonundaki kayma açısı LMA-grubunda 1 hastada, OTE-grubunda ise 2 hastada 10 prizma diyoptri üstünde



Address for Correspondence: Bülent Barış Güven, University of Health Sciences Turkey Hamidiye Faculty of Medicine; İstanbul Sultan 2. Abdülhamid Han Training and Research Hospital, Clinic of Anesthesiology and Reanimation, İstanbul, Turkey

Phone: +90 533 638 34 45 E-mail: barguv@gmail.com **ORCID ID:** orcid.org/0000-0002-3628-7408

Received: 04.02.2022 **Accepted:** 11.02.2022

öçüldü. Preoperatif ve postoperatif kayma açıları ile postoperatif kayma açısı ölçümlerindeki azalma açısından gruplar arasında istatistiksel olarak anlamlı bir fark gözlenmedi.

Sonuç: Şaşılık cerrahisi anesteziinde NMA'ların kullanılması ile göz kaslarında oluşan tonus değışikliğı postoperatif kayma açılarını etkilememiş ve cerrahi sonucu değıştirmemiştir.

Anahtar Kelimeler: Anestezi, cerrahi, şaşılık

Introduction

Children with strabismus account for 3-5% of all children, with correction surgeries the most prevalent among eye operations (1). Most pediatric ophthalmic procedures are done under general anesthesia (2). Under general anesthesia, airway patency is maintained through endotracheal intubation or the use of a laryngeal mask airway (LMA). LMA is a more practical and less invasive procedure in the current anesthetic practices preferred over endotracheal intubation, especially in short-term surgeries and patients with difficult airways in emergencies. The LMA's design maintains epiglottis patency, allowing for easier ventilation. Air leakage is avoided using the inflatable cuff. The main benefit is that there is no requirement for muscle relaxation, and there are fewer postoperative respiratory problems than orotracheal intubation (OTE) (3,4).

In strabismus surgery, the extraocular muscles are millimetrically shortened or lengthened to treat the impairment. Different anesthetic procedures may produce varying degrees of muscle tone loss during the application of anesthesia, resulting in inaccurate surgical correction results. The systemic effects of various anesthetic procedures in strabismus surgery have been mainly studied in patients in the literature so far (1,2,4). However, there is limited research on the outcomes of anesthetic procedures in strabismus surgery (5,6). The present study aims to evaluate the effects of OTE using a neuromuscular blocking agent (NMA) in general anesthesia and LMA application using total intravenous anesthesia (TIVA) without using NMA on the deviation angles measured following strabismus surgery.

Material and Methods

This study was designed as a single-center prospective observational study. Ethics committee approval was obtained at the Clinical Research Ethics Committee of Okmeydanı Training and Research Hospital meeting, dated 07/08/2018, with the decision numbered 48670771-514.10.

The study was carried out with patients who had strabismus surgery under general anesthesia in Okmeydanı

Training and Research Hospital Kasımpaşa Additional Service Building between September 2018 and September 2019. SA I-II group patients aged under 14, who were scheduled for surgery under general anesthesia and had deviation angles of 15 prism diopter (Δ D) and above due to strabismus, as well as non-paralytic horizontal strabismus, were assessed in the study. In cooperative patients, the visual acuity was assessed using the Snellen chart and form optotypes, while in non-cooperative patients, it was inferentially determined according to the fixation quality. After securing cycloplegia, a refraction was examined using a retinoscope and auto refractometer. Anterior and posterior segments were evaluated using slit-lamp and ophthalmoscope. The fixation and object tracking features were examined in non-cooperative cases, while the Worth 4-point test was employed to identify the eye to be operated on. Globe movements were assessed in nine cardinal gaze positions, and instances were classified as esotropia or exotropia. In cooperative cases, deviation angles were evaluated using the prism cover test, while in non-cooperative cases, they were assessed using the Krinsky or Hirschberg test. Those with lens, cornea, retina, or optic nerve pathologies that may cause visual impairment, those with restrictive type strabismus, those with vertical shifts or those who had intraoperative vertical muscle interventions, those with mental retardation, those with over 2 Δ D refractive deficiency (anisometropia) between both eyes or in one eye, those with corrected visual acuity less than 0.8, those who were re-operated for strabismus and those who refuse to participate in the study were excluded.

The patient's parents were informed about the study, and their written consent was obtained. None of the patients were given pre-medication. General anesthesia was induced with intravenous propofol (2-3 mg/kg) and maintained with sevoflurane. Consecutively, either OTE (OTE-group) was applied to patients using rocuronium under maintenance anesthesia with sevoflurane (1-2 minimal alveolar concentration), and remifentanyl (0.1-0.4 μ g/kg/min), or LMA (LMA-group) was applied under TIVA using propofol (6-9 mg/kg/hour) and remifentanyl (0.1-0.4 μ g/kg/min). Horizontal (lateral and medial rectus) muscle resections with or without recession were performed in all the

cases. In all cases, it is intended for post-surgery deviation angles to be ten ΔD and below. All these procedures were performed by the same specialist using the same methods. Side effects were recorded, including postoperative nausea and vomiting (PONV) and oculocardiac reflex (OCR). OCR was defined as a reduction of at least 15% in baseline heart rate or dysrhythmia. Those with a heart rate of less than <65 beats/min (bradycardia) were given 0.02 mg/kg atropine intravenously. Those experiencing complications of nausea and vomiting within the 2nd post-operative period were identified as PONV, and they were treated with intravenous 0.1 mg/kg ondansetron. Patients were routinely called for an examination on the post-operative 1st day, 1st week, and the 1st month. A physician who was not involved in the study kept track of the patients' postoperative complications and deviation angles at the end of the first month. Demographic data, anesthetic duration, complications, and preoperative and postoperative deviation angles of patients in both groups were compared statistically.

Statistical Analysis

Number Cruncher Statistical System software was used for statistical analyses. Descriptive statistical methods (mean, standard deviation, median, frequency, percentage, minimum, maximum) were used to evaluate the data. The conformity of quantitative data with the normal distribution was tested with the Kolmogorov-Smirnov and Shapiro-Wilk tests and visual examinations. The student t-test was used for the binary intergroup comparisons of the quantitative data with a normal distribution. In contrast, the Mann-Whitney U test was used for the binary inter-group comparisons of the quantitative data without normal distribution. The Pearson chi-square test was used to compare the qualitative data. The Wilcoxon Signed Ranks test was used for non-normally

distributed parameters intergroup comparisons. The significance was evaluated as minimum $p < 0.05$.

Results

Eighty-nine pediatric patients diagnosed with strabismus were assessed during the study. Out of 36 patients who did not meet the study's criteria, 12 had anisometropia, 9 had inadequate visual acuity, 12 had vertical strabismus, and 3 had restrictive strabismus. Fifty-three patients were consecutively divided into the LMA group (n=26) or OTE (n=27). One patient in the LMA group and two patients in the OTE group were eliminated from the study because at least one of the intraoperative vertical muscle groups was intervened. One patient in the LMA group failed to show up for the postoperative follow-up. As a result, 24 patients in the LMA group and 25 patients in the OTE group had their data analyzed (Figure 1).

Of the cases included in the study, 33% (n=16) were female and 67% (n=33) were male. The participants' ages ranged from 2 to 14, with a mean of 8.04 ± 3.63 years. Body mass index (BMI) measurements ranged from 20.6 to 25.8 kg/m² with a mean value 22.20 ± 1.25 kg/m². There was no statistical difference between the LMA and OTE groups in terms of gender (p=0.232), age (p=0.469) and BMI levels (p=0.109) (Table 1).

While esotropia was found in 63% (n=31) of the cases, exotropia was found in 37% (n=18), and there was no statistical difference in the distribution between the two groups (p=0.769). In the study, the mean duration of anesthesia was 49.5 ± 9.9 minutes, with 7.6 ± 12.1 minutes in the LMA group and 51.1 ± 8.2 minutes in the OTE group. Bradycardia OCR was noted in 16 patients throughout the intraoperative phase, and 10 cases were treated with intravenous atropine, whereas bradycardia resolved

Table 1. Demographic and preoperative clinical data of the patients

		Total (n=49)	LMA-group (n=24)	OTE-group (n=25)	p
Age (year)	Mean ± SD	8.04±3.63	7.78±3.74	8.30±3.55	0.469
Gender; n (%)	Female	16 (33)	10 (42)	6 (24)	0.232
	Male	33 (67)	14 (58)	19 (76)	
BMI (kg/m ²)	Mean ± SD	22.20±1.25	22.00±1.05	22.40±1.40	0.109
Anesthesia time (min)	Mean ± SD	49.5±9.9	47.6±12.1	51.1±8.2	0.456
PONV (n)		20	6	14	0.042*
OCR (n)		16	9	7	0.763
Type of strabismus; n (%)	Esotropia	31 (63)	16 (67)	15 (60)	0.769
	Exotropia	18 (37)	8 (33)	10 (40)	

LMA: Laryngeal mask airway, OTE: Orotracheal intubation, BMI: Body mass index, PONV: Postoperative nausea or vomiting, OCR: Oculocardiac reflex, SD: Standard deviation, min: Minute, *p<0.05

spontaneously in 6 cases. Both groups had statistically similar distributions ($p=0.763$). PONV was found in 6 patients in the LMA group and 14 cases in the OTE group, and the difference was statistically significant ($p=0.042$).

The preoperative degrees of deviation of the cases in the LMA group ranged between 12-50 Δ D, with a mean of 28.48 Δ D. At the end of the first postoperative month, the degree of deviation ranged between 0-11 Δ D, with a mean

of 1,2 Δ D. The preoperative and postoperative degrees of deviation in the LMA group were statistically different ($p=0.001$) (Table 2).

The preoperative degrees of deviation of the cases in the OTE group ranged between 20-55 Δ D, with a mean of 31.50 Δ D. At the end of the first postoperative month, the degree of deviation ranged between 0-13 Δ D, with a mean of 1,5 Δ D. The OTE group's preoperative and postoperative degrees




Figure 1. CONSORT flow diagram of the study participants
OTE: Orotracheal intubation, LMA: Laryngeal mask airway

Table 2. Evaluation of deviation angles in the postoperative 1st month

		LMA-group (n=24)	OTE-group (n=25)	p
The deviation angle before surgery (Δ D)	Mean \pm SD	28.48 \pm 8.54	31.50 \pm 8.88	0.135
	Median (min/max)	30 (12-50)	30 (20-55)	
The deviation angle after surgery (Δ D)	Mean \pm SD	1.20 \pm 1.98	1.50 \pm 2.53	0.784
	Median (min/max)	0 (0-11)	0 (0-13)	
	p	0.001**	0.001**	
The corrected deviation angle after surgery (Δ D)	Mean \pm SD	-27.28 \pm 7.70	-30.00 \pm 7.95	0.125
	Median (min/max)	-27 (-39/-12)	-30 (-42/-20)	

LMA: Laryngeal mask airway, OTE: Orotracheal intubation, Δ D: Prism diopter, SD: Standard deviation, ** $p<0.01$




Figure 2. Distribution of deviation angles by groups one month after surgery

LMA: Laryngeal mask airway, OTE: Orotracheal intubation, Preop: Preoperative period, Postop: Postoperative period

of deviation were statistically different ($p=0.001$) (Figure 2). One patient in the LMA group and two patients in the OTE group had a degree of deviation than $10 \Delta D$ at the end of the postoperative first month. There was no statistically significant difference between the groups regarding preoperative and postoperative deviation angles, the decrease in postoperative deviation angle measurements (Table 2).

Discussion

In ophthalmic surgeries, as in many other surgical procedures, anesthetists and surgeons use the technique and approach used to treat their patients. There is currently no consensus, particularly in strabismus surgery. Some surgeons prefer NMA to perform strabismus surgery, mainly while doing the Forced duction test. They are not affected by muscle tone fluctuations caused by changing the anesthetic type and depth (7). However, some clinicians do not opt for NMA. Migliorini et al. (5) studied the effects of the inhalation agent sevoflurane and desflurane in 42 patients undergoing strabismus surgery without NMA. They found that desflurane increased ocular deviation substantially more than sevoflurane. No study in the literature investigated the effect of employing NMA on the outcomes of strabismus surgery. According to the present study, the use of NMAs in strabismus surgery anesthetic did not affect the operative outcome, which is the first in this area.

Strabismus surgery rectifies the shift in the visual axes, give single binocular vision, and achieve a fair physical appearance. Several elements influence the success of the surgery. Patients with the same degree of strabismus may respond differently to the same amount of surgical correction. This indicated that the degree of deviation or surgery amount (operated muscle) is not the sole factor

affecting strabismus surgery. As a result, the inability to predict the outcome of strabismus surgeries becomes a significant issue. The globally approved scales given by the American Academy of Ophthalmology are frequently used to guide the amount of muscle intervention to be made in strabismus surgeries (8). Because the values in the scales are often organized according to the preoperative deviation angles, full correction may not be possible after surgery. The influence of alterations in ocular muscle tone, which occurred with two distinct anesthetic types, on postoperative deviation angles was explored in this study, and no difference in postoperative deviation angle remission was found between the NMA (OTE) and the NMA-free (LMA) groups.

Many factors influence the degree of deviation in the postoperative period, including age, refractive problems, decreased visual acuity, type and duration of strabismus, preoperative deviation angles, surgeon experience, and how the suture is passed through the sclera (9,10,11). Although most strabismus surgeries are operated under general anesthesia, eligible and compatible adult patients can be operated on with local or topical anesthetic (2,12). Because all the patients in our study were children (aged between 2 and 14), they were all operated on under general anesthesia by the same surgeon using the same procedure. Esotropia was found in 63% of the patients, while exotropia was found in 37%, with a similar distribution between the two groups. Patients with a corrected visual acuity of greater than 0.8 were also included in the study, excluding most factors that could alter the surgical outcome. While the rates of decrease in deviation angles were similar in both groups in the postoperative 1st-month follow-up, only three (3.4%) patients had a value over the predicted ($<10 \Delta D$) deviation angle. The three patients who did not have the expected outcome were over eight and had a preoperative deviation angle of more than $40 \Delta D$. Kampanartsanyakorn et al. (10) also presented the first surgical results of 304 patients with horizontal strabismus (61.5% esotropia, 38.5% exotropia), reporting that success rates were low in patients aged over six and in patients with preoperative deviation angles greater than $30 \Delta D$. As a result, we were unable to acquire the desired postoperative values in our patients with advanced age (>8 years) and sizeable preoperative deviation angles ($>40 \Delta D$) in our study.

LMA is regularly and safely used for pediatric anesthesia (13,14). LMA in pediatric anesthesia was considered an alternative to tracheal intubation in the early 2000s; however, with the advent of next-generation LMAs such as l-gel and Cobra perilyngeal airway, LMA has now become the preliminary choice in general anesthetic applications. Gulati et al. (15) compared the use of LMA and OTE in

strabismus surgery in 2004. They found that the LMA group had lower intraocular pressure increase and greater PONV due to LMA-induced gastric inflation. Nevertheless, PONV was shown to be lower in the LMA group in our study. The lack of a gastric drainage canal and inhalation medications utilized for anesthesia maintenance in first-generation classical LMAs could cause this paradox. In our study, we employed propofol (TIVA) to maintain anesthesia in the LMA group, and we used the new generation l-gel LMA with gastric drainage channel as the LMA. As a result, we observed statistically considerably less PONV than the OTE group, where anesthesia was maintained with an inhalation drug. Kranke et al. (16) also reported that when propofol was utilized for strabismus surgery anesthesia, there was less PONV than inhalation agents.

Some authors report that propofol has the same effect as metoclopramide, an antiemetic agent, at sub-hypnotic doses (17). The mechanism of this effect is not precise. It has been postulated to be an antagonist at the 5HT₃ receptor. Other reports suggest that the anti-emetic effect of propofol is due to modulation of the subcortical pathways (16,17). In many recent reports, propofol should be the first choice among anesthetic agents, especially since it prevents the development of PONV in strabismus surgery of pediatric patients (18).

OCR is identified with a reduction in heart rate of more than 15% or the onset of a new arrhythmia. During the traction and manipulation of the extraocular muscles in strabismus surgery, OCR can develop at rates ranging from 14 percent to 90 percent (19). OCR was detected at a rate of 33% in our investigation, with no differences between the groups. Aydın et al. (20) presented the results of 86 patients who underwent pediatric strabismus surgery and reported that 34% of OCR developed. The same report also stated that the group utilizing NMA developed statistically more negligible OCR. In our study, the NMA group (OTE) developed quantitatively more minor OCR, although there was no statistical significance. In this study, rocuronium was used as NMA. It is well known that rocuronium has vagolytic effects (21). We think that this effect reduced the development of OCR in our study. Additionally, since some studies reported that the OCR was statistically lower in patients receiving premedication, we did not administer premedication to any patient in our study to not impair the standardization of our research (20).

Study Limitations

The most significant limitation of the study is the small number of cases and one-month postoperative follow-up. The success of strabismus surgery was also assessed in terms of anatomical success in our research. Because the

patients were not followed up for an extended period, functional benefits such as binocularity and stereopsis were removed from the evaluation. Thus, randomized controlled studies with a larger sample size and more extended follow-up periods (>12 months) are required.

Conclusion

The tone changes in the eye muscles with the use of NMAs in strabismus surgery anesthesia caused no changes in the postoperative deviation angles or the surgical result.

Acknowledgments

We want to thank the eye clinic that operated and measurements on the cases in our study for their essential efforts.

Ethics

Ethics Committee Approval: All procedures performed in studies involving human participants were in accordance with the national ethical standards and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. The study was approved by the Ethics Committee of Okmeydanı Training and Research Hospital (approval no: 943, approval date: August 7, 2018).

Informed Consent: Informed consent was obtained from all individual participants included in the study.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: M.M., Concept: B.B.G., M.M., Design: B.B.G., M.M., Data Collection or Processing: M.M., S.T., Analysis or Interpretation: M.M., S.T., Literature Search: B.B.G., Writing: B.B.G.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Chua AW, Chua MJ, Leung H, Kam PC. Anaesthetic considerations for strabismus surgery in children and adults. *Anaesth Intensive Care*. 2020;48:277-288. [\[Crossref\]](#)
2. Lewis H, James I. Update on anaesthesia for paediatric ophthalmic surgery. *BJA Educ*. 2021;21:32-38. [\[Crossref\]](#)
3. Arican S, Pekcan S, Hacibeyoglu G, Yusifov M, Yuce S, Uzun ST. The place of ultrasonography in confirming the position of the laryngeal mask airway in pediatric patients: an observational study. *Braz J Anesthesiol*. 2021;71:523-529. [\[Crossref\]](#)
4. Luce V, Harkouk H, Brasher C, Michelet D, Hilly J, Maesani M, et al. Supraglottic airway devices vs tracheal intubation in children: a quantitative meta-analysis of respiratory complications. *Paediatr Anaesth*. 2014;24:1088-1098. [\[Crossref\]](#)

5. Migliorini R, Collini S, Malagola R, Servidio A, Cannata R, Arrico L. Anesthesia in the Surgery of Strabismus: Role of Anesthetic Agents in the Ocular Deviation and Surgical Outcome. *J Anesth Clin Res.* 2015;10:572. [\[Crossref\]](#)
6. Seijas O, Gómez de Liaño P, Merino P, Roberts CJ, Gómez de Liaño R. Topical anesthesia in strabismus surgery: a review of 101 cases. *J Pediatr Ophthalmol Strabismus.* 2009;46:218-222. [\[Crossref\]](#)
7. Dell R, Williams B. Anaesthesia for strabismus surgery: a regional survey. *Br J Anaesth.* 1999;82:761-763. [\[Crossref\]](#)
8. American Academy of Ophthalmology. Basic and Clinical Science Course Subcommittee. Basic and Clinical Science Course. Pediatric Ophthalmology and Strabismus: Section 6, 2016-2017. San Francisco, CA: American Academy of Ophthalmology; 2016:84-85. [\[Crossref\]](#)
9. Astudillo PP, Cotesta M, Schofield J, Kraft S, Mireskandari K. The Effect of Achieving Immediate Target Angle on Success of Strabismus Surgery in Children. *Am J Ophthalmol.* 2015;160:913-918. [\[Crossref\]](#)
10. Kampanartsanyakorn S, Surachatkumtonekul T, Dulayajinda D, Jumroendararamee M, Tongsaee S. The outcomes of horizontal strabismus surgery and influencing factors of the surgical success. *J Med Assoc Thai.* 2005;88(Suppl 9):S94-99. [\[Crossref\]](#)
11. Kekunnaya R, Mendonca T, Sachdeva V. Pattern strabismus and torsion needs special surgical attention. *Eye (Lond).* 2015;29:184-190. [\[Crossref\]](#)
12. Chua AW, Chua MJ, Leung H, Kam PC. Anaesthetic considerations for strabismus surgery in children and adults. *Anaesth Intensive Care.* 2020;48:277-288. [\[Crossref\]](#)
13. Tekin B, Hatipoğlu Z, Türktan M, Özcengiz D. Comparing the Laryngeal Mask Airway, Cobra Perilaryngeal Airway and Face Mask in Children Airway Management. *Turk J Anaesthesiol Reanim.* 2016;44:81-85. [\[Crossref\]](#)
14. Su MP, Hu PY, Lin JY, Yang ST, Cheng KI, Lin CH. Comparison of laryngeal mask airway and endotracheal tube in preterm neonates receiving general anesthesia for inguinal hernia surgery: a retrospective study. *BMC Anesthesiol.* 2021;21:195. [\[Crossref\]](#)
15. Gulati M, Mohta M, Ahuja S, Gupta VP. Comparison of laryngeal mask airway with tracheal tube for ophthalmic surgery in paediatric patients. *Anaesth Intensive Care.* 2004;32:383-389. [\[Crossref\]](#)
16. Kranke P, Apfel C, Papenfuss T, Metterlein T, Schuster F, Muellenbach R, et al. Nausea and vomiting after strabismus-surgery: a randomized comparison of isoflurane, enflurane, sevoflurane and propofol. *Open Clin Trials J.* 2009;1:1-6. [\[Crossref\]](#)
17. Kampo S, Afful AP, Mohammed S, Ntim M, Buunaaim ADB, Anabah TW. Sub-hypnotic dose of propofol as antiemetic prophylaxis attenuates intrathecal morphine-induced postoperative nausea and vomiting, and pruritus in parturient undergoing cesarean section - a randomized control trial. *BMC Anesthesiol.* 2019;19:177. [\[Crossref\]](#)
18. Kovac AL. Postoperative Nausea and Vomiting in Pediatric Patients. *Paediatr Drugs.* 2021;23:11-37. [\[Crossref\]](#)
19. Aletaha M, Bagheri A, Roodneshin F, Mosallaei M, Yazdani S. Oculocardiac Reflex during Strabismus Surgery: Experience from a Tertiary Hospital. *Strabismus.* 2016;24:74-78. [\[Crossref\]](#)
20. Aydın BG, Küçükosman G, Pişkin Ö, Aktaş B, Okyay RD, Uğurbaşı SC, et al. Factors affecting oculocardiac reflex incidence in pediatric strabismus surgery: Retrospective study. *JARSS.* 2021;29:58-64. [\[Crossref\]](#)
21. Aranson JK. Meyler's Side Effects of Drugs: The International Encyclopedia of Adverse Drug Reactions and Interactions. 16th edition, In: Aranson JK editor. Elsevier Science; 2016;231-235. [\[Crossref\]](#)