



ANALYZING THE PREOPERATIVE AND POSTOPERATIVE SPINOPELVIC PARAMETERS IN LENKE TYPE 3 AND TYPE 6 ADOLESCENT IDIOPATHIC SCOLIOSIS PATIENTS

Kerim SARIYILMAZ¹,
Okan ÖZKUNT¹,
Turgut AKGÜL²,
Fatih DIKICI¹,
Ünsal DOMANIÇ¹

¹ Acibadem University Atakent Hospital
Department of Orthopedics and
Traumatology

² Istanbul University Istanbul Medical
School Department of Orthopedics and
Traumatology

Address: Kerim Sarıyılmaz,
Acibadem University Atakent Hospital
Department of Orthopedics and
Traumatology, Turgut Özal Bulvarı
No:16, 34303, Halkalı, Kucukcekmece,
Istanbul.

E-mail: ksariyilmaz@gmail.com

Tel: +90 533 541 66 03

Fax: +90 212 404 44 45

Received: 22nd May, 2017.

Accepted: 14th July, 2017.

ABSTRACT

Objective: To analyze the pre and postoperative changes of sagittal spinopelvic parameters in Lenke type 3 and 6 AIS patients.

Methods: Thirty-two Lenke 3 and 6 AIS patients evaluated retrospectively. Thoracic kyphosis, lumbar lordosis, pelvic incidence, sacral slope and pelvic tilt angles were measured on preoperative and last follow-up standing full-length lateral radiographs. Kolmogorov-Smirnov test was utilized to assess distribution of study parameters. Preoperative and postoperative results were compared with Wilcoxon Sum Rank test. $p < 0.05$ considered as statistically significant.

Results: There were 27 females and 5 males. Mean age was 14 years, mean follow-up was 37 months. Mean preoperative thoracic kyphosis, lumbar lordosis, pelvic incidence, sacral slope and pelvic tilt were $38.3^\circ \pm 13.9^\circ$, $51.9^\circ \pm 9.4^\circ$, $49^\circ \pm 8.5^\circ$, $34.5^\circ \pm 6.2^\circ$, and $14.4^\circ \pm 6.9^\circ$, respectively. Mean postoperative thoracic kyphosis, lumbar lordosis, pelvic incidence, sacral slope and pelvic tilt were $26.4^\circ \pm 6.6^\circ$, $46.1^\circ \pm 7^\circ$, $49.7^\circ \pm 8.5^\circ$, $34.1^\circ \pm 5.7^\circ$, and $15.5^\circ \pm 6.7^\circ$, respectively. Preoperative and last follow-up thoracic kyphosis and lumbar lordosis comparison showed that there is a significant difference, however there is no difference in comparison preoperative and last follow-up pelvic incidence, sacral slope and pelvic tilt.

Conclusion: Lenke type 3 and 6 AIS are double structural curves and fusing the thoracic and lumbar region may change the spinopelvic parameters thus compensatory mechanisms should not be corrupted during the surgery.

Key words: adolescent idiopathic scoliosis; Lenke type 3; Lenke type 6; sagittal balance, spinopelvic parameters

Level of Evidence: Retrospective clinical study, Level III

INTRODUCTION

Adolescent idiopathic scoliosis (AIS) is a 3-dimensional deformity of the spine with an incidence of 1-3 % and an etiology that is not well yet understood. Treatment depends on age, curve type, curve size and other several factors. Several classification systems have been proposed to better understand different curve types in AIS. In 2001, Lenke⁽⁸⁾ introduced a new classification for AIS with the aim to account for all types of curves and this classification system guides selection of fusion levels. The recommended fusion is to fuse the structural curves and leave unfused the nonstructural curves.

Recent studies have shown the importance of the sagittal spinal and pelvic alignments for the treatment of spinal pathologies⁽¹³⁾. Duval-Beaupère⁽³⁾

described the relationship between the pelvis and lumbar lordosis. They introduced the pelvic incidence (PI) angle, which is an anatomical parameter characteristic of the pelvic shape of each individual, unaffected by the orientation of the pelvis. Many authors^(1,5,7,12) showed that there are correlations between the adjacent pelvic and spinal sagittal parameters: PI and sacral slope (SS), PI and pelvic tilt (PT), PI and lumbar lordosis (LL), SS and LL and finally LL and thoracic kyphosis (TK). PI is the only parameter that is anatomically fixed and it determines all spine and pelvis balances. PI determines SS and PT. SS, which is also the distal part of the lordosis, determines LL. In turn, LL balances with TK. This upward successive influence provides the spino-pelvic balance of the upright position.

Lenke type 3 curves are double major curves with a major main thoracic and a structural thoracolumbar or lumbar curve. Type 6 curves are major thoracolumbar or lumbar and structural main thoracic curves. The importance of these two curves is the necessity of the fuse the two structural thoracic and thoracolumbar/lumbar curves ⁽⁹⁾, however this may cause decompensation of spinopelvic alignment due to restriction of mobile segments close to spinopelvic region.

Thus, we aimed to analyze the pre and postoperative changes of sagittal spinopelvic parameters in Lenke type 3 and 6 AIS patients.

MATERIAL - METHODS

A retrospective study of Lenke type 3 and 6 AIS patients treated at a single institution between 2008-2013 by a single surgeon was conducted. Inclusion criteria included: 1) a diagnosis of Lenke type 3 and 6 AIS, 2) patients treated with posterior pedicle screw only instrumentation, 3) no previous spine surgery 4) full sets of preoperative and last follow-up standing full-length AP and lateral radiographs. Patients who had previous spinal surgery, suffered from congenital deformities, hybrid constructs, anterior surgery and osteotomy were excluded. Those whose radiographs did not meet standards were also excluded in order to discard measurement error.

All surgical procedures were performed by the same attending senior spinal surgeon. Segmental pedicle screws with 6.0-mm titanium rod were used in all patients. All patients received standard posterior surgery. After facetectomies within the fusion

levels, pedicle screws were placed. The lowest instrumented vertebrae (LIV) was chosen according to the stable vertebra theory, which is the first-touched vertebra by the central sacral vertical line and also lateral side-bending radiographs prior to surgery. Several surgical maneuvers were used in combination, including rod-rotation, apical vertebral derotation, convex compression, and concave distraction.

The Surgimap software (New York, New York, USA) was used to measure the sagittal spinal and pelvic parameters. Standing full-length lateral radiographs before surgery and at the last follow-up measured by the author who did not attend the surgeries. In the lateral standing radiographs, five sagittal parameters were measured: thoracic kyphosis (TK), lumbar lordosis (LL), pelvic incidence (PI), sacral slope (SS), and pelvic tilt (PT). The LL was measured as the Cobb angle between the upper endplate of the L1 and S1. The TK was measured as the Cobb angle between the upper endplate of the T5 and T12. The PI, PT, and SS were measured as the angle between the vertical line of the sacral plate and the line connecting the midpoint of the sacral plate to the mid-point of the bilateral femoral head center, the angle between the plumb line and the line connecting the midpoint of the sacral plate with the midpoint of the bilateral femoral head center, and the angle between the sacral plate and the horizontal line, respectively. Measurement values were analyzed with SPSS software (version 15.0, SPSS Inc., Chicago, IL, USA). Kolmogorov-Smirnov test was utilized to assess distribution of study parameters. Preoperative and postoperative results were compared with Wilcoxon Sum Rank test. $p < 0.05$ considered as statistically significant.




Figure-1. Preoperative and postoperative photos of a 16 year-old Lenke type 3 patient

RESULTS

A total of 218 AIS patients in the database were assessed and finally 32 patients (27 females, 5 males) with mean ages 14.7 ± 2.36 were included in the study who met all the criteria. Twenty-two patients were Lenke type 3 and 10 patients were Lenke type 6. The mean follow-up was 37.3 ± 7.2 months. Lower instrumented vertebra (LIV) was L3 in twenty-three patients and L4 in 9 patients. Mean preoperative thoracic kyphosis, lumbar lordosis, pelvic incidence, sacral slope and pelvic tilt were $38.3^\circ \pm 13.9^\circ$, $51.9^\circ \pm 9.4^\circ$, $49^\circ \pm 8.5^\circ$, $34.5^\circ \pm 6.2^\circ$, and $14.4^\circ \pm 6.9^\circ$, respectively. Mean postoperative thoracic kyphosis, lumbar lordosis, pelvic incidence, sacral slope and pelvic tilt were $26.4^\circ \pm 6.6^\circ$, $46.1^\circ \pm 7^\circ$, $49.7^\circ \pm 8.5^\circ$, $34.1^\circ \pm 5.7^\circ$, and $15.5^\circ \pm 6.7^\circ$, respectively.

Preoperative and last follow-up thoracic kyphosis and lumbar lordosis comparison showed that there is a significant difference ($p < 0.05$), however there is no difference in comparison preoperative and last follow-up pelvic incidence, sacral slope and pelvic tilt. ($p = 0.65$, $p = 0.81$ and $p = 0.38$).

DISCUSSION

After the introduction of the important effect of pelvic morphology in the regulation of an adequate sagittal balance by Duval-Beaupère et al.⁽³⁾, a great number of studies investigated the sagittal spinopelvic parameters in adults but little research has been devoted to examining this relationship on patients with AIS. Some authors have studied the spinal balance in adolescent idiopathic scoliosis (AIS)^(2,10), but its relation to pelvic configuration is poorly defined in the literature.

In the literature, debate continues as to deviations on the pelvic parameters between patients with AIS who underwent posterior instrumentation surgery and those of the healthy population. Although some authors stated no difference between the two populations^(6,14,16), some authors⁽¹⁰⁻¹¹⁾ found the PI to be higher in patients with AIS compared to the normal population.

Farshad et al.⁽⁴⁾ investigate the spinopelvic parameters in different types of AIS curves, and found that the spinopelvic balance was not statistically distinguishable in different Lenke curve types. They found a slight difference of spinopelvic balance only in Lenke type 5 and 6 (major curve at the thoracolumbar/lumbar region) with a pelvic incidence of 44° , sacral slope of 34° and pelvic tilt of 10° , when compared with normal population values. In our study investigating the Lenke type 3 and 6, those also have the structural lumbar curves, when compared with Farshad's study, we found a slight higher values in pelvic incidence, similar values in sacral slope and a slight higher values in pelvic tilt. However, when compared with normal population⁽¹¹⁾, pelvic incidence and sacral slope were similar but pelvic tilt was slightly higher.

There are explicit data in the literature regarding variances relevant to some spinopelvic parameters in patients with AIS following surgical management. For instance, La Maida et al. reported a statistically significant increase in pelvic tilt (PT)⁽⁶⁾. Similarly, Tanguay et al. obtained a significant relationship between lumbar lordosis (LL) and pelvic parameters below and within the fusion from the analysis of 60 patients with AIS following posterior spinal instrumentation and fusion surgery⁽¹⁴⁾. In contrast to the literature, in our data, PT following surgery showed no statistically significant difference, however, LL showed a statistically significant decrease.

Our study showed that pelvic parameters did not change significantly after the surgery. This indicates that the decrease in the thoracic kyphosis was compensated by the decrease in lumbar kyphosis and the spinopelvic compensatory mechanisms worked only in the spine and did not extend to the pelvic region. Furthermore, our results contradicts to the common opinion that AIS generally represents with thoracic hypokyphosis^(8,15). Mean thoracic kyphosis value for our patients was 38.3° and it changed to 26.4° postoperatively. While these pre-surgical values do not represent hypokyphosis, surgery seems to be further decreasing amount of kyphosis. This we believe may be a result of compensation to correction of lumbar lordosis or simple over correction by the surgeon.

Nevertheless, there are some limitations to our study as well. First, this is a retrospective study and it lacks randomization. The measurements were done by a computer-based software and there could be some measurement errors. Finally, we did not evaluate the costs and patient reported outcome parameters, for the reason that they are out of the scope of our aim.

CONCLUSION

Sagittal spinopelvic parameters are important in treating adolescent idiopathic scoliosis and must be taken into account before and after surgery. Lenke type 3 and 6 AIS are double structural curves and fusing the thoracic and lumbar region may change the spinopelvic parameters thus compensatory mechanisms should not be corrupted during the surgery.

REFERENCES

1. Boulay C, Tardieu C, Hecquet J, Benaim C, Mouilleseaux B, Marty C, Prat-Pradal D, Legaye J, Duval-Beaupère G, Pélissier J. Sagittal alignment of spine and pelvis regulated by pelvic incidence: standard values and prediction of lordosis. *Eur Spine J* 2006; 15(4): 415-422.
2. de Jonge T, Dubousset JF, Illes T. Sagittal plane correction in idiopathic scoliosis. *Spine* 2002; 27(7): 754-760.

-
3. Duval-Beaupere G, Schmidt C, Cosson P. A Barycentremetric study of the sagittal shape of spine and pelvis: the conditions required for an economic standing position. *Ann Biomed Eng* 1992; 20(4): 451-462.
 4. Farshad M, Catanzaro S, and Schmid SL. The spinopelvic geometry in different lenke curve types of adolescent idiopathic scoliosis. *Spine Deform* 2016; 4(6): 425-431.
 5. Jackson RP, Peterson MD, McManus AC, Hales C. Compensatory spinopelvic balance over the hip axis and better reliability in measuring lordosis to the pelvic radius on standing lateral radiographs of adult volunteers and patients. *Spine* 1998. 23(16): 1750-1767.
 6. La Maida GA, Zottarelli L, Mineo GV, Misaggi B. Sagittal balance in adolescent idiopathic scoliosis: radiographic study of spino-pelvic compensation after surgery. *Eur Spine J* 2013; 22(Suppl.-6): S859-867.
 7. Legaye J, Duval-Beaupere G, Hecquet J, Marty C. Pelvic incidence: a fundamental pelvic parameter for three-dimensional regulation of spinal sagittal curves. *Eur Spine J* 1998; 7(2): 99-103.
 8. Lenke LG, Betz RR, Harms J, Bridwell KH, Clements DH, Lowe TG, Blanke K. Adolescent idiopathic scoliosis: a new classification to determine extent of spinal arthrodesis. *J Bone Joint Surg* 2001; 83-A(8): 1169-1181.
 9. Lenke LG, Betz RR, Haher TR, Lapp MA, Merola AA, Harms J, Shufflebarger HL, Multisurgeon assessment of surgical decision-making in adolescent idiopathic scoliosis: curve classification, operative approach, and fusion levels. *Spine* 2001; 26(21): 2347-2353.
 10. Mac-Thiong JM, Labelle H, Charlebois M, Huot MP, de Guise JA. Sagittal plane analysis of the spine and pelvis in adolescent idiopathic scoliosis according to the coronal curve type. *Spine* 2003; 28(13): 1404-1409.
 11. Mac-Thiong JM, Labelle H, Berthonnaud E, Betz RR, Roussouly P. Sagittal spinopelvic balance in normal children and adolescents. *Eur Spine J* 2007; 16(2): 227-234.
 12. Marty C, Boisauvert B, Descamps H, Montigny JP, Hecquet J, Legaye J, Duval-Beaupere G. The sagittal anatomy of the sacrum among young adults, infants, and spondylolisthesis patients. *Eur Spine J* 2002; 11(2): 119-125.
 13. Schwab F, Lafage V, Patel A, Farcy JP. Sagittal plane considerations and the pelvis in the adult patient. *Spine* 2009; 34(17): 1828-1833.
 14. Tanguay F, Mac-Thiong JM, de Guise JA, Labelle H. Relation between the sagittal pelvic and lumbar spine geometries following surgical correction of adolescent idiopathic scoliosis. *Eur Spine J* 2007; 16(4): 531-536.
 15. Vaz G, Roussouly P, Berthonnaud E, Dimnet J. Sagittal morphology and equilibrium of pelvis and spine. *Eur Spine J* 2002; 11(1): 80-87.
 16. Yong Q, Zhen L, Zezhang Z, Bangping Q, Feng Z, Tao W, Jun J, Xu S, Xusheng Q, Weiwei M, Weijun W. Comparison of sagittal spinopelvic alignment in Chinese adolescents with and without idiopathic thoracic scoliosis. *Spine* 2012; 37(12): E714-720.