

Relationship Between Overactive Bladder and Anxiety in Young Men Population in Turkey

Türkiye’de Genç Erkek Popülasyonda Aşırı Aktif Mesane ve Anksiyete İlişkisi

Onur Karşlı¹, Fatih Gokalp², Ahmed Ömer Halat¹, Ömür Memik¹, Bekir Voyvoda¹, Murat Ustuner¹

¹Health Science University, Kocaeli Derince Training and Research Hospital, Department of Urology, Kocaeli, Turkey

²Osmaniye State Hospital, Department of Urology, Osmaniye, Turkey

Corresponding Author Information

Onur Karşlı

onurkarşlı@yahoo.com

+90 533 636 98 23

orcid.org/0000-0002-6451-9149

19.05.2020

01.11.2020

ABSTRACT

INTRODUCTION: Overactive bladder (OAB) symptoms are known to affect quality of life. The aim of this study is to evaluate the relationship between anxiety and OAB in young male patients in Turkish society.

METHODS: Male patients aged 18-35 years, diagnosed with OAB between 2015 and 2020, were included in the study. OAB query form (OAB-V8) was filled by participants. Hamilton anxiety scale (HAM-A) was used to evaluate the anxiety levels of OAB patients. OAB-V8 scores were compared statistically before and after anxiolytic treatment of anxious patients.

RESULTS: The mean age of 87 patients was 26.28 ± 4.53 (95% CI: 25.5-27.2) and the mean OAB-V8 symptom score was 24.96 ± 5.7 (14-36). The OAB-V8 symptom score in the group without anxiety (group 1) and the group with moderate anxiety (group 2) improved significantly at 1 month compared to the pre-treatment period ($p < 0.001$). When the OAB-V8 symptom scores in the third month were analyzed, the 1st group who didn't receive anxiolytic treatment and the 2nd and 3rd group scores receiving anxiolytic treatment were 9.6 ± 1.8 , 11.2 ± 3.2 , 10.6 ± 2.8 , and significant change in all three groups compared to the 1st month was observed ($p < 0.001$).

DISCUSSION AND CONCLUSION: In patients with anxiety, the problems caused by OAB are significantly higher than normal population. In our study, HAM-A anxiety scores were found higher in patients with OAB. OAB is a common condition that can be seen in young male population and may accompany anxiety. Treatments combined with multidisciplinary approaches are highly effective in these patients.

Keywords: Overactive bladder, anxiety, male, young.

ÖZET

GİRİŞ ve AMAÇ: Aşırı aktif mesane (AAM) semptomlarının günlük hayatı etkilediği, yaşam kalitesini azalttığı, iş verimliliğini olumsuz yönde etkilediği, aile hayatı, sosyal ilişkileri ve uyku düzenini bozduğu bilinmektedir. Bu çalışmanın amacı Türk toplumunda genç erkek

hastalarda anksiyete ile AAM arasındaki ilişkiyi değerlendirmektir.

YÖNTEM ve GEREÇLER: 2015 – 2020 Tarihleri arasında AAM tanısı alan 18-35 yaş arası erkek hastalar çalışmaya dahil edildi. Tüm çalışma katılımcılarına, alt üriner sistem semptomlarını değerlendirmek amacıyla AAM sorgu formu (OAB-V8) doldurtuldu. AAM hastalarının anksiyete düzeylerini değerlendirmek için Hamilton anksiyete derecelendirme skalası (HAM-A) kullanıldı. Anksiyöz hastaların anksiyolitik tedavi öncesi ve tedavi sonrası OAB-V8 semptom skorları istatistiksel olarak karşılaştırıldı.

BULGULAR: Çalışmaya katılan 87 hastanın ortalama yaşı 26.28 ± 4.53 (%95 CI: 25.5-27.2) yıl ve hastaların OAB-V8 semptom skoru ortalama 24.96 ± 5.7 (14-36) olarak bulundu.

Anksiyetesi olmayan grup (1. grup) ile orta şiddette anksiyetesi olan grupta (2. grup) OAB-V8 semptom skorunda 1. ayda tedavi öncesi döneme göre anlamlı olarak iyileşme sağlandı ($p<0.001$). Üçüncü aydaki OAB-V8 semptom skorları incelendiğinde, anksiyolitik tedavi almayan 1. grup ve anksiyolitik tedavi alan 2. ve 3. grup skorları 9.6 ± 1.8 , 11.2 ± 3.2 , 10.6 ± 2.3 şeklinde idi ve her üç grupta da 1. aya göre anlamlı değişim gözlemlendi ($p<0.001$).

TARTIŞMA ve SONUÇ: Klinik olarak anlamlı anksiyete ve depresyonu olan hastalarda AAM ve AAM'nin yarattığı sıkıntılar, normal popülasyona göre anlamlı derecede yüksek bulunmaktadır. Bizim çalışmamızda da AAM'si olan hastaların HAM-A anksiyete skorları daha yüksek bulunmuştur. AAM, genç erkek popülasyonda görülebilen ve anksiyetenin de eşlik edebileceği yaygın bir durumdur ve bu hastalarda multidisipliner yaklaşımlarla kombine tedavilerin erken dönem yanıtları oldukça etkindir.

Anahtar Kelimeler: Aşırı aktif mesane, anksiyete, erkek, genç.

Introduction

Overactive bladder (OAB) is a condition characterized by a sudden feeling of urination, accompanied by urinary frequency and nocturia, with incontinence. The frequency of OAB in the USA is 16% for men and 17% for women (1). According to the EpiLUTS study conducted in the USA, 27% of men and 43% of women experienced symptoms of OAB at some point in their lives (2).

Studies conducted indicate that many patients with OAB also have non-urolological symptoms. It has been reported that symptoms of urination, storage and urine are significantly associated with arthritis, asthma, chronic anxiety, depression, heart disease, irritable bowel syndrome (IBS), neurological diseases and sleep disorders (3). It is also known that OAB symptoms affect daily life, decrease the quality of life, affect work efficiency negatively, disrupt family life, social relationships and sleep patterns (4). Besides all these negativities exacerbate each other, these conditions also negatively affect the mental health of the patients. However, it is still a matter of debate whether anxiety is the cause or the result of OAB (5) (6). Studies are generally directed at the fact that OAB impairs the psychosocial situation. However, the number of studies conducted in the literature that the mental state causes OAB is low. Also, most of the studies in the literature are related to female patients. The number of studies on male patients is low. The aim of this study is to evaluate the relationship between anxiety and OAB in young male patients in Turkish society.

Material and Methods

Male patients between the ages of 18-35 who were diagnosed with overactive bladder between 2015 and 2020 were included in this retrospective study. Inclusion criteria; According to the International Continence Association (ICS) 2002 criteria, it was determined that there was a diagnosis of overactive bladder with or without urinary incontinence, infection that could make a feeling of urgent urination, pelvic radiotherapy history, no neurological diseases, and not diagnosed with anxiety and / or treatment in the last 6 months. Clinical evaluation was made according to the AUA guideline with a history, physical examination, urinalysis, post-voiding residual volume. Written consent was obtained from all

patients who participated in the study. This study was approved by the Health Sciences University Kocaeli Derince SUAM Ethics Committee; 2020/52, and conducted in accordance with the provisions of the Helsinki Declaration.

Evaluation

The clinical diagnosis of OAB was made using the 2002 International Continence Association diagnostic criteria and published AUA guidelines were followed. All study participants filled the Overactive Bladder query form (OAB-V8), the Turkish version of which was validated in order to evaluate lower urinary tract symptoms (7). For the score, 11 was accepted as the threshold value and 2 additional points were added to the score for the male gender.

The patients were consulted to the psychiatry clinic for the diagnosis of anxiety. Hamilton anxiety rating scale (HAM-A) was used to evaluate the anxiety of OAB patients. HAM-A was one of the first grading scales developed to measure the severity of anxiety symptoms and is still widely used in both clinical and research fields today. The scale consists of 14 items, each defined by a series of symptoms, and measures both psychic anxiety (mental agitation and psychological distress) and somatic anxiety (physical anxiety-related complaints). Each item is scored on a scale of 0 (not available) to 4 (severe), and the total score range is 0-56. Patients diagnosed with overactive bladder and included in the study were divided into 3 groups according to Hamilton anxiety scores. Those with a score below 17 had no anxiety, 18-24 were moderate anxious and over 25 were considered severe anxious. The group without anxiety was called group 1 (mild), the group with moderate anxiety (group 2) (group) and the group with severe anxiety group 3 (severe).

Patients diagnosed with overactive bladder according to the OAB-V8 symptom score were filled with the HAM-A scale before starting anticholinergic treatment. Solifenacin was started at a dose of 5mg / day for overactive bladder treatment. OAB-V8 symptom scores were again filled in the patients who were called for control one month later. The 1 month OAB-V8 scores of the patients who were divided into groups according to anxiety severity were compared statistically. Patients who had no improvement in symptom score and had moderate and severe anxiety according to the HAM-A scale before anticholinergic treatment were consulted to the psychiatry department. OAB-V8 symptom scores were reassessed in the 3rd month of patients who were started on anxiolytic treatment appropriate to the patient and continued concomitant anticholinergic treatment by psychiatry. OAB-V8 symptom scores were compared statistically before and after anxiolytic treatment of anxious patients.

Statistical analysis

Statistical Package 23.0 (Social SPSS Statistics; New York, USA) was used for statistical analysis. Data for continuous variables are expressed as mean and standard deviation. Data for categorical variables were expressed as percentage and frequency. For continuous variables, the differences between the three groups analyzed using the Kruskal-Wallis test, and meaningful two groups were analyzed using the Dunnett test. In the OAB score analysis, it was analyzed using Friedman test and Dunn test. Statistical significance level was accepted as $p < 0.05$.

Results

Ninety five patients who applied to the outpatient clinic and whose OAB-V8 symptom score was above 11 participated in the study. Three of the patients who participated in the study were excluded from the study because they did not come for control and 5 of them left the drug due to anticholinergic side effects. The mean age of 87 patients participating in the study was 26.28 \pm 4.53 (95% CI: 25.5-27.2) years, and the patients' OAB-V8 symptom score was 24.96 \pm 5.7 (14-36). Before anticholinergic treatment, HAM-A scale was completed. Mild

anxiety was detected in 49 patients, moderate in 22 patients, severe anxiety in 16 patients and patients were divided into 3 groups accordingly. The median OAB-V8 symptom score before treatment was found as 20 (IQR 18.0-26.0), 28 (IQR 26.0-30.0), 31 (IQR 28.0-33.0) in groups 1, 2 and 3 respectively. OAB-V8 score before treatment was significantly lower in group 1 ($p < 0.001$), but there was no significant difference between groups 2 and 3 ($p = 0.22$). A significant improvement was achieved in the OAB-V8 symptom score at 1 month compared to the pre-treatment period in the group of patients without anxiety (group 1) and group with moderate anxiety (group 2) ($p < 0.001$) (Figure 1). However, in the group with severe anxiety (group 3), it was observed that there was a statistically insignificant decrease in the OAB score at 1 month ($p = 0.64$).

When the OAB-V8 symptom scores in the third month were analyzed, the 1st group who did not receive anxiolytic treatment and the 2nd and 3rd group scores receiving anxiolytic treatment were 10 (IQR 9.0-10.0), 10 (IQR 9.0-13.0), 10.5 (IQR 9.0-11.5), and the significant change was observed in all three groups at 3rd month compared to the pretreatment ($p < 0.001$). There was no statically considerable change found between 1st month and 3rd months scores ($p=0.10$). Additionally, there was no statistically significant found between 1st month and pretreatment scores in group 3 ($p=0.64$).

Discussion

In a study conducted by the world health organization, anxiety was reported to be the most common psychiatric disorder globally (8). The prevalence of lifelong anxiety disorder in the USA is around 18%. Some researchers thought that psychosomatic disorders such as anxiety may play a role in the pathogenesis of OAB (9), and this has been shown in clinical-based and epidemiological studies (10). In our study, a relationship was found between OAB and anxiety. The relationship between affective disorders and lower urinary tract symptoms was first described by Engel in 1964. The link between the brain and bladder has not been clearly clarified today, but it has been reported that it may be using the same neuropharmacological pathways according to the hypotheses (10) (11). It is unclear which OAB and anxiety that are thought to be related trigger the other.

In a study by Milsom et al., OAB and OAB in participants with HADS (HADS) -anxiety and HADS-depression scores and who showed clinically significant anxiety and depression levels. The problems caused by the patients were found to be significantly higher than those with low anxiety and depression scores ($p < 0.001$) (12). In logistic regression analysis, the strongest ailment markers associated with OAB for both men and women were determined as urine urgency, urinary incontinence, urinary frequency and nocturia, respectively. A significant increase in OAB-related disorders was observed in male participants as the level of education, HADS-anxiety score and general health problems increased. In our study, as the level of anxiety increases in male patients, the OAB-V8 symptom score increases and the treatment of OAB becomes difficult. With anxiolytic treatment given in addition to anticholinergic treatment, a significant improvement in OAB-V8 symptom score is observed in the 3rd month. In the EpiLUTS study, in the interviews with patients with OAB symptoms, the reason for the anxiety of the patients was stated as the concern of finding a toilet in time and the urinary incontinence. At the same time, constant anxiety of going to the toilet in some participants led to a feeling of hopelessness and depression in patients (12).

In the literature, Lai et al. In a study he conducted and compared the psychological scales, OAB patients with common systemic symptoms had worse depressive symptoms (HADS-D), anxiety symptoms (HADS-A), fatigue symptoms (PROMIS-Fatigue) and higher than OAB patients without common systemic symptoms. They detected psychological stress (PSS) (13).

Results were statistically significant for all comparisons adjusted for age and gender ($p < 0.05$). Similar to the literature, HAM-A scores were higher in patients with OAB with urinary incontinence in our study.

In a study of Alves et al., Hopkins investigated the relationship of OAB with anxiety symptoms using three items of the Symptom Checklist, and although the cause-effect relationship could not be established, the authors concluded that individuals with OAB are more prone to anxiety symptoms (14). Similar to these results, although we showed the relationship between anxiety and OAB in our study, we could not show the cause-effect relationship between the two disorders.

Conclusion

OAB is a common condition that can also be seen in the young male population and may accompany anxiety. Questioning the psychosomatic complaints of patients and treating them with a multidisciplinary approach may provide better results.

References

1. Stewart WF, Van Rooyen JB, Cundiff GW. Prevalence and burden of overactive bladder in the United States. *World J Urol.* 2003. 20:327-336.
2. Coyne KS, Sexton CC, Vats V. National community prevalence of overactive bladder in the United States stratified by sex and age. *Urology.* 2011. 77:1081-1087.
3. Coyne KS, Kaplan SA, Chapple CR. Risk factors and comorbid conditions associated with lower urinary tract symptoms: EpiLUTS. *BJU Int.* 2009. 103: 24.
4. Coyne KS, Sexton CC, Irwin DE. The impact of overactive bladder, incontinence and other lower urinary tract symptoms on quality of life, work productivity, sexuality and emotional well-being in men and women: results from the EPIC study. *BJU Int.* 2008. 101:1388-1395.
5. Bogner HR, Gallo JJ, Swartz KL, Ford DE. Anxiety disorders and disability secondary to urinary incontinence among adults over age 50. *Int J Psychiatry Med.* 2002. 32(2):141-54.
6. Felde G, Bjelland I, Hunskaar S. Anxiety and depression associated with incontinence in middle-aged women: a large Norwegian cross-sectional study. *Int Urogynecol J.* 2012. 23(3):299-306.
7. Tarcan T, Mangır N, Özgür MÖ, Akbal C. OAB-V8 Aşırı Aktif Mesane Sorgulama Formu Validasyon Çalışması. *Üroloji Bülteni.* 2012. 21, 113-116.
8. Wang PS, Lane M, Olfson M, Pincus HA, Wells KB, Kessler RC. Twelve-month use of mental health services in the United States: results from the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005. 62:629-40.
9. Freeman RM, McPherson FM, Baxby K. Psychological features of women with idiopathic detrusor instability. *Urol Int.* 1985. 40(5):257-9.
10. Zorn BH, Montgomery H, Pieper K. Urinary incontinence and depression. *J Urol.* 1999. 162: 82.
11. Klausner AP, Steers WD. Corticotropin releasing factor: a mediator of emotional influences on bladder function. *J Urol.* 2004. 172: 2570.
12. Milsom I, Kaplan SA, Coyne KS, Sexton CC, Kopp ZS. Effect of Bothering Overactive Bladder Symptoms on Health-related Quality of Life, Anxiety, Depression, and Treatment Seeking in the United States: Results From EpiLUTS. *Urology.* 2012. 80: 90-96.
13. Lai HH, Vetter J, Jain S, Andriole GL. Systemic Nonurological Symptoms in Patients with Overactive Bladder. *J Urol.* 2016. 196(2):467-72.
14. Alves AT, Jácomo RH, Gomide LB, Garcia PA, Bontempo AP, Karnikoskwi MG. Relationship between anxiety and overactive bladder syndrome in older women. *Rev Bras Ginecol Obstet.* 2014. 36(7):310-4.

Table 1. OAB-V8 Scores of anxiety groups.

	HAM-A Score Groups			
	Mild	Moderate	Severe	p
OAB Score pretreatment	20 (18.0-26.0)	28 (26.0-30.0)	31 (28.0- 33.0)	p<0.001
OAB Score 1 st month	11 (9.0 -15.0)	23 (21.0-26.0)	30 (27.0-30.5)	p<0.001
OAB Score 3 rd month	10 (9.0-10.0)	10 (9.0-13.0)	10.5 (9.0-11.5)	p=0.16

*All data was expressed as median + interquartile range

OAB: over active bladder

Figure 1. OAB-V8 Score changes of groups after treatment.

