

Cough Syncope in the Times of COVID-19

© Sadaf Sheikh

Department of Emergency Medicine, Sultan Qaboos University Hospital, Muscat, Oman

Dear Editor,

In the times of Coronavirus disease-2019 (COVID-19) pandemic, cough is a common clinical presentation, however its vasomotor effects on circulatory systems are underappreciated. A healthy 39-year-old gentleman with no comorbidities presented to the emergency room during the night shift with few episodes of syncope during bouts of coughing over a week. His father suffered from upper respiratory tract symptoms for a week and recovered. He was a non-smoker and denies any use of recreational drugs. There was no fever, shortness of breath, chest pain or chronic cough. He suffered from a ground-level fall at home while coughing with brief duration of loss of consciousness for few seconds and sustained 2 centimeters linear laceration over the occipital region. His physical examination, electrocardiogram, chest X-ray and laboratory investigations were within normal limits. Prior to this presentation, he had never experienced any syncopal episode. He was negative for COVID-19 polymerase chain reaction. On further investigation, computed tomography head, echocardiogram and holter monitoring were unremarkable. His laceration was repaired and later discharged with cough suppressants and oral antibiotics to treat presumed upper respiratory tract infection.

Cough syncope or laryngeal ictus (from Latin ictus means stroke or thrust) or laryngeal vertigo coined in 1876 by Charcot, is a phenomenon with episodic cough with loss of consciousness for few seconds followed by full recovery (1). In mid-19th century, the profile of cough-induced syncope emerged in middle aged muscular built males with underlying obstructive lung disease. Presumably, such individuals generate very high intra-thoracic pressures with coughing associated with low venous return and cardiac output, decreased cerebral perfusion, increased extravascular pressure around cranial vessels resulting in cerebral concussion like effect from rapid rise in cerebrospinal

fluid pressure. It is a neutrally mediated reflex vasodepressor-bradycardia response to coughing (2,3). This case scenario suggest cough induced syncope can be a clinical problem. Reduced cerebral flow due to vagal modulation seems as a primary cause. Syncopal episode is a result of cough however elimination of cough is important hence evaluation and treatment of potential underlying causes of cough is required. The lack of robust research is due to dearth of large data sets and it is needed to investigate several pathophysiological processes. Some literatures further put forward interesting questions about vulnerable individuals with raised intra-thoracic pressures and their predisposition for this condition and suggested to incorporate new diagnostic modalities (3-5).

Keywords: Cough, syncope, COVID-19

Ethics

Peer-review: Externally peer-reviewed.

Financial Disclosure: The author declared that this study received no financial support.

References

1. Whitty CWM. On the so called "laryngeal epilepsy". Brain. 1943;66:43-54.
2. Roberto M, Patricia T, Arunashis S. Diagnostic role of head-up tilt test in patients with cough syncope. Europace. 2016;18:1273-9.
3. Chung CP. Jugular venous reflux and plasma endothelin-1 are associated with cough syncope: a case control pilot study. BMC Neurol. 2013;13:9.
4. Sutton R. Reflex syncope: Diagnosis and treatment. J Arrhythm. 2017;33:545-52.
5. Rook A.F. Coughing and unconsciousness: the so-called laryngeal epilepsy. Brain. 1946;69:138-48.



Corresponding Author: Sadaf Sheikh, M.D., Department of Emergency Medicine, Sultan Qaboos University Hospital, Muscat, Oman
E-mail: sheikh.sadaf@gmail.com ORCID ID: orcid.org/0000-0001-7457-0012

Cite this article as: Sheikh S. Cough Syncope in the Times of COVID-19. Eurasian J Emerg Med. 2022;21(1):76-76.

Received: 20.05.2020
Accepted: 23.06.2020